

fizjoterapia polska



POLISH JOURNAL OF PHYSIOTHERAPY

OFICJALNE PISMO POLSKIEGO TOWARZYSTWA FIZJOTERAPII

THE OFFICIAL JOURNAL OF THE POLISH SOCIETY OF PHYSIOTHERAPY

NR 5/2023 (23) KWARTALNIK ISSN 1642-0136



**Wpływ stresu prenatalnego
na rozwój motoryczny
niemowląt**

**Effects of prenatal stress
on infant motor
development**

Urazowość u osób biegających rekreacyjnie
Injury prevalence in recreational runners

ZAMÓW PRENUMERATĘ!
SUBSCRIBE!

www.fizjoterapiapolska.pl
www.djstudio.shop.pl
prenumerata@fizjoterapiapolska.pl





MATIO sp. z o.o.

to sprawdzony od 7 lat dystrybutor
urządzeń do drenażu dróg oddechowych
amerykańskiej firmy Hillrom

Hill-Rom.

The
Vest
Airway Clearance System

model 205



MetaNeb™



**sprzęt medyczny do drenażu i nebulizacji dla pacjentów w warunkach szpitalnych
– ze sprzętu w Polsce korzysta wiele oddziałów rehabilitacji i OIOM**

MATIO sp. z o.o., ul. Celna 6, 30-507 Kraków, tel./fax (+4812) 296 41 47,
tel. kom. 511 832 040, e-mail: matio_med@mukowiscydoza.pl, www.matio-med.pl

NOWOŚĆ W OFERCIE

ASTAR.

Tecaris



SKUTECZNA I BEZPIECZNA TERAPIA PRĄDEM O CZĘSTOTLIWOŚCI RADIOWEJ

Urządzenie przeznaczone do przeprowadzania profesjonalnych zabiegów prądem o częstotliwości radiowej (terapia TECAR).



Dowiedz się więcej
terapiatecar.astar.pl



Aparat umożliwia pracę z elektrodami rezystancyjnymi (o średnicy 25, 40, 55 lub 70 mm), pojemnościowymi (o średnicy 25, 40, 55 lub 70 mm) oraz z elektrodą typu IASTM do terapii tkanek miękkich

Tecaris generuje sinusoidalny prąd zmienny o częstotliwościach 300, 500, 750 lub 1000 kHz, dostarczanego do tkanek pacjenta za pomocą uniwersalnego aplikatora kąowego lub prostego.



Prąd o częstotliwości radiowej wywołuje efekty w głębszych warstwach tkanek, czyli kościach, ścięgnach lub więzadłach. Umożliwia to leczenie zwichnięć i zwyrodnień tkanek w przewlekłych stanach chorobowych.



Terapia wpływa przede wszystkim na tkanki powierzchniowe, czyli mięśnie (rozluźnienie) i układ limfatyczny, przyspieszając regenerację komórek.

ul. Świt 33
43-382 Bielsko-Biała

t +48 33 829 24 40
astarmed@astar.eu

POLSKI PRODUKT **WYBIERASZ I WSPIERASZ**

wsparcie merytoryczne
www.fizjotechnologia.com

www.astar.pl

Series of exercise therapy in order to make the medial longitudinal arch in changing flat foot conditions, systematic literature review

Seria terapii ruchowej w celu zmiany przyśrodkowego łuku podłużnego w schorzeniach płaskostopia, przegląd literatury systematycznej

**Ridwan Abdul Rachman^(A,B,C,D,E,F), Wawan Sundawan Suherman^(A,B,C,D,E,F),
Budi Calamita Sandra^(A,B,C,D,E,F), Ahmad Nasrulloh^(A,B,C,D,E,F)**

¹Faculty of Sport and Health Science, Universitas Negeri Yogyakarta, Indonesia

Abstract

Flat foot is a condition in which the foot has no or less arch in the arches longitudinal medial that cause the footprints become flatter than usual or seem to appear have no arch. Arches longitudinal medial is one of the pedis arches which plays an important role in the support and balance of the foot when walking. If this part becomes flat and being ignored, it will cause the limitation of walking activities, muscle imbalance, the changes of body alignment, and it can cause a risk of injury to someone as well. The aim of this paper is to identify whether the series of exercise therapy in order to make the Medial Longitudinal Arch in changing Flat Foot Conditions is an appropriate exercise method that can be carried out. This research is using the systematic literature review method with preferred reporting items for systematic reviews and meta-analytic. The data that is used in the article's findings in Sinta accredited journals is Google Scholar & Pubmed. The articles searched were based on the keywords, in which flat foot exercise therapy and changes in arches longitudinal medial, then being filtered based on the inclusion and exclusion criteria. It was 6 out of 20 articles were used as the main source in this study. The first finding of this research shows that a series of strengthening exercises such as tower curl exercise, heel raises exercise, short foot exercise, strengthening Ball Roll Exercise, calf stretch, and towel stretch can provide a curving effect or additional of the arches longitudinal medial on the flat foot. The second finding of this research is that providing training with the addition of kinesio taping can improve postural control, increase proprioception, and increase static stability in the intrinsic muscles of the foot.

Keywords

exercise therapy, flat foot, arches longitudinal medial, systematic literature review

Streszczenie

Płaskostopie to stan, w którym stopa nie ma żadnego łuku lub ma go mniej w łuku podłużnym przyśrodkowym, co powoduje, że ślady stają się bardziej płaskie niż zwykle lub wydają się nie mieć łuku. Łuk podłużny przyśrodkowy to jeden z łuków stopy, który odgrywa ważną rolę w podparciu i równowadze stopy podczas chodzenia. Jeśli ta część stania się płaska i ignorowana, spowoduje to ograniczenie aktywności chodu, brak równowagi mięśniowej, zmiany w ułożeniu ciała, a także może spowodować ryzyko kontuzji. Celem pracy jest ustalenie, czy seria ćwiczeń ruchowych mających na celu wywołanie zmian przyśrodkowego łuku podłużnego w warunkach płaskostopia jest odpowiednią metodą ćwiczeń, którą można przeprowadzić. W badaniu tym zastosowano metodę systematycznego przeglądu literatury z preferowanymi pozycjami raportowania dla przeglądów systematycznych i metaanalitycznych. Dane wykorzystane w wynikach artykułu w czasopismach akredytowanych przez Sinta pochodzą z Google Scholar i Pubmed. Wyszukiwane artykuły opierały się na słowach kluczowych, w których terapia płaskostopia oraz zmiany w łuku podłużnym przyśrodkowym, a następnie filtrowano je w oparciu o kryteria włączenia i wyłączenia. Jako główne źródło w tym badaniu wykorzystano 6 z 20 artykułów. Pierwsze wyniki tych badań pokazują, że seria ćwiczeń wzmacniających, takich jak ćwiczenia z uginaniem wieży, ćwiczenia unoszenia pięty, ćwiczenia krótkich stóp, wzmacniające ćwiczenia z piłką, rozciąganie łydek i rozciąganie rękami, mogą zapewnić efekt zakrzywienia lub dodatkowego podłużnego przyśrodkowego łuku na płaskiej stopie. Drugim wnioskiem płynącym z tych badań jest to, że zapewnienie treningu z dodatkiem kinesiotapingu może poprawić kontrolę postawy, zwiększyć propriocepcję i zwiększyć stabilność statyczną wewnętrznych mięśni stopy.

Słowa kluczowe

terapia ruchowa, płaskostopie, podłużne łuki śródstopia, systematyczny przegląd literatury

Introduction

The foot is a part of the body that has a function to support the body weight and maintain the body balance when standing. The foot has two primary functions, namely the body foundation (base of support) and the lever to move forward the body while walking or running [1]. If the support is not firm, it is not impossible that the body will often fall and end up damaging the body structure as a whole.

One of the important parts of the foot is arches. Houghlum & Bertoti said that arches are divided into three parts, namely arches longitudinal medial, arches longitudinal lateral, and arches transversal [2]. Arches serve several functions. First, they can help the foot adapt to various land surfaces. Next, they can be the power absorption that is given to the foot and base of support as well. Then, they give strength and balance when walking and disseminate the mass of the body equally on the foot. Lastly, arches keep the energy while running [3].

Arches longitudinal medial is an arch that has the highest arch composed of calcaneus, talus, navicular, cuneiform, first metatarsal, second metatarsal, and third metatarsal [4]. This arch is powered and stabilized by ligaments and intrinsic muscles. The ligaments that play a role in powering and stabilizing are the calcaneonavicular and deltoid ligaments, and the intrinsic muscles of the foot used to power and stabilize are the abductor hallucis, flexor digitorum brevis, and quadratus plantae [5]. As the arches longitudinal medial lowers or even has no arch, it may cause flat foot or it is known as pes planus. Arches longitudinal medial is one of the pedis arches which plays an important role in the support and balance of the foot when walking [6]. In the condition of flat foot, the ability of push off while walking becomes less effective due to the weakening of the plantar flexor muscle [7].

Flat foot can occur in one or both feet. This disorder has a special characteristic, namely hyperpronation of the feet [8]. Flat foot can be caused by genetic factors (congenital), the ligament weakness of the foot, or the tension in the extrinsic muscle of the ankle joint [2]. Arinaa said that the impacts that can occur due to flat foot are lessening the speed and agility while walking, getting tired easily, and changing the biomechanics of the body that can cause imbalancing [9].

The research of Nakhostin-Roohi et al. in 2013 shows that 50 teenage girls aged 14-17 with flat foot have bad static and dynamic balance [10]. Balance is needed when doing some activities such as walking, running, and standing. If someone has low balance, it will make it easy to fall and cause disruption while walking [11]. Arches that grow abnormally evoke balancing disorders, sustainability deformity, muscle weakness that make people with flat foot easy to get injured and painful [12]. Flat foot is divided into two types: flexible flat foot and rigid flat foot. Flexible flat foot is mostly due to physiological reasons, so it does not need surgery [13].

However, rigid flat foot is caused by bone abnormality [14]. Flat foot deformity first appears at the age of 10 years or more. At the age of 9 or 10 years, the arches are supposed to be firmed. It can also be given some treatments to prevent the deformity when they are adults; the prevalence of flat foot among children worldwide is around 20%-30%. In the year

2020, the child prevalence in Indonesia had not been discovered yet [15]. However, one of the research held in Surakarta showed that 27.55% of kids aged 6-12 have flat foot [9]. The case number of flat foot in Indonesia is still low. Yet, there is one research that showed 40% of teenagers aged 9-16 have flat foot [16]. The condition of flat foot can still develop during the growth period up to the age of 8 years. Nevertheless, the condition in some kids still needs some treatments so that it can prevent them from having growth problems [17].

The intervention that can be done in handling flat foot or flexible flat foot is exercise therapy. There are many exercises to improve the balance and strength [18]. One of them is strengthening exercise, the use of kinesiotaping, and medial arch support, and any other exercises. This form of strengthening exercise can improve muscle performance by increasing strength in connective tissue such as tendons, ligaments, and intramuscular connective tissue, increasing the body balance, making positive changes in body composition, and improving physical performance in daily activities [19]. The purpose of this literature review is to evaluate the treatments that can be done for the flat foot condition and explain the types of exercises that can be applied to help arch in the foot, specifically the medial longitudinal arch, by collecting and reviewing related articles over the past eight years. It is because the literature that discussed the way to handle or cure flat foot specifically reviewed from exercise therapy is still limited.

Methods research design

This research is a systematic literature review that is done by reviewing the results of previous research which discussed the series of exercise therapy to make the medial longitudinal arch in changing flat foot conditions. A systematic literature review study is a research design that uses related secondary data to a specific topic [20]. The aim of this literature review is to summarize previous research findings.

Discussion regarding the series of exercise therapy to change the medial longitudinal arch in flat foot is deemed very necessary since it considers that whether this case is a small matter but it may have fatal consequences if it was not being noticed or treated further. So the type of exercise therapy used is one of the components as well as the supporting factors to the success of physiological changes in the medial longitudinal arch in children or adults who experienced flat foot.

The data used in the article's findings are Google Scholar & Pubmed. The inclusion criteria in the study were selecting articles related to exercise therapy for flat foot. The published articles are Bali Medical Journal, Respati Jogjakarta University, Journal of nr 2/2023 [23], Physical Education, Health and Sport, FisioMU, and Kemenkes Surakarta which are accredited by Sinta. Meanwhile, the exclusion criteria in the study were articles that are not accredited by Sinta and some articles that cannot be downloaded in full text.

Result

The data collection resulted in information from research articles, including articles published in Indonesian or English Language, full text articles, articles that discuss exercise therapy in order to make the medial longitudinal arch in changing flat fo-

ot conditions, and the impact included or results from the application of exercise therapy to the condition of flat foot. The collection of the data taken from articles in the last five years, which is from 2018-2023. The first findings show the flow of data collection articles which is presented in Figure 1. Then in Table 1, it explains the analysis review of related articles about the course of exercise therapy in order to make the medial longitudinal arch in changing flat foot conditions.

Figure 1. Flow of Data Collection Explanation of the analysis results improve proprioception [29]. Using kinesio taping can also increase stimulation of mechanoreceptors that help

strengthen the posterior tibial muscle, activate sensory receptors and neuromuscular function, so that it can improve static and dynamic balance [30]. In flat foot conditions, applying kinesio taping can correct hyperpronation, so that it can improve the medial longitudinal arch [27].

One regularly with varying sets or repetitions aimed at forming the curve in the arches of the foot and strengthen the foot. By increasing muscle strength and flexibility, there will be some additional arches or an increase in the degree of arch. Therefore, it will provide good results in improving and changing the medial longitudinal arch in children who have flat foot.




Figure 1. Flow of data Collection Explanation of the analysis results

Discussion

Based on Table 1, from 6 articles that have been reviewed and analyzed, it obtained 4 discussions about exercise therapy in order to make the medial longitudinal arch in changing flat foot conditions and it will be discussed as follows:

Strengthening Exercise

Strengthening exercises, including Tower Curl Exercise, Short Foot Exercise, Heel Raises Exercises, Calf Stretch, Towel

Stretch, and Strengthening Ball Roll Exercise can help in making the medial longitudinal arch in flat foot conditions become curved [21]. This is because these exercises aim to activate the muscles, which focuses on movements involving the core muscles, especially the intrinsic parts of the foot or metatarsals [22].

Short Foot Exercise, Towel Curl Exercise, Towel Stretch are sensorimotor training that can improve the functionality of the ankle by strengthening the activation of the intrinsic muscles of the foot such as the ankle, abductor digiti minimi, abductor

hallucis, and flexor digitorum [23]. When the intrinsic muscles of the foot are activated, it will cause changes that the medial longitudinal arch becomes curved [24].

In other forms of strengthening exercises, such as Heel Raises Exercises, Calf Stretch and Strengthening Ball Roll Exercise are forms of exercise that aim to stretch the plantar fascia and increase the arch of the foot [25]. Additionally, this exercise refers to the ability of muscle contractile tissue to produce tension and resultant force in the muscle which will have an impact on changes in the medial longitudinal arch. When biomechanical changes occur, this situation can increase the tension of the intrinsic muscles in the foot so that it can help in forming the medial longitudinal arch in a flat foot condition [26].

Using Kinesio Taping

Kinesiotaping is a material similar to duct tape that was created using high technology, first developed by a chiropractor from Japan named Dr. Kenzo Kase in the 1970s [27]. This tape is made from special, highly elastic materials like cotton

with an acrylic adhesive backing. Kinesio taping has a different mechanism than ordinary foot bandages [28]. Apart from supporting muscle performance, kinesio taping also functions as a natural therapeutic tool. The function of kinesio taping is; [1] Corrects muscle function by strengthening weakened muscles. [2] Improve blood and lymph circulation by eliminating bleeding under the skin by moving the muscles. [3] Relieve pain through nerve compression. [4] Reposition subluxated joints by eliminating abnormal muscle tension, helps to restore fascia and muscle function [7]. An important function of most kinesio taping is to provide support during movement. Some kinesio taping aims to reduce injury and improve proprioception [29]. Using kinesio taping can also increase stimulation of mechanoreceptors that help strengthening the posterior tibial muscle, activates sensory receptors and neuromuscular function, so that it can improve static and dynamic balance [30]. In flat foot conditions, applying kinesio taping can correct hyperpronation, so that it can improve the medial longitudinal arch [27].

Table 1. Explanation of the analysis results

Researcher	Objective	Methods	Discussion
Putri Hapsari Srirahayuningsih & Fitri Yani 2020	To find out whether there is an effect of increasing dynamic balance in flat foot condition by giving towel curl exercise and short foot exercise	Narrative Review	Providing intervention strengthening exercise, namely with towel curl exercise and short foot exercise regularly can form curves in the middle longitudinal arch in flat foot conditions
Herta Meisatama et al., 2022	To determine intervention the effectiveness of heel raises exercises and towel curl exercises on stork stand test scores in the cases of flat foot	Experiment and quantitative analysis	This research shows that strengthening exercises such as heel raises exercises and towel curl exercises, both of them can provide strengthening of the intrinsic muscles of the foot which refers to muscle contractile tissue to produce tension and resultant muscles which will have an impact in the medial longitudinal arch of the flat foot and can help children to improve static balance of the body
Nanang Indardi, 2018	To determine the effect of strengthening exercises with kinesio taping on balance in flat foot condition	Pre and Post-test group	Providing strengthening exercise intervention with three types of exercises, namely: calf stretch, towel stretch, and towel curl which combined with using kinesiotaping can provide the effect to make change the Medial Longitudinal Arch in Flat Foot Conditions
Imam Haryoko, 2023	To determine the effect of Strengthening Ball Roll Exercise, Kinesio Taping and Towel Curl Exercise on movement disorders and ankle joint function due to flat foot	Case Study	The interventions provided by strengthening exercise are Strengthening Ball Roll Exercise and Towel Curl Exercise combined with kinesio taping. The results obtained are increased arches caused by strengthening exercise modalities which can increase the arch of the foot and The plantar fascia muscle becomes stretched and can improve the functional activities of the human body
Amirah Zahidah et al., 2020	To determine the effect of short foot exercise with using kinesio taping. in order to make the Medial Longitudinal Arch in changing flat foot Conditions	Eksperimen with randomized control trial design	Providing short foot exercises with the use of kinesio taping can influence in the medial longitudinal arch to a flat foot condition changes, increase proprioception, increase static stability in the intrinsic muscles of the foot
Rayhan Dhafa Ali & Melya Rossa., 2022	To find out the management of Exercise Therapy for Flat Foot conditions	Assessment, diagnosis, anamnesis, and intervention	Providing exercise therapy such as strengthening exercise with Heel Raises Exercises & Toe Curl Exercises involving stabilizer muscles as the shape of the arches

Medial Longitudinal Arch Change in Flat Foot Conditions

Problems in the medial longitudinal arch will cause foot abnormalities, namely flat foot. Firstly, flat foot is a condition where the sole of the foot does not have an inner arch [31]. Flat feet are caused by the bones' curve becoming flatter than usual. This case can happen because there are usually injuries in the leg or the presence of balance disorders caused by trauma or the changing of the body's spine [32]. Secondly, flat foot is considered as pes planus, which means a condition where the arch of the foot is flat, the entire sole of the foot is attached to the ground when walking or standing in a static position [33]. Lastly, factors that influence the formation of arches are genetic, complication, gender, hormones, metabolic disorders, and mechanical stress [34].

Strengthening training modalities like exercises can be applied daily with a regular training program and the body will have a physiological response and the muscles become flexible so that the medial longitudinal arch in flat foot become high [23].

Improved Muscle Strength and Balance

Doing stretching exercises will increase muscle strength and capillary blood circulation which triggers additional recruitment of motor units in the muscles so that the muscles activate the Golgi tendon organ, then the muscles will work more optimally and generate good stability [5]. In this case, the balance of flexible flat foot patients will have an increase in strength that can improve static, dynamic and postural control [23].

Strengthening exercises can also help in increasing the strength of the intrinsic muscles of the legs and around the pelvis, thighs and knees because the impacts of biomechanical changes result in muscle imbalance and cause effects on the nerves and skeleton from proprioceptive stimulation to maintain a balanced position [5, 35].

Strengthening exercises can make significant improvements in the muscles trained as well as static and dynamic balance and postural control. By having an increase in balance that has been facilitated by the twitch speed of the motor unit, the

muscle contractions stimulate gamma efferent activity in the muscle spindle. The results of strengthening exercise show an increase in strength and improvement in static and dynamic balance as well as postural control.

Conclusion

Based on the results of the explanation, it can be concluded that providing a series of exercise therapy interventions with strengthening exercises therapy such as doing towel curl exercises, heel raises exercises, short foot exercises, strengthening ball roll exercises, calf stretches, and towel stretches can give an impact toward the changes of the medial longitudinal arch in flat foot conditions. This is based on the fact that these exercises can provide stimulation to the foot and can help in making the curve in the medial longitudinal arch of the foot for flat foot condition. Basically, these movements directly provide functional strengthening and muscle flexibility in the legs as a whole. Apart from providing muscle strengthening, this exercise can also improve the arch of the foot and stretch the plantar fascia muscle. Apart from doing exercise therapy intervention, the use of kinesio taping on the foot combined with exercise therapy can improve postural control, increase proprioception, increase static stability in the intrinsic muscles of the foot. In flat foot conditions, applying kinesio taping can correct hyperpronation, so that it can repair a flat medial longitudinal arch [37]. These exercises can be done regularly with varying sets or repetitions aimed at forming the curve in the arches of the foot and strengthen the foot. By increasing muscle strength and flexibility, there will be some additional arches or an increase in the degree of arch. Therefore, it will provide good results in improving and changing the medial longitudinal arch in children who have flat foot.

Adres do korespondencji / Corresponding author

Ridwan Abdul Rachman

E-mail: ridwan.abdrchmn@gmail.com

Piśmiennictwo/ References

1. Derby H, Conner NO, Talukder A, Griffith A, Freeman C, Burch R, et al. Impact of Sub-Clinical and Clinical Compression Socks on Postural Stability Tasks among Individuals with Ankle Instability. *Healthcare (Switzerland)*. 2022 Jul 1;10(7).
2. Wicaksono A, Kusumaningtyas S, Tulaar AB. Foot Arch and Plantar Pressure in the Age of 17-21 Years. *Mutiara Medika: Jurnal Kedokteran dan Kesehatan*. 2021 Jul 8;21(2):124–9.
3. Pes Planus (Flat Feet).
4. France RC. Introduction to sports medicine and athletic training. Delmar Cengage Learning; 2011. 695 p.
5. Namsawang J, Eungpinichpong W, Vichiansiri R, Rattanathongkom S. Effects of the short foot exercise with neuromuscular electrical stimulation on navicular height in flexible flatfoot in thailand: A randomized controlled trial. *Journal of Preventive Medicine and Public Health*. 2019 Jul 14;52(4):250–7.
6. Ezema CI, Abaraogu UO, Okafor GO. Flat foot and associated factors among primary school children: A cross-sectional study. *Hong Kong Physiotherapy Journal*. 2014;32(1):13–20.
7. Halseth T, Mcchesney JW, Debeliso M, Vaughn R, Lien J. THE EFFECTS OF KINESIO TM TAPING ON PROPRIOCEPTION AT THE ANKLE [Internet]. Vol. 3, ©Journal of Sports Science and Medicine. 2004. Available from: <http://www.jssm.org>
8. Kim EK, Kim JS. The effects of short foot exercises and arch support insoles on improvement in the medial longitudinal arch and dynamic balance of flexible flatfoot patients.

9. Haq AN, Rizki Imania D. Pengaruh Short Foot Exercisedan Tibialis Posterior Strengthening Terhadap Keseimbangan Dinamis Pada Penderita Flat Foot Remaja Narrative Review [Internet]. Vol. 1, Journal Physical Therapy UNISA) ISSN XXXX-XXXX. 2021. Available from: <https://ejournal.unisayogya.ac.id/ejournal/index.php/JITU/index>
10. Nakhostin-Roohi B, Hedayati S, Aghayari A. The effect of flexible flat-footedness on selected physical fitness factors in female students aged 14 to 17 years. *Journal of Human Sport and Exercise*. 2013;8(3 SUPPL):788–96.
11. Antara KA, Adiputra N, Sugiritama W. PADA ANAK SEKOLAH DASAR NEGERI 4 TONJA KOTA DENPASAR.
12. 13. HUMAIRAH SAHABUDDIN HUBUNGAN ANTARAFAT FOOTDENGAN.
13. Studi DIII Fisioterapi P, Ilmu Kesehatan F, Muhammadiyah Palembang Ik. Penerapan Strengthening Ball Roll Exercise, Kinesiotaping dan Towel Curl Exercise Pada Gangguan Gerak dan Fungsi Sendi Pergelangan Kaki Akibat Flat Foot: Study Case Report Imam Haryoko. 2022(1):64–9.
14. Lee MS, Vanore J V., Thomas JL, Catanzariti AR, Kogler G, Kravitz SR, et al. Diagnosis and treatment of adult flatfoot. *Journal of Foot and Ankle Surgery*. 2005;44(2):78–113.
15. Rohmah Mutnawasitoh A, Pamungkas Adi Surya M, Sri Widayati R, Linia Romadhoni D, Nur Ramadhani A, Mirawati D, et al. The Effect of Otago Home Exercise Programme on Decreasing the Risk of Falling in the Elderly The Effect of Back Massage on Decreasing Insomnia in the Elderly at the Posyandu Lansia Marsudi Waras Jebres Surakarta Prevention Children Sexual Abuse in Preschool with Picture Story Book GASTER JOURNAL OF HEALTH SCIENCE The Effect of Strengthening Ball Roll Exercise and Strengthening Heel Raises Exercise on Static Balance in Children with Flat Foot in Sragi Subdistrict. 2021;19(1). Available from: <https://jurnal.aiska-university.ac.id/index.php/gasterhttps://doi.org/10.30787/gaster.v19i2.571>
16. PROFIL KESEHATAN INDONESIA TAHUN 2017.
17. Octavius GS, Sugiarto T, Agung FH, Natasha R. Flat foot at 5 to 6-year-old and history of delayed walking. *Paediatrica Indonesiana(Paediatrica Indonesiana)*. 2020;60(6):321–7.
18. Golchini A, Rahnama N, Lotfi-Foroushani M. Effect of systematic corrective exercises on the static and dynamic balance of patients with pronation distortion syndrome: A randomized controlled clinical trial study. *Int J Prev Med*. 2021 Jan 1;12(1).
19. 21. Kisner Therapeutic-exercise.-Foundations-and-techniques-by-Colby-Lynn-Allen-Kisner-Carolyn-z-lib.org__2.
20. Snyder H. Literature review as a research methodology: An overview and guidelines. *J Bus Res*. 2019 Nov 1;104:333–9.
21. 23. Differences in the Effect of Giving Heel Raises Exercise and Tighrope Walker on Improving Static Balance in Flat Foot Children. Publication manu.
22. Mckee PO, Hertel J, Bramble D, Davis I. The foot core system: a new paradigm for understanding intrinsic foot muscle function. Available from: <http://bjsm.bmj.com/>
23. Ita Mahendrayani L, Putu Gede Purwa Samatra D, Wayan Tianing N, Nyoman Ayu Dewi N. *Sport and Fitness Journal*. Vol. 6. 2018.
24. Gooding TM, Feger MA, Hart JM, Hertel J. Intrinsic foot muscle activation during specific exercises: A T2 time magnetic resonance imaging study. *J Athl Train*. 2016 Aug 1;51(8):644–50.
25. Yulianti A, Yunia Harlin Indarti B, Aulia Rahmadanti R, Ainun Ma'rufa S, Imanurrohman Lubis Z. The Effect of Strengthening Ball Roll Exercise on the Static Balance and Flexibility in Children with Flatfoot at 4-6 Years. *KnE Medicine*. 2023 Jun 23;
26. Banwell HA, Paris ME, Mackintosh S, Williams CM. Paediatric flexible flat foot: How are we measuring it and are we getting it right? A systematic review. Vol. 11, *Journal of Foot and Ankle Research*. BioMed Central Ltd.; 2018.
27. Cavalcante JGT x, Silva M do DC, Silva JT da F, dos Anjos CC, Soutinho RSR. Effect of Kinesio Taping on Hand Function in Hemiparetic Patients. *World J Neurosci*. 2018;08(02):293–302.
28. Putu Sutjana D, Irfan M. PENGGUNAAN KINESIOTAPE SELAMA TIGA HARI TIDAK BERBEDA DENGAN PEREKAT PLASEBO DALAM MENGURANGI RESIKO CEDERA BERULANG DAN DERAJAT Q-ANGLE PADA PENDERITA PATELLOFEMORAL PAIN SYNDROME. Vol. 2, *Sport and Fitness Journal*. 2014.
29. 32. Putri Hapsari Srirahayuningsih_1610301061_Fisioterapi S1 - Putry Hapsari.
30. Kim DJ, Choi IR, Lee JH. Effect of balance taping on trunk stabilizer muscles for back extensor muscle endurance: A randomized controlled study [Internet]. Available from: <http://www.ismni.org>
31. Aenumulapalli A, Kulkarni MM, Gandotra AR. Prevalence of flexible flat foot in adults: A cross-sectional study. *Journal of Clinical and Diagnostic Research*. 2017 Jun 1;11(6):AC17–20.
32. Kennedy B, Tinduh D, Utami DA, Pawana IPA, Melaniani S. Comparison of agility between the flat foot annormal foot in East Java Puslatda athletes. *Bali Medical Journal*. 2023;12(1):369–73.
33. Ariani L, Wibawa A, Made Muliarta) I. APLIKASI HEEL RAISES EXERCISE DAPAT MENINGKATAN LENGKUNGAN KAKI DAN KESEIMBANGAN STATIS PADA ANAK-ANAK FLAT FOOT USIA 4-5 TAHUN DI TK AISYIYAH BUSTANUL ATHFAL 3 DENPASAR.
34. Dabholkar T, Agarwal A. Quality of Life in Adult Population with Flat Feet [Internet]. Vol. 10, *International Journal of Health Sciences and Research (www.ijhsr.org)*. 2020. Available from: www.ijhsr.org
35. Yuvraj Babu K, Ganesh K. Assessment of Plantar Arch Index and Prevalence of Flat Feet among South Indian Adolescent Population.
36. Adegoke BO, Alumona CJ, Adeyemo AA, Adeyinka AO. Flatfoot and balance performance among junior secondary school students in Ibadan, Nigeria. *New Zealand Journal of Physiotherapy*. 2021;49(2):82–8.
37. Sivachandiran S, Kumar V. Effect of corrective exercises programme among athletes with flat feet on foot alignment factors. ~ 193 ~ *International Journal of Physical Education, Sports and Health [Internet]*. 2016;3(6):193–6. Available from: www.kheljournal.com