

fizjoterapia polska



POLISH JOURNAL OF PHYSIOTHERAPY

OFICJALNE PISMO POLSKIEGO TOWARZYSTWA FIZJOTERAPII

THE OFFICIAL JOURNAL OF THE POLISH SOCIETY OF PHYSIOTHERAPY

NR 3/2022 (22) DWUMIESIĘCZNIK ISSN 1642-

The use of deep friction massage techniques and Kinesiology Taping® (KT) in patients with faulty body posture

Zastosowanie masażu głębokiego i technik Kinesiology Tapingu® (KT) u pacjentów z wadliwą postawą ciała



Post-COVID-19 rehabilitation in the post-hospitalization period

Postępowanie rehabilitacyjne po przebytej chorobie COVID-19 w okresie poszpitalnym

ZAMÓW PRENUMERATE!

SUBSCRIBE!

www.fizjoterapiapolska.pl

www.djstudio.shop.pl

prenumerata@fizjoterapiapolska.pl



ULTRASONOGRAFIA W FIZJOTERAPII



Autoryzowani dystrybutorzy

Mar-Med

+48 22 853 14 11

info@mar-med.pl

Ado-Med

+48 32 770 68 29

adomed@adomed.pl


MAR-MED
OD 1995 ROKU

 **ADO-MED**
APARATURA MEDYCZNA



Zawód Fizjoterapeuty dobrze chroniony

Poczuj się bezpiecznie



INTER Fizjoterapeuci

Dedykowany Pakiet Ubezpieczeń

Zaufaj rozwiązaniom sprawdzonym w branży medycznej.

Wykup dedykowany pakiet ubezpieczeń INTER Fizjoterapeuci, który zapewni Ci:

- ochronę finansową na wypadek roszczeń pacjentów
— **NOWE UBEZPIECZENIE OBOWIĄZKOWE OC**
- ubezpieczenie wynajmowanego sprzętu fizjoterapeutycznego
- profesjonalną pomoc radców prawnych i zwrot kosztów obsługi prawnej
- odszkodowanie w przypadku fizycznej agresji pacjenta
- ochronę finansową związaną z naruszeniem praw pacjenta
- odszkodowanie w przypadku nieszczęśliwego wypadku

Nasza oferta była konsultowana ze stowarzyszeniami zrzeszającymi fizjoterapeutów tak, aby najskuteczniej chronić i wspierać Ciebie oraz Twoich pacjentów.

► Skontaktuj się ze swoim agentem i skorzystaj z wyjątkowej oferty!

Towarzystwo Ubezpieczeń INTER Polska S.A.
Al. Jerozolimskie 142 B
02-305 Warszawa
www.interpolska.pl

inter
UBEZPIECZENIA



zabezpiecz się przed potencjalnymi **roszczeniami** **pacjentów**

program ubezpieczeń dla fizjoterapeutów
pod patronatem PTF

dla kogo?

Zarówno dla fizjoterapeutów prowadzących własną działalność w formie praktyki zawodowej, podmiotu leczniczego jak również tych, którzy wykonują zawód wyłącznie na podstawie umowy o pracę lub umowy zlecenie.

co obejmuje program ubezpieczeń?

- igłoterapie
- zabiegi manualne (mobilizacje i manipulacje)
- leczenie osteopatyczne
- naruszenie praw pacjenta i szkody w mieniu pacjentów

oraz szereg innych rozszerzeń ukierunkowanych na zawód fizjoterapeuty



kontakt w sprawie ubezpieczeń:

Piotr Gnat

+48 663 480 698

piotr.gnat@mentor.pl

[linkedin.com/in/piotrgnat](https://www.linkedin.com/in/piotrgnat)

ubezpiecz się **on-line** na **PTFubezpieczenia.pl**



THUASNE

SpryStep®

Dynamiczne ortezy
stawu skokowego i stopy

UŁATWIA
PORUSZANIE SIĘ
Z KAŻDYM KROKIEM



© Makenzie Hollar - © Studio Caterin - © Jennifer Williams - © M-1 STUDIOS

SpryStep® flex i SpryStep® mają na celu zapewnienie wsparcia pacjentom cierpiącym z powodu opadającej stopy pochodzenia neurologicznego, pourazowego lub mięśniowego. SpryStep® plus i SpryStep® max mają na celu zapewnienie podprarcia stopy oraz dolnej kończynie pacjenta z osłabieniem dolnych kończyn pochodzenia neurologicznego, urazowego lub mięśniowego. Produkty z grupy SpryStep® są wyrobami medycznymi, posiadającymi oznakowanie CE. Przed użyciem należy zapoznać się z instrukcją oraz zasięgnąć porady specjalisty. SpryStep®, SpryStep® flex, SpryStep® plus i SpryStep® max : Thuasne Deutschland GmbH, Germany

TANITA

Healthy Habits for Happiness



JAPANESE
TECHNOLOGY

WORLD NO. 1
FOR NEW HEALTH INSIGHTS

Analizatory Składu Ciała

Mierz, monitoruj, motywuj.

Daj swoim pacjentom informacje, których potrzebują do osiągnięcia sukcesu!

Analiza składu ciała wykonywana jest w około 30 sekund, a wyniki przedstawiane są na przejrzystym raporcie.



Produkty profesjonalne TANITA wykorzystywane są przez szpitale, ośrodki badawcze, centra diagnostyczne, placówki rehabilitacyjne, kluby sportowe, osoby pracujące ze sportowcami różnych dyscyplin na całym świecie.

Więcej na tanitapolska.pl

MICROGATE



OPTOGAIT

INNOWACYJNA DIAGNOSTYKA ZDOLNOŚCI MOTORYCZNYCH I ANALIZA CHODU

Systemy MICROGATE wspierają diagnozę, ocenę postępów oraz proces rehabilitacji.

Modelowanie programów rehabilitacyjnych i kontrola procesu rehabilitacji są ułatwione dzięki obiektywnej ocenie sposobu ruchu, wykrywaniu problematycznych obszarów, ocenie biomechanicznych braków oraz ocenie asymetrii.

Możliwe parametry pomiarowe:

- fazy chodu lub biegu
- długość kroku
- prędkość i przyspieszenie
- równowaga i symetria ruchu
- video Full HD

... i wiele innych, w zależności od przeprowadzonych testów.

W połączeniu z GYKO, możliwa jest ocena stabilności dynamicznej tułowia podczas chodu/biegu, analiza skoku, analiza stabilności posturalnej, analiza zakresu ruchomości stawów (ROM), ocena siły mięśniowej.

Więcej na microgatepolska.pl

EXXENTRIC

FLYWHEEL TRAINING

Trening siłowy i rehabilitacja z użyciem zmiennej bezwładności kół zamachowych.

Exxentric wykorzystuje moment bezwładności koła zamachowego zamiast zwykłej grawitacji. To daje możliwość wykonywania ćwiczeń standardowych oraz zaawansowanych metod treningu ekscentrycznego, koncentrycznego i izometrycznego.

Jako skuteczna metoda poprawy siły i stabilności, trening ekscentryczny ułatwi pacjentom osiągnięcie zamierzonych efektów – poprawy ogólnego stanu zdrowia, wyników sportowych, rehabilitacji, czy zapobieganiu urazom.

Szeroki wybór akcesoriów i dodatków do treningu z kołem zamachowym pomoże w stworzeniu idealnego rozwiązania dla Ciebie.

Więcej na treningekscentryczny.pl



POWRÓT DO SPRAWNOŚCI PO AMPUTACJI

Czym jest program po amputacji?

Po Amputacji to unikalne w skali kraju rozwiązanie dla osób, u których konieczna była operacja odjęcia kończyn. Celem programu jest kompleksowe wsparcie w procesie odzyskania sprawności po amputacji, niezależnie od jej przyczyny. Pomagamy pokonywać granice, osiągać kolejne cele, kreować bardziej przyjazną rzeczywistość.



W ramach programu oferujemy:

- Opieka Menadżera Rehabilitacji
- Dofinansowanie do zakupu protez
- Wsparcie psychologiczne
- Bezpłatne konsultacje protetyczne
- Rehabilitacja w ośrodkach na terenie kraju
- Pomoc Asystentów Wsparcia

**Skontaktuj się z nami i zapytaj
o bezpłatne egzemplarze
Poradnika dla
osób po
amputacji
do Twojej
placówki.**



**Masz pytanie odnośnie programu.
Napisz do nas lub skontaktuj się z nami
telefonicznie:**

+48 793 003 695

biuro@poamputacji.pl www.poamputacji.pl



POAMPUTACJI
Razem możemy więcej

SPRZEDAŻ I WYPOŻYCZALNIA ZMOTORYZOWANYCH SZYN CPM ARTROMOT®

Nowoczesna rehabilitacja **CPM** stawu kolanowego, biodrowego, łokciowego, barkowego, skokowego, nadgarstka oraz stawów palców dłoni i kciuka.



ARTROMOT-H



ARTROMOT-F



ARTROSTIM
FOCUS PLUS

ARTROMOT-K1 ARTROMOT-SP3 ARTROMOT-S3 ARTROMOT-E2

Najnowsze konstrukcje ARTROMOT zapewniają ruch bierny stawów w zgodzie z koncepcją **PNF** (Proprioceptive Neuromuscular Facilitation).

KALMED Iwona Renz
ul. Wilczak 3
61-623 Poznań
www.kalmed.com.pl

tel. 61 828 06 86
faks 61 828 06 87
kom. 601 64 02 23, 601 647 877
kalmed@kalmed.com.pl



Serwis i całodobowa
pomoc techniczna:
tel. 501 483 637
service@kalmed.com.pl

mindray

healthcare within reach

ULTRASONOGRAFIA

W FIZJOTERAPII



Autoryzowani dystrybutorzy

Mar-Med

+48 22 853 14 11

info@mar-med.pl

Ado-Med

+48 32 770 68 29

adomed@adomed.pl


MAR-MED
OD 1995 ROKU

 **ADO-MED**
APARATURA MEDYCZNA

NOWOŚĆ W OFERCIE

ASTAR.

PhysioGo.Lite SONO

**NIEWIELKIE URZĄDZENIE
EFEKTYWNA TERAPIA ULTRADŹWIĘKOWA**

Zaawansowana technologia firmy Astar to gwarancja niezawodności i precyzji parametrów. Urządzenie, dzięki gotowym programom terapeutycznym, pomaga osiągać fizjoterapeucie możliwie najlepsze efekty działania fal ultradźwiękowych.

Głowica SnG to bezobrotowe akcesorium o dużej powierzchni czola (17,3 cm² lub 34,5 cm² w zależności od wybranego trybu działania). Znajduje zastosowanie w klasycznej terapii ultradźwiękami, fonoforezie, terapii LIPUS i zabiegach skojarzonych (w połączeniu z elektroterapią).



wsparcie merytoryczne
www.fizjotechnologia.com



ul. Świt 33
43-382 Bielsko-Biała

t +48 33 829 24 40
astarmed@astar.eu

**POLSKI
PRODUKT**  **WYBIERASZ
I WSPIERASZ**

www.astar.pl

Dr. Comifort®

Nowy wymiar wygody.

Obuwie profilaktyczno-zdrowotne
o atrakcyjnym wzornictwie



APROBATA
AMERYKAŃSKIEGO
MEDYCZNEGO
STOWARZYSZENIA
PODIATRYCZNEGO



WYRÓB
MEDYCZNY

**Stabilny, wzmocniony
i wyścielany zapętek**
Zapewnia silniejsze
wsparcie łuku
podłużnego stopy

**Miękki, wyścielany
kołnierz cholewki**
Minimalizuje podrażnienia

Wyścielany język
Zmniejsza tarcie
i ulepsza dopasowanie

Lekka konstrukcja
Zmniejsza codzienne
zmęczenie

**Antypoślizgowa,
wytrzymała podeszwa
o lekkiej konstrukcji**
Zwiększa przyczepność,
amortyzuje i odciąża stopy

**Wysoka jakość materiałów
- oddychające siatki i naturalne skóry**
Dostosowują się do stopy,
utrzymują je w suchości
i zapobiegają przegrzewaniu

**Zwiększona
szerokość i głębokość
w obrębie palców
i przodostopia**
Minimalizuje ucisk
i zapobiega urazom

Trzy
rozmiary
szerokości

Podwyższona
tęgłość

Zwiększona
przestrzeń
na palce

**Ochronna przestrzeń
na palce - brak szwów
w rejonie przodostopia**
Minimalizuje możliwość zranień

WSKAZANIA

- haluksy • wkładki specjalistyczne • palce młotkowate, szponiaste • cukrzyca (stopa cukrzycowa) • reumatoidalne zapalenie stawów
- bóle pięty i podeszwy stopy (zapalenie rozciągnięcia podeszwowego - ostroga piętowa) • płaskostopie (stopa poprzecznie płaska)
- bóle pleców • wysokie podbicie • praca stojąca • nerwiak Mortona • obrzęk limfatyczny • opatrunki • ortezy i bandaż • obrzęki
- modzele • protezy • odciski • urazy wpływające na ścięgna, mięśnie i kości (np. ścięgno Achillesa) • wrastające paznokcie



ul. Wilczak 3
61-623 Poznań
tel. 61 828 06 86
fax. 61 828 06 87
kom. 601 640 223, 601 647 877
e-mail: kalmed@kalmed.com.pl
www.kalmed.com.pl



www.butyladzrowia.pl

www.dr-comfort.pl


fizjoterapia polska



sklep internetowy:
www.djstudio.shop.pl

w sklepie dostępne między innymi:

- archiwalne numery Fizjoterapii Polskiej w wersji papierowej
- artykuły w wersji elektronicznej
- książki poświęcone fizjoterapii
- prenumerata Fizjoterapii Polskiej





Polisa**Med**

program
**ubezpieczeń
dla studentów**
kierunków medycznych



Drodzy Studenci

szukający artykułów do pracy naukowej.

**Przypominamy o dobrowolnym ubezpieczeniu
OC studentów kierunków medycznych!**

dlatego warto je mieć?

- ponieważ bywa wymagane w trakcie praktyk, staży czy wolontariatu
- niektóre Uczelnie wymagają je do udziału w zajęciach praktycznych
- działa na całym świecie, a dodatkowo otrzymasz certyfikat w języku angielskim w razie wyjazdu na ERASMUS-a
- wywołuje uśmiech na twarzy Pań z dziekanatów – sami sprawdziliśmy!



**posiadamy również w ofercie
ubezpieczenia dla masażystów
i techników masażystów.**



Polisa**Med**

**kontakt w sprawie
ubezpieczeń:**


+48 56 642 41 82

kontakt@polisa.med.pl

Ubezpiecz się **on-line** na

polisa.med.pl

NOWOŚCI



FUNKCYJNA BIELIZNA LECZNICZA

PRZECIWŻYLAKOWA

Przeciwwżylakowe wyroby pończosznice włoskich producentów, bardzo skuteczne i niezwykle eleganckie. Dostępne **w I, II oraz III klasie kompresji** w wielu modelach, w różnym stopniu przezroczystości (m. in. wyjątkowo przezroczyste w II kl. ucisku), w szerokiej gamie kolorystycznej, w różnych wersjach długości, z palcami zamkniętymi lub otwartymi

- podkolanówki ● pończochy ● legginsy ● rajstopy ● rękawy kompresyjne

ANTYCELLULITOWA, NA LIMFODEMIĘ I LIPODEMIĘ

Bielizna i odzież wykonana jest z mikrofibry. Unikalny splot nawet przy najmniejszym ruchu wywołuje **efekt masażu**. Dziańska stymuluje cyrkulację podskórną i drenaż limfatyczny. Prowadzi to do poprawy jakości skóry

- z włókna emana®
- z kofeiną i wit. E
- z nanosrebrem

NA NIETRZYMANIE MOCZU

Wyroby medyczne **wielokrotnego użytku** z dyskretną stałą wszywką o właściwościach chłonnych. Polecane jako codzienna bielizna gwarantująca ochronę przed przemakaniem - 100% absorpcji cieczy, zapewniająca całkowitą suchość warstw: zewnętrznej i wewnętrznej

- do wielokrotnego prania (min. 100 prań)

ART & COLL
M E D I C A L E

artcoll.pl

e-sklep@artcoll.pl

tel. 22 720 35 96

+48 510 160 100

Impact of acupuncture dry needle versus progressive pressure technique on pain severity and pressure pain threshold in patients with lower back myofascial pain syndrome: A randomized clinical trial

Wpływ suchego igłowania i progresywnego ucisku na nasilenie bólu i próg bólu uciskowego u pacjentów z zespołem bólu mięśniowo-powięziowego dolnej części pleców: randomizowane badanie kliniczne

**Ghada Mohamed Rashad Koura^{1,2(A,B,C,E,F,G)}, Saud Mashi Alrawaili^{3(A,C,D,E)},
Amr Moustafa Yehia Mohammed^{4(A,B,D,E,F)}, Sara Awad Gharama Alasmri^{2,5(B,D,E,F)},
Raghad Mudawi Yahya Asiri^{2,5(B,E,F)}, Ahmed Mohamed Fathi Elshawi^{5,6(A,B,C,D,E,F)}**

¹Department of Physical Therapy for Musculoskeletal Disorders and its Surgery, Faculty of Physical Therapy, Cairo University, Egypt

²Department of Medical Rehabilitation, College of Applied Medical Sciences, King Khalid University, Saudi Arabia

³Department of Physical Therapy and Health Rehabilitation, College of Applied Medical Sciences, Prince Sattam bin Abdulaziz University, Saudi Arabia

⁴Department of Musculoskeletal Disorders and its Surgery, Faculty of Physical Therapy, 6 October University, Giza, Egypt

⁵Saudi German Hospital, Aseer, Saudi Arabia

⁶Faculty of Physical Therapy, Cairo University, Egypt

Abstract

Background. Chronic lower back pain contributes to the global disability burden, and continues to be a factor in the number of years spent disabled; Myofascial pain syndrome (MPS) is an inflammatory condition that may cause chronic localized pain and stiffness. Repetitive movements and incorrect posturing habits contribute to MPS advancement by formation of trigger points (TrPs) on the muscles.

Purpose. To investigate the impact of acupuncture dry needle versus progressive pressure technique on pain severity and pressure pain threshold (PPT) in patients with lower back MPS.

Methods. A randomized clinical trial. Thirty patients suffering from lower back MPS with the presence of TrPs on iliocostalis lumborum, quadratus lumborum, piriformis, and gluteus medius muscles were divided into two equal groups. Group (A) received acupuncture dry needle over TrPs followed by stretching exercise for 2 weeks (n = 15), and group (B) received progressive pressure technique over the same TrPs followed by stretching exercise for 2 weeks (n = 15). Pain severity and PPT at the TrPs were evaluated before and after treatment.

Results. There were statistically significant improvements in pain severity and PPT at the TrPs of the four muscles within both groups (p < 0.05), except the PPT at iliocostalis lumborum that showed a non-significant difference within group (A) (p > 0.05). Comparing both groups pre- and post-treatment showed statistically non-significant differences in all outcome measures (p > 0.05).

Conclusion. Both acupuncture dry needle and progressive pressure technique were effective in reducing pain severity and increasing PPT in patients with lower back MPS.

Keywords

acupuncture dry needle, progressive pressure technique, pain severity, pressure pain threshold, myofascial pain syndrome

Streszczenie

Informacje wprowadzające. Przewlekły ból dolnej części pleców przyczynia się do ogólnego obciążenia niepełnosprawnością i jest czynnikiem wpływającym na liczbę lat spędzonych z niepełnosprawnością; Zespół bólu mięśniowo-powięziowego (MPS) to stan zapalny, który może powodować przewlekły ból i sztywność. Powtarzające się ruchy i nieprawidłowe nawyki postawy przyczyniają się do postępu zespołu MPS poprzez powstawanie punktów spustowych (TrPs) na mięśniach.

Cel. Zbadanie wpływu suchego igłowania i progresywnego ucisku na nasilenie bólu i próg bólu uciskowego (PPT) u pacjentów z zespołem MPS w dolnej części pleców.

Metody. Randomizowane badanie kliniczne. Trzydziestu pacjentów cierpiących na MPS dolnej części pleców z wyczuwalnymi punktami spustowymi na mięśni biodrowo-lędźwiowym, czworobocznym lędźwi, gruszkowatym i pośladkowym podzielono na dwie równe grupy. Grupa (A) była poddawana suchemu igłowaniu punktów spustowych, a następnie wykonywała ćwiczenia rozciągające przez 2 tygodnie (n=15), a grupa (B) była poddawana progresywnemu uciskowi tych samych punktów spustowych, a następnie wykonywała ćwiczenia rozciągające przez 2 tygodnie (n=15). Nasilenie bólu i PPT w punktach spustowych oceniano przed i po leczeniu.

Wyniki. Wystąpiła statystycznie istotna poprawa w zakresie nasilenia bólu i PPT w punktach spustowych czterech mięśni w obu grupach (p < 0,05), z wyjątkiem PPT na mięśniu biodrowo- lędźwiowym, który wykazał nieistotną różnicę w grupie (A) (p > 0,05). Porównanie obu grup przed i po leczeniu wykazało statystycznie nieistotne różnice we wszystkich pomiarach wyników (p > 0,05).

Wniosek. Zarówno suche igłowanie jak i progresywny ucisk były skuteczne w zmniejszaniu nasilenia bólu i zwiększaniu PPT u pacjentów z MPS w dolnej części pleców.

Słowa kluczowe

suche igłowanie, progresywny ucisk, nasilenie bólu, próg bólu uciskowego, zespół bólu mięśniowo-powięziowego

Introduction

Chronic pain is a major cause of morbidity and has a significant impact on quality of life [1]. Chronic lower back pain (LBP) contributes to the global disability burden and continues to be a factor in the number of years spent disabled [2, 3]. From a global standpoint, it is also regarded as a global public health problem [4].

Myofascial pain syndrome (MPS) is an inflammatory condition that causes recurrent chronic pain in muscles and fascia or associated soft tissues. It can be accompanied by obvious emotional disorders or dysfunction [5-7]. The muscular fascia is a connective tissue in which pathological changes appear under overload. That damage leads to the biomechanical change of muscle contraction force and flexibility [8]. Repetitive movements and incorrect posturing habits contribute to the advancement of MPS by the formation of TrPs on participating muscles [9].

A trigger point is a hypersensitive spot in a taut dense band of skeletal muscle. When stimulated, it causes referred pain and motor phenomena [10]. Depending on their clinical characteristics, TrPs are classified as active or latent. An active TrP causes spontaneous pain enhanced with palpation. The pain pattern is similar to the pain complaint of the patient. Unprompted pain that only appears in the presence of a local stimulus seems to be a latent TrP. It can, however, have repercussions such as limited range of motion (ROM) and muscle weakness [11]. Clinical symptoms related to myofascial TrPs (pain, limitation of ROM, paresthesia, and weakness) occur due to the shortening of muscle fibers and pressure on close-up nerves and adjacent tissues [12]. These symptoms may have a large impact on life quality, health status, and mood [13]. Quadratus lumborum (QL) is one of the many significant causes of LBP [14]. Poor posture and body positioning change the body mechanics that appear in muscle trigger points (MTrPs). There are four potential TrPs in the QL muscle, including upper, lower, middle, or deep. QL MTrPs were seen to be more common in the younger age group, with the highest prevalence, i.e., 48%, in patients 20–30 years of age [12].

The new secondary type of LBP is gluteal syndrome (GS), which was included in the 11th Revision of the International Classification. GS is a good example of nociplastic pain that can imitate sciatica or overlap with real sciatica. It is characterized as a local syndrome of the lower limbs that enhances myofascial pain from one of the three TrPs in gluteal muscles (i.e., gluteus maximus, medius, or minimus) [15].

The most important differential diagnosis of LBP is piriformis syndrome, and it should not be ruled out with proper examination and assessment. Clinicians must consider medical management and conservative management in the initial treatment plan. Conservative management includes physical therapy, steroid injections, botulinum toxin injections, and dry needling, which are all potentially effective [16].

Different therapeutic methods are advocated for TrPs management [17], Kinesio tape and dry needling have more positive effects in terms of pain and disability [18], with needling interventions being the most commonly used [17]. Extracorporeal shock wave therapy [19] and manual therapy interventions are effective for improving ROM in cases with

myofascial TrPs. Massage, cupping, and ischemic compression are the most common conventional approaches for treating TrPs [20, 21]. Neuromusculoskeletal pain and movement impairments can be managed with needles that penetrate the skin and stimulate TrPs and muscular tissues [22]. Dry needling utilization by physical therapists in the United States has increased over the last five years. It has been certified by over 4,000 physiotherapists in the United States [23].

The TrPs progressive pressure technique is effective and safe for successfully reducing pain and tenderness [24]. The clinician applies direct digital pressure on each TrP until a state of relief tension is reached and inactivates the TrP [25]. As demonstrated by Holey and Cook [26], stretching complements myofascial TrP pressure release by effectively deactivating a TrP. Stretching is a beneficial intervention to treat chronic MPS. Stretching cervical muscles for 30 seconds was optimal in reducing the negative effects of MPS on the affected nerve roots and central nervous system [27].

Until now, no study compared the effects of acupuncture dry needle versus progressive pressure technique on pain severity and pressure pain threshold in patients with lower back MPS. Thus, this study was conducted to investigate which treatment protocol is more effective.

Subjects and methods

Study Design

The study was designed as an experimental randomized clinical trial. It based on the Helsinki Declaration principles for human research. It was carried out between October 2017 and March 2022.

Participants

Thirty patients (male & female), with an age ranging from 18 to 43 years, were recruited from the out clinic of the Faculty of Physical Therapy, Cairo University, Egypt. They had lower back MPS for 3 months, and MTrPs in iliocostalis lumborum, quadratus lumborum, piriformis, and gluteus medius muscles on any side, with a pain level of 3 cm to 7 cm on visual analog scale (VAS). The diagnosis of MPS was confirmed through both proper history taking and clinical examination. The history identified the areas affected by pain and helped in finding the predisposing factors that led to MPS. In clinical examination, muscles whose TrPs could refer pain in the affected areas were examined; muscles were palpated searching for taut bands, using either flat palpation or pincer palpation; fingers were moved along the taut band to find the hardest and most tender spot (TrP); TrPs were compressed manually and the patient was asked if the spot was tender or painful, and if so the pain resembled his/her usual pain; TrPs were compressed for 5–10 seconds and the patient was asked if there was pain radiating from the TrP. Exclusion criteria were history of previous back surgery, neurologic deficit, current lower extremity symptoms, a cardiovascular disease with decreased activity tolerance, rheumatologic conditions, polyarticular osteoarthritis, rheumatoid arthritis, and advanced lumbar degenerative disease, as well as participants taking other treatment, whether physical therapy or medication that might interfere with the results of our study.




Figure 1. Participant flow diagram

Randomization

Each participant was notified of the nature, purpose, and value of the study, as well as the right to withdraw from it at any time and the confidentiality of all data acquired. After all baseline criteria had been met, the random permuted block method was the way to randomize the participants. They were randomized in our study to ensure that the treatment groups were balanced at the end of every block. Participants were registered in blocks using random number generator software (figure 1). All patients were asked to sign a consent form for ethical issues.

Interventions

Group (A) consisted of 15 patients receiving acupuncture dry needle over TrPs of iliocostalis lumborum, quadratus lumborum, gluteus medius, and piriformis muscles followed by stretching exercise for 2 weeks. Group (B) consisted of 15 patients receiving progressive pressure technique over the same muscles followed by stretching exercise for 2 weeks.

Acupuncture dry needle

Each participant in group (A) received acupuncture dry needle over TrPs in iliocostalis lumborum, quadratus lumborum, gluteus medius, and piriformis muscles, 2 treatment sessions per week, for 2 weeks. Before the application of acupuncture dry needle, a detailed explanation of the procedure to the patient was performed. Possible fear of needling and pain associated with dry needle were in our consideration. The pain associated with needling could be inhibited by activating patients' conditioned pain modula-

tion system, as demonstrated by research [28]. Initially, the TrPs were determined by palpating taut bands in the belly of the iliocostalis lumborum, quadratus lumborum, gluteus medius, and piriformis muscles. Then, dry needling was performed with a solid filiform needle in a tube, 0.30 mm in diameter, 50 and 70 mm long. The filiform needle in its tube was steady with the non-needling hand against the targeted area by using a pincer grip or flat palpation depending on the muscle orientation, location, and direction of needle penetration. In order to draw the needle out, the needling hand gently pulled it from the tube. In order for the needle to penetrate the skin, the top of the needle was tapped or flicked. To achieve deep needling, the needle was directed in the direction of the TrP. Once the needle felt resistance against the TrP, a local twitch response (LTR) was provoked and that was considered fundamental in obtaining a desirable therapeutic effect [29, 30]. The needle was inserted deep enough to fully penetrate the taut band region and then pulled back to the subcutaneous tissue layer but not out of the skin, then the needle was kept in the TrP for seven to ten minutes [31]. Generally, numerous LTRs could be elicited. Discontinuation of a given dry needling procedure might happen as a result of prominent decreased frequency or removal of LTRs, decreased resistance to palpation of the underlying tissue, or patient lack of tolerance of continued needling at that site. Once the needle had come out completely from the skin, pressure (hemostasis) was applied directly to the skin over the insertion site of the needle with an alcohol swab to prevent possible swelling or soreness. The muscle was reassessed by palpation of taut bands and TrPs [32].

Regarding iliocostalis lumborum, TrP was identified via flat palpation from a side-lying position. A needle of 5cm in length was inserted slightly superior to the TrP, perpendicular to the skin, and directed anteromedially for about 30°. Care was taken to avoid penetration of the lung. Regarding quadratus lumborum, the patient was placed in a lying position, with the arm in extension to elevate the rib cage and the leg in extension and adduction to lower the iliac crest. Using a pillow under the sound side to make the space wider where TrPs were identified was easier. Caudal to the 12th rib and anterior to the paraspinal muscle mass, the 7-cm needle was inserted in a parallel direction to the back's plane (in the frontal plane) toward the transverse processes of L2 and L3. Regarding gluteus medius, the patient was in a side-lying position. A needle of 5cm in length was used along the contour of the iliac crest. The muscle was needled perpendicularly with flat palpation. The strong dropping of the subcutaneous tissue was required to minimize the distance from the skin to the muscle. It was common if the needle contacted the periosteum. Precautions were taken to avoid needling the nerves (sciatica- superior gluteal) and deep branch of the superior gluteal vessels. The amount of adipose tissue determined the depth of penetration. Regarding piriformis, the patient was in a side-lying position. The bony landmarks for the greater trochanter and sacrum at S2, S3, and S4 were detected. Then, the needle 5 cm length was inserted perpendicular to the muscle surface directly into the TrP taut band from the sciatic notch toward the pubic symphysis, with taking care to avoid nerve needling [31].

Progressive pressure technique

Each participant in group (B) received progressive pressure technique over TrPs in iliocostalis lumborum, quadratus lumborum, gluteus medius, and piriformis muscles, 2 treatment sessions per week, for 2 weeks. The patient position was side lying for the application of progressive pressure technique over the TrPs in the four muscles. A sustained gentle pressure was applied using the thumb or four fingers for 90-120 seconds moving inward toward the center of the MTrp. Once tissue resistance was felt, pressure was maintained until resistance dissipated (melting away). This cycle was repeated 3 times with a 90-second resting period between each compression. By the end of treatment, relaxation of the tissue was felt or no benefit was achieved [33].

Stretching exercise

Groups (A) received stretching exercise for iliocostalis lumborum, quadratus lumborum, gluteus medius, and piriformis muscles following acupuncture dry needle, while group (B) received the same stretching exercise following progressive pressure technique, 2 sessions per week, for 2 weeks. Concerning iliocostalis lumborum stretching, it was achieved through reaching with the arm to the opposite side from long sitting and trunk flexion. Concerning quadratus lumborum stretching, the patient was in a semi-prone position with one leg in adduction and extension, and the therapist supported the lower thoracic cage and iliac crest while spreading the hands. The therapist could also support the lower thoracic cage and

iliac crest with his hands while spreading the hands apart in semi-supine with the leg in adduction and flexion. Concerning gluteus medius stretching, the patient was supine with the treated side in hip flexion and adduction. The patient eased movement using one hand to help hip flexion and the other to assist hip adduction. Concerning piriformis stretching, the patient was supine with the treated side in hip flexion above 90 degrees, adduction, and external rotation. Assurance was on external rotation. The patient eased movement using the hands and the other leg to help hip flexion, adduction and external rotation [34, 35].

Outcome measures

Pain severity assessment

For pain assessment, the VAS was used to evaluate pain severity pre- and post-treatment for both groups (A) and (B). The VAS is a valid, responsive, and frequently used pain outcome measure [36]. The validated subjective measure of chronic and acute pain is the visual analog scale (VAS). The scores were recorded by putting a handwritten mark on a 10-cm line between "no pain" and "worst pain" [37].

Pressure pain threshold (PPT) measurement

The PPT at the TrPs of the iliocostalis lumborum, quadratus lumborum, gluteus medius, and piriformis muscles were measured pre- and post-treatment for both groups (A) and (B), using a pressure algometer (Baseline® Pressure algometer Fabrication Enterprises, Inc. PO Box 1500 White Plains, NY 10602). The pressure algometer has a high degree of validity and reliability in measuring PPT for the diagnosis of MPS in back and trunk muscles [38]. The tip of the algometer was placed precisely on the MTrP region to avoid any difference in readings, the pressure was increased by 1 kg/cm² steady and perpendicular to the identified TrPs [39]. Patients were instructed to say "now" when they first began to feel a change of pressure to either discomfort or pain [40]. Pressure thresholds lower than 3 kg were considered to be abnormally low [41].

Statistical analysis

Statistical Package for Social Sciences (SPSS) for Windows, version 20 (SPSS, Inc., Chicago, IL) was used to analyze the data. Prior to final analysis, data were screened for normality assumption, homogeneity of variance, and presence of extreme scores. This exploration was done as a pre-requisite for parametric calculations of the analysis of difference. Results were expressed as mean and standard deviation for baseline and demographic data as well as PPT at the TrPs of the iliocostalis lumborum, quadratus lumborum, gluteus medius, and piriformis muscles, while they were expressed as median for pain severity. In normally distributed data, unpaired t test was used for comparison of different variables between groups, while paired t test was used for comparing "pre" and "post" treatment within the same group for different variables. In non-normally distributed data, unpaired Mann-Whitney test was used for comparison of different variables between groups, while Wilcoxon test was used for comparing "pre" and "post" treatment within the same group. The alpha level was set at 0.05.

Results

There were no statistically significant differences ($p > 0.05$) between subjects in both groups concerning age, body mass, height and all outcome measures at baseline (Tables 1–3).

The pain severity showed a statistically significant reduction ($p < 0.05$) within the two groups (A) and (B). The post-treatment comparison of both groups showed a statistically non-significant difference ($p > 0.05$) (Table 2).

The PPT showed statistically significant increases at the TrPs of the four muscles ($p < 0.05$) within both groups, except the PPT at iliocostalis lumborum that showed a non-significant difference within group (A) ($p > 0.05$). The post-treatment comparison of both groups showed statistically non-significant differences at the TrPs of the four muscles ($p > 0.05$) (Table 3).

Table 1. Physical characteristics of patients in both groups

	Group (A) (n = 15)	Group (B) (n = 15)	P value
Age [years]	29 ± 6.17	25.26 ± 6.30	0.113 ^{NS}
Body mass [kg]	84.8 ± 11.43	79.2 ± 10.94	0.181 ^{NS}
Height [cm]	174.06 ± 8.48	172.36 ± 8.39	0.585 ^{NS}

NS, $p > 0.05$ = non-significant, p = probability

Table 2. Pain severity for both groups

		Group (A) (n = 15)	Group (B) (n = 15)	P value*
Pain severity	Pre-treatment	6	6	0.964 ^{NS}
	Post-treatment	3	4	0.549 ^{NS}
	P value**	0.001 ^S	0.001 ^S	

Data were expressed as median

* Inter-group comparison; ** intra-group comparison of the results pre- and post-treatment

NS, $p > 0.05$ = non-significant; S, $p < 0.05$ = significant; p = probability

Table 3. The PPT at the TrPs of iliocostalis lumborum, quadratus lumborum, gluteus medius, and piriformis muscles for both groups

		Group (A) (n = 15)	Group (B) (n = 15)	P value*
PPT at iliocostalis lumborum [kg/cm ²]	Pre-treatment	3.58 ± 1.68	3.14 ± 1.37	0.521 ^{NS}
	Post-treatment	4.76 ± 1.58	4.94 ± 1.63	0.804 ^{NS}
	Improvement%	32.96%	57.32%	
	p value**	0.071 ^{NS}	0.006 ^S	
PPT at quadratus lumborum [kg/cm ²]	Pre-treatment	2.47 ± 0.75	3.14 ± 1.37	0.866 ^{NS}
	Post-treatment	4.35 ± 2.3	4.94 ± 1.63	0.687 ^{NS}
	Improvement%	76.11%	57.32%	
	p value**	0.004 ^S	0.006 ^S	
PPT at gluteus medius [kg/cm ²]	Pre-treatment	3.59 ± 1.13	3.09 ± 0.88	0.253 ^{NS}
	Post-treatment	5.39 ± 2.42	4.8 ± 1.97	0.531 ^{NS}
	Improvement%	50.1%	55.3%	
	p value**	0.029 ^S	0.012 ^S	
PPT at piriformis [kg/cm ²]	Pre-treatment	3.15 ± 1.4	2.68 ± 1.2	0.387 ^{NS}
	Post-treatment	5.56 ± 2.15	4.13 ± 1.9	0.086 ^{NS}
	Improvement%	76.5%	54.1%	
	p value**	0.002 ^S	0.011 ^S	

Data were expressed as mean and standard deviation

* Inter-group comparison; ** intra-group comparison of the results pre- and post-treatment

Discussion

To our knowledge, this is the first randomized trial demonstrating the impact of acupuncture dry needling versus progressive pressure technique on pain severity and PPT in patients with lower back MPS.

In the present study, we observed that both acupuncture dry needle and progressive pressure technique followed by stretching exercise were effective in reducing pain severity and increasing PPT in patients with lower back MPS and resulted in similar outcomes after treatment.

Regarding the positive effect of acupuncture dry needle followed by stretching exercise on reducing pain severity and increasing PPT, the results of this study are consistent with a systematic review by Hu et al. [42] that revealed the effects of dry needling for reducing pain and disability for treating LBP. Another meta-analysis included 45 articles by Sánchez-Infante et al. [43] analyzed the effect of dry needling in the short, medium, and long term and the results suggest that dry needling is effective in reducing pain in the short (immediately to 72 hours), medium (1–12 weeks), and long (13–24 weeks) terms. In addition to the effectiveness of dry needling in reducing pain better than other modalities such as transcutaneous electrical nerve stimulation, compression technique, superficial dry needling, conventional rehabilitation, massage, stretching.

The dry needling chemical effect in some studies has demonstrated that enhanced levels of bradykinin, calcitonin gene-related peptide, substance P, and other TrP chemicals are directly corrected by eliciting LTR after dry needling [44].

Regarding the beneficial influence of progressive pressure technique followed by stretching exercise on lowering pain severity and improving PPT, the findings of the current study come in agreement with other studies showing the effects of trigger point release in patients with MPS. Kostopoulos et al. [45] compared the efficacy of ischemic compression, passive stretching, and the combination of ischemic compression and passive stretching for the first time and reported that the combination was the most effective for pain relief. In addition, Lake et al. [46] investigated the efficacy of ischemic compression on 13 patients with 40 myofascial TrPs and observed that ischemic compression was significantly efficient for treatment in comparison with the control group, but the optimal level of ischemic compression was not defined. Moreover, Hanten et al. [47] experimented with the efficacy of the combination of ischemic compression and stretching for patients with MPS on the neck and upper back for 5 days and then the duration of pain sensations in 24 hours. PPT and VAS measured 3 days after treatment were compared with those

measured for pre-treatment. Results revealed that a home program, consisting of ischemic pressure and sustained stretching, is effective in reducing TrPs sensitivity and pain intensity in neck and upper back pain. Furthermore, Hou et al. [48] examined various combinations of physiotherapy treatment methods for active upper trapezius trigger points and found that ischemic compression with specific pressure and duration provided immediate pain relief and decreased TrP sensitivity.

Shortened sarcomeres are the main cause of TrP formation and local ischemia at the site of TrPs. Therefore, by the use of progressive pressure technique and stretching in group (B), the shortened sarcomeres will flatten and lengthen. This local stretch reduces the overlapping of actin and myosin and also causes a flush of blood at the area of compression after the pressure is removed. This improves the local circulation and thus reduces the release of noxious, painful substances. All of this leads to inhibiting the TrP activity and reduces the sensitivity of myofascial TrPs [40].

The current study has some limitations. The first limitation is its short duration. As a result, future studies are needed to evaluate the long-term impact of acupuncture dry needling versus progressive pressure technique in patients with lower back MPS. Another limitation is the lack of control group. So, it would be useful for future trials to include a control or placebo group. The third limitation is the lack of the underlying mechanisms attributing to these findings. Thus, further research is needed to investigate the effect of acupuncture dry needling versus progressive pressure technique on various TrP chemicals in individuals suffering from lower back MPS. At last, the management of MTrPs is not limited to MTrP inhibition, but it requires correction of aggravation factors that lead to the activate it.

Conclusion

The results of the current study showed significant improvements in pain severity and pressure pain threshold in patients with lower back myofascial pain syndrome. No differences were found between the improvements in both groups. Acupuncture dry needle and progressive pressure technique were equally effective in reducing pain level and increasing pressure pain threshold in the treatment of patients with myofascial back pain.

Adres do korespondencji / Corresponding author

Ahmed Mohamed Fathi Elshiwi

E-mail: asd2511978@gmail.com

Acknowledgements

The authors would like to thank all individuals who participated in this study.

Piśmiennictwo/ References

1. World Health Organization. The burden of musculoskeletal conditions at the start of the new millennium: Report of a WHO scientific report. Genève, Switzerland: World Health Organization; 2003.
2. Hurwitz EL, Randhawa K, Yu H, Côté P, Haldeman S. The Global Spine Care Initiative: a summary of the global burden of low back and neck pain studies. *Eur Spine J*. 2018;27(Suppl 6):796-801.
3. James SL, Abate D, Abate KH, Abay SM, Abbafati C, Abbasi N, Abbastabar H, Abd-Allah F, Abdela J, Abdelalim A, Abdollahpour I. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *The Lancet*. 2018;392(10159):1789-1858.

4. Sowden G, Hill JC, Morso L, Louw Q, Foster NE. Advancing practice for back pain through stratified care (STarT Back). *Braz J Phys Ther.* 2018;22(4):255-264.
5. Lu Y, Cheng J, Fan B, Liu YQ, Yu SY, Zhang DY, Fu ZJ, Song XJ, Yi XB, Cheng ZX, Liu XL. ICD-11 Chinese compilation of chronic pain classification. *Zhongguo Tengtong Yixue Zazhi.* 2018;24:801-805.
6. National Center for Health Statistics 2021 release of International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM) FY 2021, Geneva, Switzerland: World Health Organization; 2020.
7. Weller JL, Comeau D, Otis JAD. Myofascial Pain. *Semin Neurol.* 2018;38(6):640-643.
8. Devereux F, O'Rourke B, Byrne PJ, Byrne D, Kinsella S. Effects of Myofascial Trigger Point Release on Power and Force Production in the Lower Limb Kinetic Chain. *J Strength Cond Res.* 2019;33(9):2453-2463.
9. Bautista A, Webb C, Rosenquist R. Serratus Anterior Muscle Pain Syndrome: A Diagnostic Conundrum. *Pain Med.* 2017;18(8):1600-1602.
10. Simons DG, Travell JS. Myofascial Pain and Dysfunction. *The Trigger Point Manual.* 3rd edition. Philadelphia: Wolters Kluwer; 2019.
11. Travell JG, Simons DG, Simons LS. Travell & Simons' myofascial pain and dysfunction: the trigger point manual. Second edition. Philadelphia: Lippincott Williams & Wilkins; 1998.
12. Shah D, Sheth M, Shah A. Effect of static versus Dynamic Ultrasound on pain in subjects with trigger point in Quadratus Lumborum having non-specific chronic low back pain. *International Journal of Health Sciences and Research* 2020; 10(12):263-269.
13. De Meulemeester KE, Castelein B, Coppieters I, Barbe T, Cools A, Cagnie B. Comparing Trigger Point Dry Needling and Manual Pressure Technique for the Management of Myofascial Neck/Shoulder Pain: A Randomized Clinical Trial. *J Manipulative Physiol Ther.* 2017;40(1):11-20.
14. Barge AS, Barge SM. Quadratus lumborum: One of the many significant causes of low back pain. *Ind J Pain* 2018;32(3):184.
15. International Association for the Study of Pain (IASP). Classification of Chronic Pain. second edition (Revised); 2021.
16. Vij N, Kiernan H, Bisht R, Singleton I, Cornett EM, Kaye AD, Imani F, Varrassi G, Pourbahri M, Viswanath O, Urits I. Surgical and Non-surgical Treatment Options for Piriformis Syndrome: A Literature Review. *Anesth Pain Med.* 2021;11(1):e112825.
17. Fernández-de-Las-Peñas C, Nijs J. Trigger point dry needling for the treatment of myofascial pain syndrome: current perspectives within a pain neuroscience paradigm. *J Pain Res.* 2019;12:1899-1911.
18. Yasar MF, Yaksi E, Kurul R, Alisik T, Seker Z. Comparison of dry needling and kinesio taping methods in the treatment of myofascial pain syndrome: A single blinded randomised controlled study. *Int J Clin Pract.* 2021;75(10):e14561.
19. Wu T, Li S, Ren J, Wang D, Ai Y. Efficacy of extracorporeal shock waves in the treatment of myofascial pain syndrome: a systematic review and meta-analysis of controlled clinical studies. *Ann Transl Med.* 2022;10(4):165.
20. Moraska AF, Schmiede SJ, Mann JD, Butryn N, Krusch JP. Responsiveness of Myofascial Trigger Points to Single and Multiple Trigger Point Release Massages: A Randomized, Placebo Controlled Trial. *Am J Phys Med Rehabil.* 2017;96(9):639-645.
21. Behrangrad S, Kamali F. Comparison of ischemic compression and lumbopelvic manipulation as trigger point therapy for patellofemoral pain syndrome in young adults: A double-blind randomized clinical trial. *J Bodyw Mov Ther.* 2017;21(3):554-564.
22. Pociask FD, Fleck TM. Iontophoresis, ultrasound, phonophoresis, and laser and light therapy. In: *Orthopaedic Physical Therapy Secrets.* Elsevier; 2017. p. 73–80.
23. Evidence in motion, home [Internet]. EIM. 2019. Available from: <https://www.kinetacore.com/about/history/>
24. Fernández-de-las-Peñas C, Alonso-Blanco C, Fernández-Carnero J, Carlos Miangolarra-Page J. The immediate effect of ischemic compression technique and transverse friction massage on tenderness of active and latent myofascial trigger points: a pilot study. *J Bodyw Mov Ther.* 2006;10(1):3–9.
25. Kisner C, Colby LA. Therapeutic exercise: foundations and techniques. FA Davis Company, Philadelphia; 2002. p. 13–24.
26. Holey EA, Cook EM. Neuromusculoskeletal disorders. Holey EA, Cook EM: Therapeutic massage. Philadelphia: WB Saunders; 1997. p. 77-100.
27. Mansoori SS, Moustafa IM, Ahbouch A, Harrison DE. Optimal duration of stretching exercise in patients with chronic myofascial pain syndrome: A randomized controlled trial. *Journal of Rehabilitation Medicine.* 2021;53(1).
28. Bjørkedal E, Flaten MA. Expectations of increased and decreased pain explain the effect of conditioned pain modulation in females. *Journal of pain research.* 2012;5:289.
29. Hong CZ. Lidocaine injection versus dry needling to myofascial trigger point. The importance of the local twitch response. *American journal of physical medicine & rehabilitation.* 1994;73(4):256-263.
30. Tekin L, Akarsu S, Durmuş O, Çakar E, Dinçer Ü, Kıralp MZ. The effect of dry needling in the treatment of myofascial pain syndrome: a randomized double-blinded placebo-controlled trial. *Clinical rheumatology.* 2013;32(3):309-315.
31. Travell JG, Simons DG. Myofascial pain and dysfunction: the trigger point manual. Lippincott Williams & Wilkins; 1983.
32. Dunning J, Butts R, Mourad F, Young I, Flannagan S, Perreault T. Dry needling: a literature review with implications for clinical practice guidelines. *Phys Ther Rev.* 2014;19(4):252-265.
33. Kostopoulos D, Rizopoulos K. The manual of trigger point and myofascial therapy. Thorofare, NJ: Slack; 2001.
34. Hanney WJ, Puenteadura EJ, Kolber MJ, Liu X, Pabian PS, Cheatham SW. The immediate effects of manual stretching and cervicothoracic junction manipulation on cervical range of motion and upper trapezius pressure pain thresholds. *Journal of back and musculoskeletal rehabilitation.* 2017;30(5):1005-1013.
35. Barbero M, Schneebeli A, Koetsier E, Maino P. Myofascial pain syndrome and trigger points: evaluation and treatment in patients with musculoskeletal pain. *Current opinion in supportive and palliative care.* 2019;13(3):270-276
36. Hjermstad MJ, Fayers PM, Haugen DF, Caraceni A, Hanks GW, Loge JH, Fainsinger R, Aass N, Kaasa S, European Palliative Care Research Collaborative (EPCRC). Studies comparing numerical rating scales, verbal rating scales, and visual analogue scales for assessment of pain intensity in adults: a systematic literature review. *Journal of pain and symptom management.* 2011;41(6):1073-1093.
37. Delgado DA, Lambert BS, Boutris N, McCulloch PC, Robbins AB, Moreno MR, et al. Validation of digital visual analog scale pain scoring with a traditional paper-based visual analog scale in adults. *J Am Acad Orthop Surg Glob Res Rev.* 2018;2(3):e088.
38. Park G, Kim CW, Park SB, Kim MJ, Jang SH. Reliability and usefulness of the pressure pain threshold measurement in patients with myofascial pain. *Ann Rehabil Med.* 2011;35(3):412-417.
39. Gemmell H, Miller P and Nordstrom H. Immediate effect of ischemic compression and trigger point pressure release on neck pain and upper trapezius trigger points: A randomized controlled trial. *Clinical Chiropractic* 2008;11(1):30-36.
40. Fryer G, Hodgson L. The effect of manual pressure release on myofascial trigger points in the upper trapezius muscle. *Journal of Bodywork and movement therapies.* 2005;9(4):248-255.
41. Fischer AA. Quantitative and objective documentation of soft tissue abnormality: pressure algometry and tissue compliance recording. *Motor vehicle collision injuries, mechanism, diagnosis, and management. Part.* 1996;2:142-148.
42. Hu HT, Gao H, Ma RJ, Zhao XF, Tian HF, Li L. Is dry needling effective for low back pain?: A systematic review and PRISMA-compliant meta-analysis. *Medicine (Baltimore).* 2018;97(26):e11225.
43. Sánchez-Infante J, Navarro-Santana MJ, Bravo-Sánchez A, Jiménez-Díaz F, Abián-Vicén J. Is dry needling applied by physical therapists effective for pain in musculoskeletal conditions? A systematic review and meta-analysis. *Physical Therapy.* 2021;101(3):pzab070.
44. Shah JP. Integrating dry needling with new concepts of myofascial pain, muscle physiology, and sensitization. In: Audette J.F., Bailey A., editors. *Integrative Pain Medicine.* Humana Press; Totowa, NJ: 2008. p. 107–121.
45. Kostopoulos D, Nelson Jr AJ, Ingber RS, Larkin RW. Reduction of spontaneous electrical activity and pain perception of trigger points in the upper trapezius muscle through trigger point compression and passive stretching. *Journal of musculoskeletal pain.* 2008;16(4):266-278.
46. Lake DA, Wright LL, Cain J, Nail R, White L. The effectiveness of ischemic pressure and ischemic pressure combined with stretch on myofascial trigger points. *J Orthop Sports Phys Ther.* 2009;39:A70.
47. Hanten WP, Olson SL, Butts NL, Nowicki AL. Effectiveness of a home program of ischemic pressure followed by sustained stretch for treatment of myofascial trigger points. *Physical therapy.* 2000;80(10):997-1003.
48. Hou CR, Tsai LC, Cheng KF, Chung KC, Hong CZ. Immediate effects of various physical therapeutic modalities on cervical myofascial pain and trigger-point sensitivity. *Archives of physical medicine and rehabilitation.* 2002;83(10):1406-1414.