

fizjoterapia polska



POLISH JOURNAL OF PHYSIOTHERAPY

OFICJALNE PISMO POLSKIEGO TOWARZYSTWA FIZJOTERAPII

THE OFFICIAL JOURNAL OF THE POLISH SOCIETY OF PHYSIOTHERAPY

NR 1/2023 (23) KWARTALNIK ISSN 1642-0136

**Restytucja wielkości cech postawy
w płaszczyźnie czołowej po obciążeniu
masą transportowanych przyborów
szkolnych w trybie ciagu lewą i prawą
ręką uczniów obojga płci w wieku 7 lat**

Restitution of the size of postural features
in the frontal plane after loading with the
weight of school items carried with the right
and left hand in 7-year-old pupils
of both sexes

Ocena postawy ciała gimnazjalistów w kontekście ich aktywności fizycznej

Assessment of body posture of junior high school students in the context of their physical activity

ZAMÓW PRENUMERATĘ!

SUBSCRIBE!

www.fizjoterapiapolska.pl

www.djstudio.shop.pl

prenumerata@fizjoterapiapolska.pl



mindray

healthcare within reach

ULTRASONOGRAFIA W FIZJOTERAPII



Autoryzowani dystrybutorzy

Mar-Med

+48 22 853 14 11

info@mar-med.pl

Ado-Med

+48 32 770 68 29

adomed@adomed.pl



MAR-MED

OD 1995 ROKU



ADO-MED®

APARATURA MEDYCZNA



Zawód Fizjoterapeuty dobrze chroniony

Poczuj się bezpiecznie



INTER Fizjoterapeuci

Dedykowany Pakiet Ubezpieczeń

Zaufaj rozwiązaniom sprawdzonym w branży medycznej.

Wykup dedykowany pakiet ubezpieczeń INTER Fizjoterapeuci, który zapewni Ci:

- ochronę finansową na wypadek roszczeń pacjentów
— **NOWE UBEZPIECZENIE OBOWIĄZKOWE OC**
- ubezpieczenie wynajmowanego sprzętu fizjoterapeutycznego
- profesjonalną pomoc radców prawnych i zwrot kosztów obsługi prawnej
- odszkodowanie w przypadku fizycznej agresji pacjenta
- ochronę finansową związaną z naruszeniem praw pacjenta
- odszkodowanie w przypadku nieszczęśliwego wypadku

Nasza oferta była konsultowana ze stowarzyszeniami zrzeszającymi fizjoterapeutów tak, aby najsuk-
teczniej chronić i wspierać Ciebie oraz Twoich pacjentów.

► Skontaktuj się ze swoim agentem i skorzystaj z wyjątkowej oferty!

Towarzystwo Ubezpieczeń INTER Polska S.A.

Al. Jerozolimskie 142 B

02-305 Warszawa

www.interpolska.pl

inter
UBEZPIECZENIA


fizjoterapia polska



sklep internetowy:
www.djstudio.shop.pl

w sklepie dostępne między innymi:

- *archiwalne numery Fizjoterapii Polskiej w wersji papierowej*
- *artykuły w wersji elektronicznej*
- *książki poświęcone fizjoterapii*
- *prenumerata Fizjoterapii Polskiej*





KALMED
Iwona Renz, Poznań

ARTROMOT®
WYŁĄCZNY PRZEDSTAWICIEL
WWW.KALMED.COM.PL



SPRZEDAŻ I WYPOŻYCZALNIA ZMOTORYZOWANYCH SZYN CPM ARTROMOT®

Nowoczesna rehabilitacja **CPM** stawu kolanowego, biodrowego, łokciowego, barkowego, skokowego, nadgarstka oraz stawów palców dłoni i kciuka.



ARTROMOT-H



ARTROMOT-F



ARTROSTIM
FOCUS PLUS

ARTROMOT-K1 ARTROMOT-SP3 ARTROMOT-S3 ARTROMOT-E2

Najnowsze konstrukcje ARTROMOT zapewniają ruch bierny stawów w zgodzie z koncepcją **PNF** (Proprioceptive Neuromuscular Facilitation).

KALMED Iwona Renz
ul. Wilczak 3
61-623 Poznań
www.kalmed.com.pl

tel. 61 828 06 86
faks 61 828 06 87
kom. 601 64 02 23, 601 647 877
kalmed@kalmed.com.pl

Serwis i całodobowa
pomoc techniczna:
tel. 501 483 637
service@kalmed.com.pl

NOWOŚĆ W OFERCIE

ASTAR.

PhysioGo.Lite SONO

NIEWIELKIE URZĄDZENIE EFEKTYWNA TERAPIA ULTRADŹWIEKOWA

Zaawansowana technologia firmy Astar to gwarancja niezawodności i precyzjności parametrów. Urządzenie, dzięki gotowym programom terapeutycznym, pomaga osiągać fizjoterapeucie możliwie najlepsze efekty działania fal ultradźwiękowych.

Głowica SnG to bezobsługowe akcesorium o dużej powierzchni czota (17,3 cm² lub 34,5 cm² w zależności od wybranego trybu działania). Znajduje zastosowanie w klasycznej terapii ultradźwiękami, fonoforezie, terapii LIPUS i zabiegach skojarzonych (w połączeniu z elektroterapią).



wsparcie merytoryczne
www.fizjotechnologia.com




ul. Świt 33
43-382 Bielsko-Biała

t +48 33 829 24 40
astarmed@astar.eu

POLSKI PRODUKT  **WYBIERASZ
I WSPIERASZ**

www.astar.pl



B.A.D. - Boundaries Are Dumb - the first Polish clothing brand dedicated primarily to all amputees but also to all open-minded people who do not recognise the limitations in their lives. In order to meet the expectations of our clients, we are creating the highest quality products with special reinforcements and systems to facilitate the removal. Stylish and comfortable - our modern, minimalist design emphasises the versatility of B.A.D. All products are made in Poland from first-class cotton because customer satisfaction is crucial to our brand. Say **“No”** to limitations and answer the question, “Are you ready to be **B.A.D.?**”



**DESIGNED FOR PEOPLE
WHO CAN'T RESPECT
BOUNDARIES**



MEN'S FABRIC TROUSERS FOR RIGHT AND LEFT-SIDED FEMORAL PROSTHESIS USERS

Comfortable fabric trousers that look ordinary on the surface. However, thanks to the **use of a zipper** placed under the flap on the outside of the left or right leg will allow you to unfasten and **freely adjust the prosthetics** without having to remove your pants

They are fitted with an elastic waistband supported by string to tie. The product is made of high-quality fabric with the addition of elastane, which improves the comfort of use.

There is a short zipper on the left leg in the inner seam at thigh level, **allowing easy access to the valve.** An additional improvement is a **reinforcement** in the area of the bend in the knee and on the back of the pants, the place that wears off more quickly from sitting.



bad_clth_
www.instagram.com/bad_clth_



BAD – Boundaries Are Dumb
www.facebook.com/BoundariesAreDumb

www.bad-clth.com






Pomoc w optymalizacji procesu rehabilitacji **chorych z COVID-19**

w różnych okresach choroby i zdrowienia
z uwzględnieniem zachowania **zasad bezpieczeństwa** związanych z potencjalnym zakażeniem terapeutów

**SPRAWDŹ
TUTAJ**



**TYLKO
W FORMIE
e-booka**



**POBIERZ
DARMOWY E-BOOK**
od PZWL w prezencie!




mindray

healthcare within reach

ULTRASONOGRAFIA

W FIZJOTERAPII



Autoryzowani dystrybutorzy

Mar-Med

+48 22 853 14 11

info@mar-med.pl

Ado-Med

+48 32 770 68 29

adomed@adomed.pl



MAR-MED

OD 1995 ROKU



ADO-MED®

APARATURA MEDYCZNA

The 2nd
International Conference On Multidiscipline Education



Kampus
Merdeka
INDONESIA JAYA



Empowering Minds: Navigating the Future of Education

Keynote Speakers



Prof. Kerry J. Kennedy

The Then Hong Kong Institute of Education
Hongkong



Prof. Madya Ts. Dr. Arumugam Raman

Universiti Utara Malaysia
Malaysia



Prof. George Arief D Liem

Nanyang Technological University
Singapore



Dr. Erick Burhaein, M.Pd., AIFO

Universitas Ma'arif Nahdlatul Ulama Kebumen
Indonesia

Sub Theme

- Blended Learning
- Curricula
- Early Childhood Education
- Educational Leadership
- Educational Psychology
- Education Programs and Teaching
- Foreign Languages Education
- Health
- Health Education
- Higher Education
- Innovative Methodologies in Learning
- Learners of Qualitative Research
- Learning Environments
- Methodology of Sport and History of Physical Culture and Sport
- Multimedia in Digital Learning
- Physical Activity and Health
- Physical Education
- Public Health
- Sport Sciences
- Teaching and Assessment
- Teaching Disability
- Virtual and Augmented Reality Learning Environments

Opening Remarks



H. Herman Suherman, S.T., M.A.P.

Regent of Cianjur
Indonesia



Deden Nasihin, S.Sos.I., M.K.P.

Deputy Chairman of DPRD
Cianjur, Indonesia



Prof. Dr. H. Dwidja Priyatno, SH., MH., Sp.N.

Rector of Suryakencana University
Cianjur



Dr. H. Munawar Rois, M.Pd.

Dean of Faculty Of Education Teacher Training
Suryakencana University Cianjur

Publication Options

- International Journal of Learning Teaching and Educational Research (Scopus Q3)
- Hong Kong Journal of Social Sciences (Scopus Q4 dan WOS)
- International Journal of Disabilitas and Health Sciences (Scopus Q4)
- Polish Journal of Physiotherapy (Scopus Q4)
- The International Society for Technology Educational and Sciences (ISTES) (Proceedings are indexed in Scopus)
- Al-Ishlah : Jurnal Pendidikan (Terindeks SINTA 2)
- Edu Sportivo: Indonesian Journal of Physical Education (Terindeks SINTA 2)
- Journal Elemen (Terindeks SINTA 2)
- Journal Sport Area (Terindeks SINTA 2)
- Journal Teori dan Aplikasi Matematika (Terindeks SINTA 2)

Conference Registration Fees



General Participants	50 K
Host Student Presenters	400 K
General Presenters	500 K



Bank Mandiri | BNI | BRI
1820006898530 | 0622468257 | 010501070965501
a.n D. Nurfajrin Ningsih

Important Dates

18 March 2023 - 30 May 2023	1 June 2023 - 20 July 2023	1 - 20 July 2023	20 July 2023 - 2 August 2023	27 July 2023 - 2 August 2023	5 August 2023	September - December 2023
Abstract Arrangement	Full Paper Acceptance	Payment Due	Full Paper Review	Announcement of Full Paper Accepted	Conference Day	Full Paper Publication

Contact Person ☎ +62 877-7879-4797 Rani Sugiarni

Organized by:

Faculty of Education Teacher Training
Suryakencana University Cianjur, Indonesia

Jl. Pasir Gede Raya, Bojongherang, Cianjur, Indonesia



Dr. Comfort®



APROBATA
AMERYKAŃSKIEGO
MEDYCZNEGO
STOWARZYSZENIA
PODIATRYCZNEGO

Nowy wymiar wygody.

Obuwie profilaktyczno-zdrowotne
o atrakcyjnym wzornictwie



WYRÓB
MEDYCZNY

**Stabilny, wzmocniony
i wyściełany zapętek**
Zapewnia silniejsze
wsparcie łuku
podłużnego stopy

**Miękki, wyściełany
kołnierz cholewki**
Minimalizuje podrażnienia

Wyściełany język
Zmniejsza tarcie
i ulepsza dopasowanie

Lekka konstrukcja
Zmniejsza codzienne
zmęczenie

**Antypoślizgowa,
wytrzymała podeszwa
o lekkiej konstrukcji**
Zwiększa przyczepność,
amortyzuje i odciąża stopy

**Wysoka jakość materiałów
- oddychające siatki i naturalne skóry**
Dostosowują się do stopy,
utrzymują je w suchości
i zapobiegają przegrzewaniu

**Zwiększona
szerokość i głębokość
w obrębie palców
i przodostopia**
Minimalizuje ucisk
i zapobiega urazom

Trzy
rozmiary
szerokości

Podwyższona
tęgłość

Zwiększona
przestrzeń
na palce

**Ochronna przestrzeń
na palce - brak szwów
w rejonie przodostopia**
Minimalizuje możliwość zranień

WSKAZANIA

- haluksy • wkładki specjalistyczne • palce młotkowate, szponiaste • cukrzyca (stopa cukrzycowa) • reumatoidalne zapalenie stawów
- bóle pięty i podeszwy stopy (zapalenie rozciągna podeszwowego - ostroga piętowa) • płaskostopie (stopa poprzecznie płaska)
- bóle pleców • wysokie podbicie • praca stojąca • nerwiak Mortona • obrzęk limfatyczny • opatrunki • ortezy i bandaże • obrzęki
- modzele • protezy • odciski • urazy wpływające na ścięgna, mięśnie i kości (np. ścięgno Achillesa) • wrastające paznokcie



ul. Wilczak 3
61-623 Poznań
tel. 61 828 06 86
fax. 61 828 06 87
kom. 601 640 223, 601 647 877
e-mail: kalmed@kalmed.com.pl
www.kalmed.com.pl



www.butydlazdrowia.pl

www.dr-comfort.pl



26. Sympozjum Sekcji Rehabilitacji Kardiologicznej i Fizjologii Wysiłku Polskiego Towarzystwa Kardiologicznego

11-13 maja 2023, Wisła, Hotel STOK

www.rehabilitacja2023ptk.pl



Rehabilitacja kardiologiczna i fizjologia wysiłku – zapraszamy do rejestracji na wyjątkową konferencję w Wiśle

W dniach 11–13 maja w Hotelu Stok Wiśle odbędzie się wyjątkowe i interdyscyplinarne spotkanie specjalistów z całej Polski – 26. Sympozjum Sekcji Rehabilitacji Kardiologicznej i Fizjologii Wysiłku Polskiego Towarzystwa Kardiologicznego. Serdecznie zapraszamy do rejestracji.

26. Sympozjum Sekcji Rehabilitacji Kardiologicznej i Fizjologii Wysiłku Polskiego Towarzystwa Kardiologicznego to coroczne spotkanie specjalistów, zajmujących się rehabilitacją kardiologiczną, prewencją chorób układu krążenia i innymi formami aktywności fizycznej, która ma prowadzić do poprawy stanu naszego zdrowia.

Ta trzydniowa konferencja przeznaczona jest dla lekarzy kardiologów, specjalistów rehabilitacji medycznej oraz innych specjalności, którzy w swojej codziennej praktyce zajmują się rehabilitacją i fizjologią wysiłku, ale także dla fizjoterapeutów, pielęgniarek, techników i przedstawicieli innych zawodów medycznych, zainteresowanych tematyką spotkania, oraz studentów.

Jakie tematy zostaną poruszone podczas konferencji?

26. Sympozjum Sekcji Rehabilitacji Kardiologicznej i Fizjologii Wysiłku to konferencja, na którą zaproszeni zostali wybitni specjaliści z dziedziny kardiologii i nie tylko. Podczas wydarzenia wygłoszonych zostanie prawie 100 wykładów merytorycznych w ciągu aż 20 sesji. Uczestnicy będą mieli również szansę na udział w sesjach przypadków klinicznych, intensywnych warsztatach, a także panelach dyskusyjnych. To wydarzenie cechujące się dużą interdyscyplinarnością, dlatego z pewnością każdy znajdzie coś dla siebie.

Podczas wydarzenia kompleksowo pochylimy się nad dziedziną rehabilitacji kardiologicznej i fizjologii wysiłku. Wśród tematów wiodących znajdują się:

- rehabilitacja w dobie pandemii i po pandemii COVID-19;
- telerehabilitacja i rehabilitacja hybrydowa;
- rehabilitacja kardiologiczna w specyficznych grupach pacjentów;
- programy KOS-zawał i KONS;
- nowe standardy ESC, PTK i SRKiFW;
- Testy wysiłkowe i testy spiroergometryczne
- monitorowanie wysiłku fizycznego;
- prewencja pierwotna i wtórna chorób sercowo-naczyniowych;
- farmakoterapia pacjentów rehabilitowanych kardiologicznie i nie tylko;
- sport i aktywność sportowa w kardiologii;
- czynniki ryzyka chorób układu krążenia.

Program merytoryczny wydarzenia jest niezwykle bogaty i angażujący. Warto podkreślić także, iż na konferencji pojawią się specjalne sesje wykładów prowadzone przez zaproszone sekcje i ассоjacje Polskiego Towarzystwa Kardiologicznego, m.in. Sekcję Kardiologii Sportowej, Asocjacje Niewydolności Serca, Asocjacje Elektrokardiologii Nieinwazyjnej i Telemedycyny, Sekcję Pielęgniarstwa Kardiologicznego i Pokrewnych Zawodów Medycznych, „Klub 30”, Sekcję Farmakoterapii Sercowo-Naczyniowej, Sekcję Prewencji i Epidemiologii, a także Polskie Towarzystwo Medycyny Sportowej.

„Pandemia wymusiła na nas zmianę paradygmatu rehabilitacji kardiologicznej”

Organizatorami wydarzenia są wydawnictwo naukowe Evereth Publishing oraz Sekcja Rehabilitacji Kardiologicznej i Fizjologii Wysiłku Polskiego Towarzystwa Kardiologicznego (SRKiFW). Przewodniczącą Komitetu Naukowego jest prof. dr hab. n. med. Małgorzata Kurpesa, Wiceprzewodniczącymi – prof. dr hab. n. med. Anna Jagier, dr hab. n. med. Dominika Szałewska, a Komitetu Organizacyjnego – dr n. med. Bartosz Szafran.

Dr n. med. Agnieszka Mawlichanów, Przewodnicząca SRKiFW, podkreśla, iż ostatnie Sympozjum miało miejsce w 2019 r. w Wiśle. W tym czasie udało się zorganizować wydarzenie w formule online, jednak zdaniem Przewodniczącej obecnie „wszyscy spragnieni jesteśmy spotkania osobistego, wymiany doświadczeń i bezpośrednich rozmów, nie tylko na sali wykładowej, ale i w kulisach”.

– Cztery lata w sporcie to pełna olimpiada, a w naszej dziedzinie kardiologii można powiedzieć – cała wieczność. Pandemia wymusiła na nas zmianę paradygmatu rehabilitacji kardiologicznej, między innymi stworzyła pole dla rozwoju modelu hybrydowego i monitorowanego telemedycznie. W tym czasie ukazało się wiele ważnych dokumentów, stworzonych przez polskie i europejskie towarzystwa kardiologiczne, dotyczące rehabilitacji, prewencji i aktywności fizycznej. Dynamicznie w naszym kraju rozwija się też program KOS-zawał, przynoszący liczne korzyści, ale też budzący kontrowersje. O tym wszystkim i jeszcze wielu innych sprawach pragniemy podyskutować w czasie naszego majowego spotkania – zapowiedziała dr Mawlichanów.

Rejestracja na 26. Sympozjum Sekcji Rehabilitacji Kardiologicznej i Fizjologii Wysiłku możliwa jest na stronie internetowej konferencji rehabilitacja2023ptk.pl/rejestracja/. Informacje na temat opłatyjazdowej i wydarzeń towarzyszących znajdują się tutaj: rehabilitacja2023ptk.pl/oplata-konferencyjna/.

Informujemy jednocześnie, iż liczba miejsc na konferencji jest ograniczona, dlatego warto zarejestrować się już dzisiaj.

Serdecznie zapraszamy do Hotelu Stok w Wiśle!

**Dołącz do najstarszego polskiego
towarzystwa naukowego
zrzeszającego fizjoterapeutów.**

Polskie Towarzystwo Fizjoterapii

od 1962 roku jako sekcja PTWzK

od 1987 roku jako samodzielne stowarzyszenie



- członek WCPT 1967-2019
- członek ER-WCPT 1998-2019
- projektodawca ustawy o zawodzie fizjoterapeuty (lipiec 2014)

Pracujemy w:

- 15 oddziałach wojewódzkich
- 10 sekcjach tematycznych

Odwiedź nas na stronie:

www.fizjoterapia.org.pl

i rozwijaj z nami polską fizjoterapię

Deklaracja członkowska dostępna jest w zakładce w menu strony

Multimodal programmes in the treatment of myofascial pain syndrome (MPS) – a two-step review

Programy multimodalne w leczeniu zespołu bólu mięśniowo-powięziowego (MPS) – przegląd dwuetapowy

Krzysztof Konior^{1(A,B,E,F,G)}, Aleksandra Bitenc-Jasiejko^{2(A,B,E,F,G)}, Danuta Lietz-Kijak^{2(A,E,G)}, Piotr Skomro^{2(B,E,G)}, Małgorzata Kowacka^{3(B,F,G)}, Konrad Kijak^{4(B,F,G)}, Zbigniew Śliwiński^{5(A,B,E)}

¹Department of Propaedeutic, Physical Diagnostics and Dental Physiotherapy, Faculty of Medicine and Dentistry Pomeranian Medical University, Szczecin, Poland

²Doctoral Study at the Department at the Propaedeutic, Physical Diagnostics and Dental Physiotherapy, Faculty of Medicine and Dentistry, Pomeranian Medical University, Szczecin; Medical Center in Nowogard, Poland

³Independent Public Health Care Complex in Żarki, Poland

⁴Student Scientific Society at the Department and Clinic of Internal Diseases, Angiology and Physical Medicine, Medical University of Silesia in Katowice, Poland

⁵Department of Physiotherapy, Collegium Medicum UJK in Kielce, Poland

Abstract

Myofascial pain syndrome (MPS) is one of the most common ailments associated with the human musculoskeletal system, characterised by the presence of the so-called trigger points (TrP – trigger point; MTrPs – myofascial trigger points). The International Association for the Study of Pain indicates that MPS may affect approximately one-third of people with chronic musculoskeletal pain, and that there is a lack of appropriate classification which can be attributed to a misunderstanding and/or misinterpretation of the pathophysiology. Given the diverse causes of pain syndromes in myofascial structures, it is vital to properly select and integrate therapeutic methods. The scientific literature indicates that treatment programmes should include a variety of manual therapy methods and rehabilitation exercises. Trigger point therapies, such as dry needling or dry cupping, are also widely used. At the heart of the success of rehabilitation programmes, in the opinion of the authors of this publication, is their multimodality, i.e. selection of therapeutic methods based on the cause of the pain, providing for measurable, reproducible diagnostic methods in therapy.

Aim of the study. The aim of this study is to analyse and infer conclusions on multimodal myofascial pain therapy programmes.

Material and methods. Given the complex research problem set as the aim, the study was carried out through a literature review in terms of two criteria:

Criterion I (C I): analysis of the literature on the etiology and pathogenesis of myofascial pain (i.e. causes and triggers, symptoms, social and environmental factors determining the onset of MPS), diagnostic procedures (initial diagnosis and ongoing monitoring of treatment outcomes), and therapeutic methods used in the course of MPS.

Criterion II (C II): a literature study of research publications addressing multimodal programmes for myofascial pain therapy, with their qualitative evaluation using the modified PEDro scale, and empirical testing of hypotheses based on the literature study and the analysis made in Part I.

Data sources: PubMed, SCOPUS, Science Direct, MEDLINE, PEDro, Cochrane, Embase, Web of Science Core Collection, Google Scholar electronic databases were searched systematically, restricting the languages to English and German only.

Results. The analysis of the literature showed that the causes, symptoms and associations of myofascial pain have been described in detail. There are also numerous reports on a variety of therapeutic methods, together with a precisely described methodology for their implementation. It is not uncommon to recommend combining methods into multimodal programmes, which unfortunately does not mean that there are many such programmes or that studies on MPS are consistent. The literature study on multimodal treatment programmes for MPS revealed that there is no correlation between its pathogenesis and a purposeful selection of specific therapeutic methods. In a small number of cases, a complex etiopathogenesis led to the formation of multidisciplinary teams. This may be associated with the absence of strict recommendations on the diagnostic methods applicable to the assessment of MPS.

Conclusions. 1. Multimodal programmes for the treatment of musculoskeletal pain, notably MPS and MTrPs, should include a detailed and comprehensive diagnosis (structural, biochemical, psycho-emotional) which should serve as the basis for the formation of interdisciplinary rehabilitation teams. 2. Musculoskeletal diagnosis, in addition to radiological assessment, should include measurable techniques of postural and functional assessment (such as pedobarography, wearable sensors, assisted anthropometry, i.e. photogrammetry, videogrammetry, etc.), aimed primarily at the ongoing assessment of posture. 3. The choice of therapeutic methods and patient education should be based on the causes of the patient's pain, taking into account systemic diseases, postural defects, lifestyle and psycho-emotional state. 4. Scientific research in multimodal treatment programmes should be carried out in randomised groups, with due attention to the methodologies of diagnostic and therapeutic procedures and group selection.

Keywords

chronic pain, myofascial pain, manual therapy, exercise, multimodal programmes

Streszczenie

Zespół bólu mięśniowo-powięziowego (MPS) jest jedną z najczęstszych dolegliwości związanych z układem mięśniowo-szkieletowym człowieka, charakteryzującą się obecnością tzw. punktów spustowych (TrP – trigger point; MTrPs – myofascial trigger points). International Association for the Study of Pain wskazuje, że MPS może dotyczyć około jednej trzeciej osób z przewlekłym bólem mięśniowo-szkieletowym oraz że brak jest odpowiedniej klasyfikacji, co można przypisać niezrozumieniu i/lub błędnej interpretacji patofizjologii. Biorąc pod uwagę różnorodne przyczyny powstawania zespołów bólowych w strukturach mięśniowo-powięziowych, istotny jest właściwy dobór i integracja metod terapeutycznych. Literatura naukowa wskazuje, że programy leczenia powinny obejmować różnorodne metody terapii manualnej oraz ćwiczenia rehabilitacyjne. Powszechnie stosowane są również terapie punktów spustowych, takie jak suche igłowanie lub suche łańki. U podstaw sukcesu programów rehabilitacyjnych, zdaniem autorów niniejszej publikacji, leży ich multimodalność, czyli dobór metod terapeutycznych w oparciu o przyczynę bólu, zapewniający mierzalne, powtarzalne metody diagnostyczne w terapii. Celem tego badania jest analiza i wyciągnięcie wniosków na temat multimodalnych programów terapii bólu mięśniowo-powięziowego. Materiał i metody. Biorąc pod uwagę złożony problem badawczy, jaki postawiono za cel, badanie przeprowadzono poprzez przegląd literatury pod kątem dwóch kryteriów:

Kryterium I (C I): analiza piśmiennictwa dotyczącego etiologii i patogenezы bólu mięśniowo-powięziowego (tj. przyczyn i wyzwalaczy, objawów, czynników społecznych i środowiskowych determinujących wystąpienie MPS), postępowania diagnostycznego (wstępna diagnostyka i bieżąca obserwacja wyników leczenia), i metody terapeutyczne stosowane w przebiegu MPS.

Kryterium II (C II): studium literaturowe publikacji naukowych dotyczących multimodalnych programów terapii bólu mięśniowo-powięziowego wraz z ich oceną jakościową przy użyciu zmodyfikowanej skali PEDro oraz empiryczne testowanie hipotez w oparciu o studium literaturowe i analizę dokonaną w części I.

Źródła danych: PubMed, SCOPUS, Science Direct, MEDLINE, PEDro, Cochrane, Embase, Web of Science Core Collection, Google Scholar systematycznie przeszukiwano elektroniczne bazy danych, ograniczając języki tylko do angielskiego i niemieckiego.

Wyniki. Analiza piśmiennictwa wykazała, że szczegółowo opisano przyczyny, objawy i powiązania bólu mięśniowo-powięziowego. Istnieją również liczne doniesienia o różnych metodach terapeutycznych wraz z dokładnie opisaną metodologią ich realizacji. Nierzadko zaleca się łączenie metod w programy multimodalne, co niestety nie oznacza, że takich programów jest wiele i że badania nad MPS są spójne. Studium literaturowe dotyczące multimodalnych programów leczenia MPS wykazało, że nie ma związku między jego patogenezą a celowym doбором określonych metod terapeutycznych. W niewielkiej liczbie przypadków złożona etiopatogeneza doprowadziła do powstania zespołów wielodyscyplinarnych. Może to być związane z brakiem ścisłych zaleceń dotyczących metod diagnostycznych mających zastosowanie do oceny MPS. Wnioski. 1. Multimodalne programy leczenia bólów narządu ruchu, zwłaszcza MPS i MPPS, powinny zawierać szczegółową i kompleksową diagnostykę (strukturalną, biochemiczną, psychoemocjonalną), która powinna stanowić podstawę do tworzenia interdyscyplinarnych zespołów rehabilitacyjnych. 2. Diagnostyka narządu ruchu, oprócz oceny radiologicznej, powinna obejmować mierzalne techniki oceny posturalnej i funkcjonalnej (takie jak pedobarografia, czujniki ubieralne, antropometria wspomagana, tj. postawy).

3. Wybór metod terapeutycznych i edukacja pacjenta powinny być oparte na przyczynach dolegliwości bólowych pacjenta, z uwzględnieniem chorób ogólnoustrojowych, wad postawy, stylu życia i stanu psychoemocjonalnego. 4. Badania naukowe nad multimodalnymi programami leczenia powinny być prowadzone w grupach losowych, z należywym uwzględnieniem metodologii postępowania diagnostyczno-terapeutycznego oraz doboru grup.

Słowa kluczowe

ból przewlekły, ból mięśniowo-powięziowy, terapia manualna, ćwiczenia, programy multimodalne

Introduction

Approximately 85% of the human population experience myofascial pain at least once in their lifetime [1]. Its prevalence varies, ranging from 21% of patients presenting with orthopedic problems up to even 93% of patients undergoing treatment at pain management centres [2]. Overall, musculoskeletal pain has been shown to affect about 26% of the adult population, 39% of the elderly population and about 86% of workers [3]. MPS often leads to disability, accounting for approx. 40–50% of the economic cost associated with disorders affecting the quality of work in the European Community. These disorders are not life-threatening, but have a definite impact on the quality of life, psychosocial status, etc. [4–6].

The onset of specific myofascial pain may be acute (although this is not a rule) with concomitant autonomic symptoms [7–8]. Chronic pain in myofascial structures is often a complex condition with no clear clinical definition, while the recurrence and persistence of symptoms complicates the process of planning and selecting therapeutic approaches [9–11].

MPS is often defined as a pain syndrome in a specific area of the body, which may coexist with one or more pain foci, or the so-called trigger points [12–13]. While the pain foci have been defined in different ways, it is becoming increasingly common to refer to them as myofascial trigger points (MTrPs) [13–16]. Recurrent MPS has been shown to be closely associated with the presence of MTrPs [9–10, 17–18]. Trigger points are a common reason for patients presenting to primary care and pain clinics [19–21].

Myofascial pain, triggered by MTrPs, manifests itself as areas/spots/bands of tense myofascial tissue [17, 22–25]. When pressed, they can elicit local and radiating pain. The pain may cause the patient to restrict their mobility, which in the long term can lead to motor dysfunction. Pain also significantly contributes to the development of symptoms associated with the autonomic system [13, 23, 26–32]. Autonomic symptoms include vasodilation or vasoconstriction, lacrimation (tearing), piloerection (hair standing) [33–36]. Active trigger points may cause spontaneous pain, while latent points are only painful upon compression [21, 23, 26–28, 37]. Pressure, needle insertion and other methods of trigger point stimulation can elicit a local twitch response, defined as an involuntary, rhythmic muscle contraction at or near the MTrP site [21, 38]. The location of the twitch response is easier to identify in the case of active trigger points [23, 39–40].

Upon physical examination, MTrP sites show an altered muscle tone, hypersensitivity to touch, often during stretching, leading to restriction of movement or eliciting jump sign. Sometimes there is a palpable or even visible tremor [7, 22]. For the most part, patients describe trigger points as a significantly unpleasant sensory and emotional experience. Depending on the actual and potential tissue damage, the patient's subjective perception may differ [41].

Myofascial pain can be classified in a number of ways, as acute or chronic, neurogenic, accompanying inflammation, visceral, somatic, nociceptive, etc. [42–43]. Depending on the location of the pain, MTrPs can trigger focal or referred pain (felt at a site distant from the site of the painful stimulus). [31, 44–46]. Sustained contractile activity of a trigger point in the

muscle induces local ischemia, hypoxia and changes at nociceptors, leading to pain in deep myofascial structures and at distant locations [1]. The muscle fascia contains mechanoreceptors sensitive to changes in length, muscle tension and neurological stimuli. Some theories cite abnormalities in fascial structures as the main factor in MPS development [47–48]. MTrPs can develop in different muscle groups, but they are most common in the sternocleidomastoid, trapezius, levator scapulae, infraspinatus, and rhomboid muscles of the upper back and neck [49–52]. Chronic headache, including migraine, and chronic complaints in the upper quadrant (quarter pain), as well as craniofacial complaints are often attributed to MTrPs [20, 53–57]. Myofascial pain has also been shown to affect 95.5% of patients with low back pain, occurring in the paraspinal, piriformis, tensor fascia lata muscles [58] and also in patients with intervertebral disc diseases [59] and spinal cord compression. [60]. According to the research literature, myofascial pain and associated trigger points can occur in almost any musculoskeletal ailment, including tendonitis [61], craniomandibular dysfunctions [62–63], in the course of carpal tunnel syndrome [64], pelvic pain and urologic diseases, etc. [65–68]. The research literature also refers to the phenomenon of chronic MPS as a postoperative complication. It is defined as moderate to severe pain lasting at least three months after surgery [69]. For instance, chronic postsurgical pain (CPSP) of neuropathic origin persisting beyond the normal healing time is reported by 6–10% of patients undergoing total knee arthroplasty [70]. This phenomenon should be recognised as a serious health problem, seeing as prolonged postoperative use of opioids and adjuvant analgesics represents a risk of addiction in total knee arthroplasty (TKA) patients, leading to significant economic costs for this group [70–71].

It should be emphasised that symptoms have been challenged to varying degrees, which may be related to misidentified pathophysiology [72–74].

Aim of the study

The aim of this study is to identify the relationships between the multifaceted etiopathogenesis of myofascial pain and the development of multimodal programmes for managing MPS and, as a consequence of the above considerations, to conduct an analysis on the selection of appropriate diagnostic and therapeutic methods and the formation of an interdisciplinary team.

Materials and methodology of the study

The papers included in the literature analysis below were dedicated to a detailed investigation of the causes, symptoms and impact of systemic diseases and external factors on the formation of MPS. The results of this analysis helped guide further research, involving on a review of the literature on multimodal treatment programmes for MPS. Given that the research and analysis was carried out in two stages, the aim was pursued through a database search according to two sets of search criteria:

Criterion I (C I): analysis of the literature on the etiopathogenesis of myofascial pain (i.e. causes and triggers, symptoms, social and environmental factors determining the onset of MPS), therapeutic methods used in the course of MPS and diagnostic procedures (initial diagnosis and ongoing monitoring of treatment outcomes).

Source of data (C I): PubMed, MEDLINE, PEDro, Web of Science Core Collection, Cochrane, Embase, Google Scholar. Search keywords: myofascial pain syndrome, myofascial trigger point, and these keywords in combination with: manual therapy, muscle energy techniques, joint mobilization, neuromobilization, dry needling, dry cupping, kinesiotaping, therapeutic exercises, home exercises, patient education.

Inclusion criteria (C I): the papers included in the analysis had to be published in peer-reviewed journals, in English and one article in German, discussing scientific research studies relevant to the aim and main issue of this review. Only content confirmed in more than 3 peer-reviewed publications was analysed.

Exclusion criteria (C I): content at the stage of hypothesis, research projects, expert opinions, etc. was not accepted for analysis.

Criterion II (C II): a literature study of research publications addressing multimodal programmes for myofascial pain therapy, with their qualitative evaluation using the modified PEDro scale, and empirical testing of hypotheses based on the literature study and analysis made in Part I.

The literature was tested through the authors' research hypotheses (H), formed on the basis of the literature study in Part I, that is:

H1: Multimodal treatment programmes for MPS rely on the strict methodology of scientific research work

a. They account for age, sex, weight, with a multi-centre approach to patient selection,
b. Participant selection takes place by random sampling, with the use of blinding (therapist/patient, sample, etc.), follow-up, intention to treat, comparison of groups, final score estimation, statistical analysis,

H2: The selection of therapeutic methods to be included in a multimodal treatment programme for MPS is based on a detailed diagnosis (measurable, reproducible, objective) of the patient's condition, including comorbidities,

H3: In the case of comorbidities associated with MPS, a multidisciplinary team is set up,

H4: Therapeutic methods and patient education are selected depending on the diagnosis.

Source of data (C II): Database search was conducted in PubMed, MEDLINE, PEDro, Web of Science Core Collection, Cochrane, Embase, Google Scholar. Searched keywords were: myofascial pain syndrome and myofascial trigger point, and these keywords in combination with: myofascial pain and myofascial pain in combination with multimodal, multi-modal, approach, integrated methods, treatment, non-invasive treatment, interdisciplinary multimodal pain therapy.

Inclusion criteria (C II): the papers included in the analysis had to be published in peer-reviewed journals, in English, discussing scientific research studies relevant to the aims and main issue of this review, in which multimodal treatment pro-

grammes were implemented and evaluated. The analysed content addressed MPS treatment programmes with a description of integrated programmes and/or provided for a role of multidisciplinary teams. Thus, our analysis included articles that described 2 or more rehabilitation methods (e.g. manual therapy, exercise, dry needling, dry cupping) and/or physical therapy and/or diet therapy, psychotherapy, etc.

Exclusion criteria (C II): content at the stage of hypothesis, research projects, expert opinions, etc. was not accepted for analysis. Also excluded were manuscripts containing recommendations for multimodal treatment programmes based on a literature review (those were included in Part I of this study). Moreover, articles on pharmacological and surgical methods and articles indicative of research work focusing on only one rehabilitation method were excluded from the analysis.

Main issue – Criterion I

Etiopathogenesis of myofascial pain

The causes of myofascial pain and MTrPs, and the resulting impairment of musculoskeletal function, are not entirely clear. One concept is the release of neurovasoactive substances (i.e. bradykinin (BK), prostaglandin, interleukin-1 β , substance P), which enhance nociceptor sensitivity and vascular permeability, leading to edema. Consequently, the pressure on the capillaries and restriction of blood supply causes local ischemia, increasing the release of substance P, which exacerbates tissue irritation. Another theory, of the so-called motor end plate, points to a neuromuscular connection by which painful contraction, triggered by the release of acetylcholine, leads to the release of calcium stored in the muscles and sarcomere shortening. There is also a theory whereby hypersensitive points arise as reflex disorders due to the improper functioning of spinal nerves and/or abnormalities in the spine [43, 75–85]. Lastly, patients with myofascial pain were found to present pathological changes in red fibres, known as “moth-eaten fibres” [86–90].

The musculoskeletal system is the largest organ in the human body, and its system of force and pressure distribution can be associated with different types of pain, often coexisting not only with problems specific to musculoskeletal structures, but also with systemic diseases, postural defects, psycho-emotional disorders, etc. [42, 50, 91]. The most commonly reported causes of MTrPs include trauma and structural overload [16, 36, 92–95], coexisting osteoarticular and neurological dysfunctions [14, 45, 96–100], psycho-emotional state and stress [101], and damage to muscle fibres [17, 102–104]. Other causes of myofascial pain include adaptive and eccentric muscle lengthening, postural defects, joint hypermobility, inappropriate physical and social conditions – these can activate MTrPs [17, 105]. Myofascial trigger points may be associated with other pain syndromes [106], i.e. postherpetic neuralgia [107–108], complex regional pain syndrome and phantom pain [109–110], muscle spasms [111], and rare diseases etc. [112–113].

As a result, each of these factors impairs the supply of oxygen and nutrients to tissues, resulting in compensatory shortening of the muscles and consequently increased metabolic demand on the tissues [23, 114]. Researchers have also implicated nutritional deficiencies (vitamin C, B, D, iron, zinc, folic acid) and fo-

od intolerances (e.g. lactose, casein, gluten) in activating MTrPs [93, 105, 115–118].

Diagnosis of myofascial structures and posture

Detailed history, gait and posture analysis, palpation and compression, functional assessment (ranges of motion), assessment of referred pain projection and local twitch response all contribute to an accurate diagnosis, but there are no clearly defined methods enabling an unambiguous diagnosis of trigger points [119–121]. Taut bands of muscle fibre can be identified by electromyography [122], sonoelastography [123], thermography (infrared imaging) [124], and magnetic resonance imaging [125]. Biochemical diagnostics have been performed, too [126].

Diagnostic procedures and monitoring are also recommended to track progress in therapy using all integrated therapeutic methods. The methods for assessing the effects of manual therapy, exercise, stretching, joint mobilization etc. should be measurable, objective and reproducible, which is where computer techniques can come in, measuring ranges of motion, muscle performance, and general movement patterns (such as pelvic kinematics, gait, balance etc.). These include sensor systems for postural and functional assessment [127–131], pedobarography [132–138], videogrammetry, photogrammetry etc. They are applicable both in the initial evaluation, ongoing monitoring and in planning further stages of therapy, postural re-education and stabilization [139–146].

Progress monitoring in therapy, but also follow-ups at specific time points after therapy, are important due to the perpetuating factors promoting the persistence of myofascial pain. These include mechanical problems (e.g. head protraction, pelvic torsion, biomechanical disorders), psycho-emotional (e.g. stress), nutritional and metabolic factors [21, 31, 147]. The management of perpetuating factors requires the inclusion of a multidisciplinary team [105].

Non-pharmacological and non-surgical therapies for myofascial structures used in myofascial pain

Manual therapy

The efficacy of various manual therapy techniques can be attributed mainly to the effects of oxygen and nutrient uptake, both during the treatment and during the release and relaxation of structures in the course of manual therapy [20]. The techniques used in the treatment of myofascial pain, in addition to classic massage, include ischemic compression for the treatment of MTrPs, muscle energy technique (MET), strain-counterstrain, spray and stretch, static stretching, transverse friction massage, joint mobilization or manipulation, and other soft tissue mobilization techniques [148–157]. Muscle energy techniques aim to cause the muscle to lengthen, strengthen, to reduce swelling and relieve passive congestion. They produce a rapid therapeutic effect in the course of acute pain in myofascial structures [158–160]. METs are also widely used for joints with restricted mobility [161–162], as are some other methods, e.g. AKA-H (arthrokinematic approach-Hakata) or HVLA (high velocity low amplitude), intended to improve joint glide, which also produces an indirect effect on the functionality of myofascial structures [163–166]. At this point, it is worth emphasising that, in terms of the range of motion, depending on

the pathophysiology of the pain, for any therapeutic intervention to be effective, manual therapy must address both myofascial and joint structures. This knowledge inspired the development of therapeutic approaches combining these two aspects, such as e.g. the Mulligan concept, the SNAG (sustained natural apophyseal glide) technique, which involves the application of accessory passive glide along the joint plane while the patient simultaneously performs an active movement, under loading conditions [167]. This consequently improves the functionality of the structures and increases the range of motion [167–170]. In pain therapy, it is important to restore nerve mobility relative to adjacent structures, which can be achieved by manual therapy using neural mobilization techniques. If the intervention is applied as early as possible, it may be possible to avoid morphological changes in neural structures [171–172].

In manual therapy procedures, apart from the local approaches, an important role is played by the tensegrity concept, i.e. a holistic approach based on structural integration. It has been shown time and again to be one of the most effective rehabilitation methods in balancing musculoskeletal strains and relieving chronic myofascial pain, including trigger point therapy [173–175]. This is important mainly because excessive tension in tissues represents a damaging stimulus which is distributed in the human body in a linear manner [176–178]. Pain originating in myofascial structures can therefore be felt at a different location (sometimes even quite distant) than the initial pain stimulus, or TrP [179–183]. The inclusion of tensegrity models in diagnosis and therapy helps restore structural balance in the patient's body. It is an important aspect of MPS therapy, with implications for increasing structural functionality, proper force/load distribution, and activation of stabilising muscles [182]. As a result of this intervention, muscles regain the important functions of cushioning, anti-gravity and stabilization of osteoarticular structures [173–174, 182, 184]. This is particularly relevant in chronic and recurrent pain, e.g. low back pain [176–178]. Integrated manual therapy methods include myofascial release (MFR) [185–161] and osteopathic techniques (osteopathic manipulative treatment – OMT). OMTs are used to treat myofascial structures, as well as joint manipulation and mobilization (including MET), and visceral manipulation (of the internal organs) [158–159, 191].

Therapeutic methods dedicated to the elimination of MTrPs

Equally effective and popular interventions are aimed at eliminating trigger points, and involve the local application of needles (dry needling) or cups (dry cupping) [20]. Dry needling can be used in the treatment of pain of myofascial origin in all regions of the body, e.g. head, back [22, 150, 192], lower and upper limbs [150, 193–194], etc. The main effects of dry needling include pain relief, through stimulating the spinal cord and sympathetic response, activation of biochemical changes in tissues through improving local microcirculation, reduction of chemical mediators and induction of an immune response [192]. Research studies have also shown that dry needling corrects the levels nociceptive chemicals, such as bradykinin, calcitonin-gene-related peptide, and substance P [22]. Stimulation of alpha-delta nerve fibres leads to the release of endorphins and enkephalins [150, 192, 195]. It has been suggested that this

process results in opioid-mediated pain suppression [197]. All these processes help reduce muscle tension and increase range of motion in myofascial structures, thereby reducing pain [198]. The insertion of a needle at the trigger point as well as palpation can elicit a twitch response [23, 26–28]. Some studies indicate that the analgesic and relaxing effect is achieved as a result of the twitching induced by dry needling [23, 39, 199].

Trigger points can also be managed by dry cupping, which are used in the treatment of painful trigger points, increased tension and other complaints related to myofascial structures mainly to increase subcutaneous blood flow and dilate blood vessels, which improves microcirculation, relieving muscle tension and promoting healing. These metabolic effects are similar to those of dry needling [200]. Dry needling treatments are usually followed by stretching exercises [32].

Improved local blood flow, regeneration of pain points and normalization of tension in the fascial system with reduced pressure on pain receptors can also be achieved with elastic therapeutic tape (Kinesio Taping – KT). It augments lymph circulation, reduces swelling and enables local decompression. It is also used to relax and mobilize long structures and in therapies based on the anatomy trains concept [201–205].

Physical therapy in the treatment of myofascial pain

Physical therapy treatments are widely used in the management of pain in myofascial structures:

- shockwave therapy – is designed to normalise tension in myofascial structures and eliminate MTrPs, both active and latent [206–207],
- magnet therapy, which through the use of a constant and slowly changing magnetic field increases the secretion of opiates from the β -endorphin group, thereby increasing pain response latency and the pain threshold [185,208–212],
- electrotherapy and electrical stimulation, which promotes the release of adenosine triphosphate (ATP), seeing as low ATP levels may be responsible for the activation of MTrPs [211–213],
- ultrasound therapy, which works on both active and latent trigger points [196, 214–215].

Pain in the pelvic area, of urologic or gynecologic origin, can be situated inside the body. This is an important aspect, mainly due to the need for adapting therapeutic methods and sometimes employing additional therapeutic tools [216–217]. Despite the fact that laser therapy, as a matter of principle, improves microcirculation and eliminates metabolic by-products from tissues, it has been demonstrated that placebo treatment produced better effects than low-level laser therapy (LLLT) [160].

Rehabilitation exercises as part of therapy and home exercise

Next to manual therapy, exercise plays an important role in the rehabilitation of myofascial pain. However, in this case, too, the type of exercise depends on the therapeutic goal, ranging from decompression and relaxation exercises for pain relief (e.g. the McKenzie method) [160, 218–220] to core stability exercise with strength training, such as Yoga [221–223]. Eccentric exercises have been shown to have a positive effect in the therapy of both active and latent trigger points [224–226]. Stretching exercises and Yoga also improve respiratory function and the psycho-emotional state of pain patients, which is an important aspect in the therapy of pain, both acute and chronic [227–228]. Core stability training has the important role of reducing structural instability, which is important in the prevention of recurrent pain syndromes in affected tissues, particularly in spinal pain [229–231]. Core stability training is more effective if performed using rehabilitative ultrasound imaging of isolated muscle groups, aiming to ensure proper contraction while maintaining joint mobility [232–235].

Patient education

Patient education, particularly in the case of chronic myofascial pain, is aimed at providing knowledge on pain management, regulation of rest and physical activity. A cognitive-behavioural intervention is also required, involving pain neuroscience education, belief revision, and helping accept and adapt to flare-ups and (temporary or permanent) disability [236–238]. Learning to manage flare-ups includes both stress management and the implementation of various elements of self-therapy, e.g. relaxation, decompression techniques etc. [239–241].

Manual therapy (MT) and exercise, performed during therapy sessions as well as at the patient's home, are effective, but recommendations and the research literature are not always clear on how often they should be applied for the optimal effect [22]. The combination of MT and exercise with dry needling produces therapeutic effects in both chronic pain and postoperative conditions [242–243]. These are relatively low-cost and minimally invasive techniques, producing similar short-term analgesic effects [244].

Our literature study according to Criterion I enabled us to identify the problem of multifaceted etiopathogenesis of MPS, which points to the need for multimodal programmes in the treatment of myofascial pain, targeting the specific cause.

Based on the literature study, the different causes (pathogenesis) of MTrPs and myofascial pain have been matched to their dedicated therapeutic approaches, as summarised in Table 1.

Table 1. An overview of the causes of MTrPs with the therapeutic approaches aimed at eliminating the cause/pathogenic factors

Cause	Therapeutic method targeting the cause
Active and/or latent trigger points in general, local pain syndromes and phantom pains (also with concomitant swelling, local fever)	<p>Methods to eliminate pain:</p> <ul style="list-style-type: none"> • manual therapy – ischemic compression, • local methods: dry needling and dry cupping, kinesiotaping, • manual therapy: MET, • strain-counterstrain, • spray and stretch technique.

Cause	Therapeutic method targeting the cause
Trigger points caused by nerve compression or entrapment (without signs of swelling or local fever, with radiating pain)	Manual therapy based on neural mobilization
Trigger points and myofascial pain caused by long-term joint immobilization.	<ul style="list-style-type: none"> • Bone and joint mobilisation: MET, HVLA, AKA-H, • myofascial therapy using structural integration procedures (tensegrity) e.g. according to the anatomy trains concept – MFR, OMT, • exercise: the McKenzie method, Yoga.
Trigger points coexisting with remote pain; Postural defects	<ul style="list-style-type: none"> • Myofascial therapy using structural integration procedures (tensegrity) e.g. according to the anatomy trains concept – MFR, OMT, • stretching exercises: the McKenzie method, Yoga, • corrective exercises for strength and stability training.
Pain of psycho-emotional origin	Therapeutic methods targeting the cause of myofascial pain and psychotherapy (a multidisciplinary team including a psychotherapist).
Pain caused by nutritional deficiencies	Therapeutic methods targeting the cause of myofascial pain and clinical nutritional therapy (a multidisciplinary team including a nutritionist, biochemistry specialist).
Systemic diseases	A multidisciplinary team including a specialist in the field dedicated to the treatment of the patient's specific disease entity

The above literature study has raised important questions about the procedures undertaken in the treatment of myofascial pain. These questions then became the foundation for the literature analysis in Part Two.

Main issue – Criterion II

The literature review in Part I revealed the high complexity and multifaceted nature of the pathogenesis of MPS. It also identified a number of therapeutic approaches to the treatment of MPS, targeting the specific cause and symptoms of myofascial pain. Most research studies mention the need for the integration of therapeutic approaches and a multidisciplinary approach to MPS. Our quantitative analysis showed, however, that there are few randomised trials investigating multimodal treatment pro-

grammes for MPS that would simultaneously address the following three important aspects: method selection based on etiopathogenesis, integration of therapeutic approaches and interdisciplinarity of the treatment team. For the most part, such comprehensive studies were based on literature reviews. There are many studies on the efficacy of a specific therapeutic method based on case studies, whose findings should be treated with caution as they do not represent the entire population. The literature analysis also revealed a great deal of inconsistencies and a lack of clear recommendations regarding diagnostic methods for MPS. For our detailed analysis, we selected seven articles on multimodal programmes which reported on research with groups of patients. The findings, together with the auxiliary questions to test the hypotheses, are shown in Tables 2–6.

Table 2. Testing H1: Background information on sample size, sex, age, weight and location of patients (i.e. single-centre or multi-centre study)

Test questions	Articles included in the analysis						
	1[245]	2[10]	3[246]	4[247]	5[248]	6[249]	7[217]
Sample size [number of people]	150	72	9	14	120	120	393
Age (minimum) [0–no data, or age given]	0	18	20–40	0	0	18	18
Age (maximum) [0–no data, or age given]	0	65	65	0	0	18 years plus	80

Test questions	Articles included in the analysis						
	1[245]	2[10]	3[246]	4[247]	5[248]	6[249]	7[217]
Number of male patients [0=no data, or number of male patients given]	0	0	0	0	0	0	314
Number of female patients [0=no data, or number of female patients given]	0	0	0	0	0	0	79
BMI [0=no data, or result given]	0	0	21 ± 3.29	0	0	0	0
Number of centres participating in the study [number]	1	1	1	1	1	1	1

Table 3. Methodology testing using the modified PEDro scale [250]

Test questions	Articles included in the analysis						
	1[245]	2[10]	3[246]	4[247]	5[248]	6[249]	7[217]
Was there a research protocol with inclusion and exclusion criteria?	1	1	1	1	1	1	0
With randomisation (random allocation of patients into groups)?	0	1	0	0	1	1	0
Was group allocation blinded?	0	1	0	0	1	1	0
Were study groups analysed statistically for homogeneity of patient characteristics and baseline measurements (in the research tools used in the study)?	0	1	0	0	1	1	0
Was a single-blind design used (with the patients unaware of their allocation to the treatment or control group)?	0	0	0	0	0	0	0
Was a double-blind design used (with the researchers responsible for therapy unaware of the patients' group allocation to the treatment or control group)?	0	0	0	0	0	0	0
Was a double-blind design used (with the researchers responsible for diagnostics unaware of the patients' group allocation to the treatment or control group)?	0	0	0	0	1	1	0
Were the patients subjected to follow-up after completion of the study?	1	1	1	0	1	1	1
Was the study conducted with the intention to treat?	1	1	1	1	1	1	1
Were the results analysed statistically by intergroup comparisons?	1	1	0	0	1	1	0
Did the analysis include variability measurements and final point estimation?	1	1	0	0	1	0	0
Were treatment outcomes analysed statistically?	1	1	1	1	1	1	1

Key – Table 3: 0 – no data relating to the test question asked, 1 – YES, i.e. an affirmative answer to the test question asked

Table 4. Testing H2: Diagnostic procedures used in the implementation of a multimodal myofascial pain programme

Test questions	Articles included in the analysis						
	1[245]	2[10]	3[246]	4[247]	5[248]	6[249]	7[217]
Were diagnostic methods for the initial assessment of the patients' condition identified?	1	1	1	1	1	1	1
Was there an assessment of pain and tension at trigger points?	1	1	1	1	1	1	0
Were pain scales used? (if so, which ones?)	VAS	VAS-P / GPE	0	VAS	Electronic pain measurement	VAS	0
Was there an assessment of posture?	0	0	0	0	1	1	0
Was there an assessment of function/fitness? (if so, what kind)	0	DASH	CF-PDI/NDI	WOMAC	MANUAL	MANUAL	0
Was there an assessment of gait determinants?	0	0	0	1	0	0	0
Were diagnostic methods for assessing ongoing	1	1	1	0	1	0	0

Key – Table 4: 0 – no data relating to the test question asked, 1 – YES, i.e. an affirmative answer to the test question asked; descriptive responses are the names of the pain scales and the diagnostic methods used in the case of affirmative answers.

Table 5. Testing H3: Participation of a multidisciplinary team in the multimodal approach to myofascial pain therapy

Test questions	Articles included in the analysis						
	1[245]	2[10]	3[246]	4[247]	5[248]	6[249]	7[217]
Was a multidisciplinary team involved in myofascial pain therapy?	0	0	1	0	0	0	1
Were pain complaints differentiated depending on their origin: myofascial, neurological, psycho-emotional, biochemical, nutritional deficiencies?	1	1	1	0	0	0	1
Was a diet therapy specialist included in the treatment team?	0	0	0	0	0	0	0
Was nutritional supplementation provided?	0	0	0	0	0	0	0
Was an assessment of psycho-emotional state included?	0	0	1	0	0	0	0
Was a psychotherapist included in the treatment team?	0	0	1	0	0	0	0
What diagnostic procedures were used?	0	0	1	0	0	0	0
What behavioural interventions?	0	0	1	0	0	0	0
Were systemic diseases included in the diagnostic process?	0	1	0	0	0	0	1
Were specialists of the relevant systemic diseases involved in the therapeutic process?	0	0	0	0	0	0	1
Were the results analysed statistically?	1	1	1	1	1	0	1

Key – Table 5: 0 – no data relating to the test question asked, 1 – YES, i.e. an affirmative answer to the test question asked.

Table 6. Testing H4: Procedures for selecting therapeutic modalities to be used in the multimodal / interdisciplinary programme for myofascial pain therapy

Test questions	Articles included in the analysis						
	1[245]	2[10]	3[246]	4[247]	5[248]	6[249]	7[217]
Was the number of therapy sessions determined? [0– no information, or number given]	5	12	6	10	30	10	4 sessions per week, for 6 months
Was the number of procedures to be applied during each session specified? [0–no information, 1–YES]	1	0	0	0	1	1	1
Duration of therapy [weeks]	4	12	0	4	10	5	“6 days at the facility 6 months at home”
Is there information on which therapies were applied during a particular session? [0 – no information, 1 – YES]	1	0	0	0	1	1	1
What therapeutic methods were used?							
Ischemic compression of trigger points	0	1	0	0	1	0	1
Manual therapy of myofascial structures (joint mobilization, neural mobilization, dynamic soft tissue mobilization)	0	0	1	0	1	1	0
Strain counterstrain (SCS)	0	0	0	0	1	0	0
Dry needling of trigger points	1	0	0	1	0	1	0
Spray and stretch	1	1	0	0	0	0	0
Kinesio taping	1	0	0	0	0	0	0
Therapeutic exercise	1	0	1	1	0	1	1
Other (which?)	0	0	0	0	Deneroll	0	Internal Myofascial Trigger Point Wand and Padoxical Relaxation Therapy (PRT)
Has the treatment methodology been described in detail?	1	0	1	1	1	1	1
Was Patient Education provided?	1	0	1	0	1	1	1
In therapy, was it specified on which areas of the body the therapy was carried out? *	0	1	1	1	1	1	1
How many people were treated in a given area of the body? [number]							
Cervical spine, shoulders, arms	102	0	9	0	120	120	0
Craniofacial (temporomandibular joint area)	1	0	9	0	0	0	0
Shoulder	0	72	0	0	0	0	0
Upper limb (elbow)	14	0	0	0	0	0	0
Lumbosacral	30	0	0	0	0	0	0
Pelvis (pain of urologic/gynecologic origin)	0	0	0	0	0	0	393
Lower limb (hip, knee)	0	0	0	14	0	0	0
Foot and ankle joint	10	0	0	0	0	0	0
TOTAL	157	72	9	14	120	120	393

Key – Table 6: 0 – no data relating to the test question asked, 1 – YES, i.e. an affirmative answer to the test question asked. The table sections below the question: In therapy, was it specified on which areas of the body the therapy was carried out? present the number of people who had therapy in the specific area of the body.

Discussion

As shown in Part I of our study, research dedicated to myofascial pain therapy programmes needs to take into account two essential aspects, i.e.:

1. this type of pain affects around 90% of the human population,
2. it has a complex etiology, involving not only short-term complaints (e.g. due to trauma, periodic dysfunctions), but mainly chronic conditions related to posture, systemic diseases, work, social status and psychophysical state.

Non-traumatic myofascial pain in most cases can be elicited by trigger points [251–252]. Many studies confirm the efficacy of MFTTrP therapies, e.g. dry needling in randomised groups, in different areas of the body [253–258]. However, empirically, it can be concluded that in the case of comorbidities (i.e. systemic diseases, postural defects, sensory-motor deficits, psycho-emotional problems, etc.) the pain may return in the same and/or different location or form. Hence, it is crucial to combine therapeutic methods in multimodal programmes, involving an interdisciplinary team. Moreover, multimodal programmes should provide for a careful selection of therapeutic modalities, based on an objective and reliable diagnosis of MPS etiopathogenesis. Despite the recommendations for measurable diagnostic methods in myofascial pain, such as EMG, sonoelastography, thermography (infrared imaging) and MRI [122–125, 259], the majority of the research studies dedicated to multimodal programmes used palpation and inspection methods in the evaluation of both function and pain [260–261].

In the majority of the articles included in our analysis, pain was assessed using the Visual Analogue Scale (VAS). Moustafa et al. (2018) used an algometric measurement of the pressure pain threshold (PPT) at the most tender point of the upper trapezius muscle before and after treatment [262]. The average value of 3 repetitive measurements with an interval of 30 to 60 s (expressed as kilograms per square centimetre) was used to analyse the PPT data [263]. It should be noted, however, that the analysis did not include an assessment of latent and referred trigger points, which was characteristic of the majority of the articles reviewed in Part II of the study. Bron et al. (2011) assessed secondary outcome measures using the VAS-P and GPE scales (which is used to differentiate changes in pain and function) [10]. They also used the DASH questionnaire, which is recognised as one of the best questionnaires for patients with shoulder symptoms [264–265].

From the clinical perspective of myofascial pain, given that trigger points in the latent phase can alter activation patterns and/or cause weakness in pain-free muscles [266–268], through structural, functional, compensatory, etc. relationships, muscle tension and range-of-motion testing should be included in the diagnosis. In three of the seven articles analysed, a manual goniometer was used to measure ranges of motion (Bron et al. 2011, Marcos-Martín et al. 2018, Moustafa et al. 2018) [10,246–248]. Núñez-Cortés et al. used the WOMAC scale and a digital inclinometer for both disability diagnosis and functional assessment [247]. WOMAC is used extensively in clinical trials, mainly in surgery and traumatology, notably for its sensitivity and validity [269]. Unfortunately, the use of both a manual goniometer and an inclinometer is fraught with scale discrepancies. In this case, the Constant-Murley scale was used, with subscales including subjective pain and mobility [270–272]. In the assessment of disability, in

turn, Marcos-Martín et al. (2011) employed the CF-PDI method for craniofacial assessment, and NDI for cervical spine assessment [247]. The Neck Disability Index (NDI) is a questionnaire with 10 items referring to the activities of daily living, in which patients rate their subjective experience [272–274]. The reliability, construct validity and responsiveness of NDI have been studied. In addition, cervical and shoulder alignment and head translation posture were assessed using photogrammetry, which provides reliable data for postural anthropometry [275–277].

In rehabilitation diagnosis, it is important to assess baseline functional parameters and then evaluate treatment progress on an ongoing basis. This is because the outcome may be affected by both compensatory movement and the original constraint. Hence the strict recommendation to use measurable, reproducible diagnostic methods in the assessment of posture and function [278–279]. Indeed, the development of MPS is heavily influenced by global postural defects, abnormal locomotor patterns, physical conditions (including weight) etc. [17, 49, 105]. The literature analysis has shown that in multimodal therapy programmes, global patterns (gait, running, ranges of motion) are assessed locally (i.e. in a particular area of the body).

Núñez-Cortés et al. (2017) proposed a walk test, pre- and post-intervention, as an objective and valid measure of exercise tolerance and exercise capacity [280]. The timed up-and-go test (TUG) was used to evaluate dynamic equilibrium, predicting the risk of falling [281]. Lower-body muscle power was assessed using a 30-second chair-stand test. This test can be influenced by range of motion in joints [282] and balance [283] and it was used to measure lower-body muscle power based on Smith et al. [284]. The use of measurable tests deserves recognition (as emphasised repeatedly in this publication) as an indication of best practices in rehabilitation diagnosis and therapy. Nowadays, there are numerous devices and sensors for postural and functional assessment [127–131] of balance, gait, force and pressure distribution [132–139], in combination with videogrammetry and photogrammetry. In most research studies, functional diagnosis is performed out locally, rarely globally. The diagnostic methods used, despite the significant advances in measurable (computer-controlled) technologies, mainly boil down to visual assessment, palpation, analogue measurements or questionnaires. The analysis of multimodal programmes is largely based on a local assessment of the pain problem [17, 20, 44–46, 49, 51, 53, 55, 62–63, 66, 188].

The literature study revealed a significant deficit in methods for a global assessment of posture, postural patterns, gait determinants, etc. This may undermine the efficacy of the therapeutic approaches, also in the assessment of referred and latent pain.

The local approach to the problem is reflected in the research literature where diagnostic methods are tested in specific ailments and isolated body areas (as seen in both Part I and Part II of this publication). The combination of manual therapy and dry needling with pain neuroscience education is probably the preferred multimodal approach [285–288]. The multimodal programmes included in our analysis appear to confirm the above assertion, although manual therapy and dry needling were used in 3 of the 7 articles, which seems puzzling, mainly due to the fact that the existence of MTrPs was reported in all of them, while ischemic compression, dry needling, dry cupping are local methods dedicated to the elimination of active (painful) trigger points. Pérez-Palomares et al. in their

findings went as far as stating that dry needling did not have a meaningful effect in MPS therapy, compared to groups that received a multimodal therapy programme. This finding can nevertheless be challenged (as the authors admitted objectively), due to the fact that patients received a personalised therapy programme. The principle of reproducibility was therefore not complied with [248].

The patient's individual complaints should also guide the choice of rehabilitation exercises (both in the process of patient education and therapist-monitored therapy), i.e. stretching, isometric exercises, proprioceptive and postural re-education, including during activities of daily living [287, 289–293].

None of the analysed programmes explicitly described the rules for method selection, making it impossible to ascertain whether the choice of method depended on the cause of pain. The literature review has led us to conclude empirically that scientific research is focused on proving the effectiveness of specific therapeutic methods rather than entire programmes. Even if the title suggests the opposite. Each of the seven articles included in the analysis presents a well-defined methodology for administering the respective therapeutic methods, patient education, etc. This has probably had a positive impact on the findings from the studies, i.e. pain relief, functional improvement. Such an approach to the evaluation of efficacy does not yield itself to a comparative analysis of treatment outcomes.

In the majority of the articles included in the analysis, the results were analysed statistically and observations were conducted according to the “follow-up” and “intention to treat” principles. In rehabilitation procedures, it is hardly possible to fulfil the condition that the specialist should not know the patient's group allocation. Hence, double blinding in rehabilitation studies is not common. However, studies investigating the efficacy of multimodal programmes for MPS treatment need to be randomised. Given the diverse etiology, these programmes should be developed with strict grouping of patients according to the cause of pain.

Randomisation of patient groups is also relevant to the issue of pain projection and latent lesions in myofascial structures, where it is important to diagnose and treat the surrounding and distant tissues and even posture as a whole. It is also necessary to select the treatment method to address the location of the pain in structural terms, i.e. muscle stretching and strengthening, fascial stretching, joint mobilization, neural mobilization, etc. (see also the summary in Table 1, Part I of this article).

As a result, detailed diagnosis, differentiation of causes and evaluation of the general condition all have implications for the composition of multidisciplinary teams in the development of multimodal myofascial pain treatment programmes. Given that pain is often associated with high levels of stress, depression, anxiety, there is an important role on the team to be played by a psychotherapist. Marcos-Martín et al. (2018) highlighted the important role of the psychotherapist in the assessment of kinesiophobia and catastrophising [294]. Thus, their multimodal therapy programme for chronic cervico-craniofacial pain employed distraction and stress management techniques, and relaxation exercises (i.e. Yoga). The team

assembled for the study by Anderson et al. (2016) included a urologist, primarily because the pain they studied involved the pelvic floor [217]. In the other articles included in the analysis, the involvement of a multidisciplinary team was not reported. Indeed, in the studies by Marcos-Martín et al. (2018), Núñez-Cortés et al. (2017), Moustafa et al. (2018), concomitant systemic diseases were listed as an exclusion criterion [246]. Having regard to the anatomical relationships of the musculoskeletal and visceral structures, as well as biochemical and lifestyle factors, it is important that further research should include treatment efficacy analysis with the participation of an interdisciplinary team.

Results of hypothesis testing

1. (+) Multimodal treatment programmes for MPS rely on the strict methodology of scientific research work, with respect to treatment methodology, including statistical analysis of the results; (-) on the other hand, research methodology (randomisation, blinding, final estimation) is not strictly adhered to.
2. In the analysed articles, the selection of therapeutic methods to be included in a multimodal treatment programme for MPS was not based on a detailed diagnosis (measurable, reproducible, objective) of the patient's condition, including comorbidities.
3. In most of the reviewed articles a multidisciplinary team was not set up in the case of comorbidities associated with MPS.
4. The selection of therapeutic methods in the analysed multimodal programmes does not indicate explicitly the correlations between the method and the cause of pain, comorbidities, psycho-emotional state, etc.

Conclusions

1. Multimodal programmes for the treatment of musculoskeletal pain, notably MPS and MTrPs, should include a detailed and comprehensive diagnosis (structural, biochemical, psycho-emotional) which should serve as the basis for the formation of interdisciplinary rehabilitation teams.
2. Musculoskeletal diagnosis in MPS therapy, in addition to radiological assessment, should include measurable techniques of postural and functional assessment (such as pedobarography, wearable sensors, assisted anthropometry, i.e. photogrammetry, videogrammetry, etc.), aimed primarily at the ongoing assessment of posture.
3. The choice of therapeutic methods and patient education should be based on the causes of the patient's pain, taking into account systemic diseases, postural defects, lifestyle and psycho-emotional state.
4. Scientific research in multimodal treatment programmes should be carried out in randomised groups, with due attention to the methodologies of diagnostic and therapeutic procedures and group selection.

Adres do korespondencji / Corresponding author

Danuta Lietz-Kijak

E-mail: danuta.lietzkijak@gmail.com

Piśmiennictwo/ References

1. Pal US, Kumar L, Mehta G, Singh N, Singh G, Singh M, Yadav HK. Trends in management of myofascial pain. *Natl J Maxillofac Surg* 2014;5(02):109–116
2. Desai MJ, Saini V, Saini S. Myofascial pain syndrome: a treatment review. *Pain Ther* 2013;2(01):21–36.

3. Jackson T, Thomas S, Stabile V, Han X, Shotwell M, McQueen K. Prevalence of chronic pain in low-income and middle-income countries: a systematic review and metaanalysis. *Lancet*. 2015;385(suppl 2): S10.
4. Disease GBD, Injury I, Prevalence C. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study. *Lancet* 2018; 392:1789–1858.
5. Bevan S. Economic impact of musculoskeletal disorders (MSDs) on work in Europe. *Best Pract Res Clin Rheumatol* 2015; 29:356–373.
6. Salaffi F, De Angelis R, Stancati A, Garssi W. Health-related quality of life in multiple musculoskeletal conditions: a cross-sectional population based epidemiological study. II. The MAPPING study. *Clin Exp Rheumatol* 2005; 23:829–839.
7. Couto C, de Souza IC, Torres IL, Fregni F, Caumo W. Paraspinal stimulation combined with trigger point needling and needle rotation for the treatment of myofascial pain: a randomized sham-controlled clinical trial. *Clin J Pain* 2014;30(03):214–223.
8. Weissmann R.D. Considerations with regard to the biomechanics related to myofascial trigger point therapy. *Physiotherapie*, 2000; 35 (10): 13–21.
9. Sergienko S, Kalichman L. Myofascial origin of shoulder pain: a literature review. *J Bodyw Mov Ther*. 2015;19(1):91–101.
10. Bron C, de Gast A, Dommerholt J, Stegenga B, Wensing M, Oostendorp RA. Treatment of myofascial trigger points in patients with chronic shoulder pain: a randomized, controlled trial. *BMC Med*. 2011;9(1):8.
11. Worsley P, Warner M, Mottram S, Gadola S, Veeger HE, Hermens H, Morrissey D, Little P, Cooper C, Carr A, Stokes M. Motor control retraining exercises for shoulder impingement: effects on function, muscle activation, and biomechanics in young adults. *J Shoulder Elb Surg*. 2013;22(4):11–19.
12. Jafri MS. Mechanisms of myofascial pain. *Int Sch Res Notices* 2014; 2014:523924.
13. Lavelle ED, Lavelle W, Smith HS. Myofascial trigger points. *Med Clin North Am*. 2007; 91:229-239.
14. Kuan, TS, Wu CT, Chen S., Chen JT, Hong CZ. Manipulation of the cervical spine to release pain and tightness caused by myofascial trigger points. *Arch Phys Med Rehabil*. 1997;78 (9), 1042.
15. Simons DG. Understanding effective treatments of myofascial trigger points. *J. Bodywork Mov Ther*. 2002; 6: 81–88.
16. Simons DG, Travell JG, Simons LS, Travell and Simons' Myofascial Pain and Dysfunction: The Trigger Point Manual. Vol. 1. 2nd ed. Baltimore, MD: Williams & Wilkins, 1999.
17. Simons DG, Travell JG, Simons LS. Travell & Simons' Myofascial Pain and Dysfunction: The Trigger Point Manual Vol. 1: Upper Half of Body. 2nd ed. Philadelphia, PA: Lippincott Williams & Wilkins; 1998.
18. Bron C, Dommerholt J, Stegenga B, Wensing M, Oostendorp RA. High prevalence of shoulder girdle muscles with myofascial trigger points in patients with shoulder pain. *BMC Musculoskel Dis*. 2011;12(1):139.
19. Skootsky SA, Jaeger B, Oye RK. Prevalence of myofascial pain in general internal medicine practice. *West J Med*. 1989; 151:157-160.
20. Kalichman L, Vulfsons S. Dry needling in the management of musculoskeletal pain. *J Am Board Fam Med*. 2010; 23:640-646.
21. Friction JR, Kroening R, Haley D, Siegert R. Myofascial pain syndrome of the head and neck: a review of clinical characteristics of 164 patients. *Oral Surg Oral Med Oral Pathol*. 1985; 60:615-623.
22. Kietrys DM, Palombaro KM, Azzaretto E, Hubler R, Schaller B, Schluskel JM, Tucker M. Effectiveness of dry needling for upper-quarter myofascial pain: a systematic review and meta-analysis. *J Orthop Sport Phys Ther*. 2013;43 (9), 620e634.
23. Hong CZ, Simons DG. Pathophysiologic and electrophysiologic mechanisms of myofascial trigger points. *Arch Phys Med Rehabil*. 1998; 79:863-872.
24. Perez-Palomares S, Oliván-Blázquez B, Amal-Burró AM, Mayoral-Del Moral O, Gaspar-Calvo E, de-la-Torre-Beldarraín ML, López-Lapeña E, Pérez-Benito M, Ara-Loriente V, Romo-Calvo L. Contributions of myofascial pain in diagnosis and treatment of shoulder pain. A randomized control trial. *BMC Musculoskel Disord*. 2009; 10:92.
25. Travell JG, Simons DG. Myofascial Pain and Dysfunction: The Trigger Point Manual Vol.2: The Lower Extremities. Media, PA: Williams & Wilkins; 1992.
26. Cummings TM, White AR. Needling therapies in the management of myofascial trigger point pain: a systematic review. *Arch Phys Med Rehabil*. 2001; 82: 986– 992.
27. Tough EA, White AR, Cummings TM, Richards SH, Campbell JL. Acupuncture and dry needling in the management of myofascial trigger point pain: a systematic review and meta-analysis of randomised controlled trials. *Eur J Pain*. 2009; 13: 3– 10.
28. Spadoni GF, Stratford PW, Solomon PE, Wishart LR. The evaluation of change in pain intensity: a comparison of the P4 and single-item numeric pain rating scales. *J Orthop Sports Phys Ther*. 2004; 34: 187– 193.
29. Meister MR, Shivakumar N, Sutcliffe S, Spitznagle T, Lowder JL. Physical examination techniques for the assessment of pelvic floor myofascial pain: a systematic review. *Am J Obstet Gynecol*. 2018, 219(5): 497.e1-497.e13.
30. Ribeiro DC, Belgrave A, Naden A, et al. The prevalence of myofascial trigger points in neck and shoulder-related disorders: a systematic review of the literature. *BMC Musculoskel Disord*. 2018, 19(1): 252.
31. Sreby JZ. New trends in the treatment and management of myofascial pain syndrome. *Curr Pain Headache Rep*. 2010, 14(5): 346-352.
32. Furlan AD, van Tulder M, Cherkin D, Tsukayama H, Lao L, Koes B, Berman B. Acupuncture and dry-needling for low back pain: an updated systematic review within the framework of the Cochrane Collaboration. *Spine (Phila Pa 1976)*. 2005; 30:944-963.
33. Lidbeck J. Central hyperexcitability in chronic musculoskeletal pain: A conceptual breakthrough with multiple clinical implications. *Pain Res Manag*. 2002; 7 (2): 81–92.
34. Ge HY, Fernández-de-las-Peñas C, Arendt Nielsen L. Sympathetic facilitation of hyperalgesia evoked from myofascial tender and trigger points in patients with unilateral shoulder pain. *Clin. Neurophysiol*. 2006; 117: 1545–1550.
35. Munglani R. Neurobiological mechanisms underlying chronic whiplash associated pain: The peripheral maintenance of central sensitization. *J Musculoskeleal Pain*. 2000; 8: 169–178.
36. Dommerholt J. Persistent myalgia following whiplash. *Curr. Pain Headache Rep*. 2005; 9: 326–33.
37. Lucas KR, Polus BI, Rich PS. Latent myofascial trigger points: Their effect on muscle activation and movement efficiency. *J. Bodywork Mov. Ther*. 2004; 8: 160–166.
38. Hsieh YL, Kao MJ, Kuan TS, Chen SM, Chen JT, Hong CZ. Dry needling to a key myofascial trigger point may reduce the irritability of satellite MTrPs. *Am J Phys Med Rehabil*. 2007; 86:397-403.
39. Hong CZ. Considerations and recommendations regarding myofascial trigger point injection. *J Musculoskel Pain*. 1994; 2:29-59.
40. Hong CZ. Lidocaine injection versus dry needling to myofascial trigger point. The importance of the local twitch response. *Am J Phys Med Rehabil*. 1994;73:256-263.
41. Fink R. Pain assessment: the cornerstone to optimal pain management. *Proc (Bayl Univ Med Cent)* 2000;13:236–239.
42. Chandola HC, Chakraborty A. Fibromyalgia and myofascial pain syndrome-a dilemma. *Indian J Anaesth* 2009;53(05):575–581.
43. Weller JL, Comeau D, Otis JAD. Myofascial pain. *Semin Neurol*. 2018 Dec;38(6):640-643.
44. Simons DG. Do endplate noise and spikes arise from normal motor endplates? *Am J Phys Med Rehabil* 2001;80:134–40.
45. Kuan TS, Hsieh YL, Chen SM, Chen JT, Yen WC, Hong CZ. The myofascial trigger point region: correlation between the degree of irritability and the prevalence of endplate noise. *Am J Phys Med Rehabil* 2007;86:183–9.
46. Dumitru D, King JC, Stegeman DF. Endplate spike morphology: a clinical and simulation study. *Arch Phys Med Rehabil*. 1998;79:634–40.
47. Borg-Stein J, Simons DG. Focused review: myofascial pain. *Arch Phys Med Rehabil*. 2002;83(03, Suppl 1):S40–S47, S48–S49.
48. Stecco A, Gesi M, Stecco C, Stern R. Fascial components of the myofascial pain syndrome. *Curr Pain Headache Rep* 2013;17(08):352.
49. Alvarez DJ, Rockwell PG. Trigger points: diagnosis and management. *Am Fam Physician*. 2002 Feb 15;65(4):653-60.
50. Bitenc-Jasieško A, Konior K, Gonta K, Duleba M, Lietz-Kijak D. Prophylaxis of Pain and Fractures within Feet in the Course of Osteoporosis: The Issue of Diagnosing. *Pain Res Manag*. 2020 Nov 29;2020:1391026. doi: 10.1155/2020/1391026. eCollection 2020
51. Fernández-de-las-Peñas C, Alonso-Blanco C, Miangolarra JC. Myofascial trigger points in subjects presenting with mechanical neck pain: A blinded, controlled study *Man Ther*. 2007 Feb;12(1):29-33.
52. Calandre EP, Hidalgo J, García-Leiva JM, Rico-Villademoros F. Trigger point evaluation in migraine patients: An indication of peripheral sensitization linked to migraine predisposition? *Eur J Neurol* 2006; 13: 244–249.
53. Headache Classification Subcommittee of the International Headache Society: The international classification of headache disorders. *Cephalalgia* 2004; 24 (Suppl I): 9-160.
54. Hesse J, Mogelvang B, Simonsen H. Acupuncture versus metoprolol in migraine prophylaxis: A randomized trial of trigger point inactivation. *J Intern Med* 1994; 235: 451–456.
55. Fernández-de-las-Peñas C, Cuadrado ML, Gerwin RD, Pareja JA. Referred pain from the trochlear region in tension-type headache: A myofascial trigger point from the superior oblique muscle. *Headache*. 2005; 45: 731–737.
56. Fernández-de-las-Peñas C, Alonso-Blanco C, Cuadrado ML, Pareja JA. Myofascial trigger points in the suboccipital muscles in episodic tension-type headache. *Man Ther*. 2006; 11: 225–230.
57. Weiner DK, Sakamoto S, Perera S, Breuer P. Chronic low back pain in older adults: Prevalence, reliability, and validity of physical examination findings. *J Am. Geriatr Soc*. 2006; 54: 11–20.
58. Hsueh TC, Yu S, Kuan TS, Hong CZ. Association of active myofascial trigger points and cervical disc lesions. *J Formosa Med Assoc*. 1998; 97(3): 174–80.
59. Fruth SJ. Differential diagnosis and treatment in a patient with posterior upper thoracic pain. *Phys Ther*. 2006; 86: 254–268.
60. Wang CF, Chen M, Lin MT, Kuan TS, Hong CZ. Teres minor tendinitis manifested with chronic myofascial pain syndrome in the scapular muscles: A case report. *J. Musculoskeletal Pain*. 2006; 14(1): 39–43.
61. Friction JR. Etiology and management of masticatory myofascial pain. *J Musculoskeletal Pain*. 1999; 7 (1/2): 143–160.
62. Teachey WS. Otolaryngic myofascial pain syndromes. *Curr Pain Headache Rep*. 2004; 8: 457–462.
63. Skubick DL, Clasby R, Donaldson CC, Marshall WM. Carpal tunnel syndrome as an expression of muscular dysfunction in the neck. *J. Occupational Rehab.*, 1993; 3: 31–43.
64. Weiss JM. Pelvic floor myofascial trigger points: Manual therapy for interstitial cystitis and the urgency-frequency syndrome. *J Urol*. 2001; 166: 2226–223.
65. Doggweiler-Wyguil R. Urologic myofascial pain syndromes. *Curr Pain Headache Rep*. 2004; 8: 445–451.
66. Jarrell JF, Vilos GA, Allaire C, Burgess S, Fortin C, Gerwin R, Lapensee L, Lea RH, Leyland NA, Martyn P, Abu-Rafea B. Consensus guidelines for the management of chronic pelvic pain. *J Obstet Gynaecol Can*. 2005; 27: 869–887.

67. Jarrell J. Myofascial dysfunction in the pelvis. *Curr Pain Headache Rep.* 2004; 8: 452–456.
68. Treede RD, Rief W, Barke A, Aziz Q, Bennett MI, Benoliel R, Cohen M, Evers S, Finnerup NB, First MB, Giamberardino MA, Kaasa S, Kosek E, Lavand'homme P, Nicholas M, Perrot S, Scholz J, Schug S, Smith BH, Svensson P, Vlaeyen JW, Wang SJ. A classification of chronic pain for ICD-11. *Pain.* 2015; 156: 1003– 1007.
69. Lavand'homme P, Thienpont E. Pain after total knee arthroplasty: a narrative review focusing on the stratification of patients at risk for persistent pain. *Bone Joint J.* 2015; 97-B: 45– 48.
70. Fuzier R, Serres I, Bourrel R, Palmaro A, Montastruc JL, Lapeyre-Mestre M. Analgesic drug consumption increases after knee arthroplasty: a pharmacoepidemiological study investigating postoperative pain. *Pain.* 2014;155:1339-1345.
71. Kurtz SM, Ong KL, Lau E, Bozic KJ. Impact of the economic downturn on total joint replacement demand in the United States: updated projections to 2021. *J Bone Joint Surg Am.* 2014;96:624-630.
72. Nice DA, Riddle, DL, Lamb RL, Mayhew TP, Rucker K. Interrater reliability of judgments of the presence of trigger points in patients with low back pain. *Arch Phys Med Rehabil.* 1992;73(10):893e898.
73. Wolfe F, Simons DG, Friction J, Bennett RM, Goldenberg DL, Gerwin R, Hathaway D, McCain GA, Russell JJ, Sanders HO. The fibromyalgia and myofascial pain syndromes: a preliminary study of tender points and trigger points in persons with fibromyalgia, myofascial pain syndrome and no disease. *J Rheumatol.* 1992;19(6):944e95.
74. Hua NK, Van der Does E. The occurrence and inter-rater reliability of myofascial trigger points in the quadratus lumborum and gluteus medius: a prospective study in non-specific low back pain patients and controls in general practice. *Pain.* 1994;58 (3), 317e323.
75. Ge HY, Arendt-Nielsen L. Latent myofascial trigger points. *Curr Pain Headache Rep.* 2011; 15(5): 386-392.
76. Malanga GA, Cruz Colon EJ. Myofascial low back pain: a review. *Phys Med Rehabil Clin N Am.* 2010, 21(4):711-724.
77. Dommerholt PT, Bron C, Franssen J. Myofascial Trigger Points: An Evidence-Informed Review. *The Journal of Manual & Manipulative Therapy,* 2006, 14 (4), 203–2216.
78. Ruiz-Sáez M, Fernández-de-las-Peñas C, Blanco CR, Martínez-Segura R, García-León R. Changes in pressure pain sensitivity in latent myofascial trigger points in the upper trapezius muscle after a cervical spine manipulation in pain-free subjects. *J Manipulative Physiol Ther.* 2007, 30(8): 578-583.
79. Henneman E, Somjen G, Carpenter DO. Excitability and inhibitory of motoneurons of different sizes. *J Neurophysiol.* 1965; 28: 599–620.
80. Hägg GM. The Cinderella Hypothesis. in: Johansson H, Windhorst U, Djupsjöbacka M, Passatore M. Chronic Work - Related Myalgia. Gävle, Sweden: Gävle University Press. 2003; 127–132.
81. Febbraio MA, Pedersen BK. Contraction-induced myokine production and release: Is skeletal muscle an endocrine organ? *Exerc Sport Sci Rev.* 2005; 33 (3): 114–119.
82. Pedersen BK, Febbraio M. Muscle-derived interleukin-6: A possible link between skeletal muscle, adipose tissue, liver, and brain. *Brain Behav Immun.* 2005; 19: 371–376.
83. Gerwin RD, Dommerholt J, Shah J. An expansion of Simons' integrated hypothesis of trigger point formation. *Curr Pain Headache Rep.* 2004; 8: 468–475.
84. Graven-Nielsen T, Arendt-Nielsen L. Induction and assessment of muscle pain, referred pain, and muscular hyperalgesia. *Curr Pain Headache Rep.* 2003; 7: 443–451.
85. Shah JP, Phillips TM, Danoff JV, Gerber LH. An in-vivo micro-analytical technique for measuring the local biomechanical milieu of human skeletal muscle. *J Appl Physiol.* 2005; 99: 1980–1987.
86. Henriksson KG, Bengtsson A, Lindman R, Thormell LE. Morphological changes in muscle in fibromyalgia and chronic shoulder myalgia. in: Vörfy H, Merskey H. Progress in Fibromyalgia and Myofascial Pain. Elsevier. 1993; 61–73.
87. Hägg GM. Human muscle fibre abnormalities related to occupational load. *Eur J Appl Physiol.* 2000; 83(2–3):159–165.
88. Kadi F, Waling K, Ahlgren C, Sundelin G, Holmner S, Butler-Browne GS, Thormell LE. Pathological mechanisms implicated in localized female trapezius myalgia. *Pain.* 1998; 78: 191–196.
89. Kadi F, Hagg G, Hakansson R, Holmner S, Butler-Browne GS, Thormell LE. Structural changes in male trapezius muscle with work related myalgia. *Acta Neuropathol.* 1998; 95: 352–360.
90. Larsson B, Björk J, Kadi F, Lindman R, Gerde B. Blood supply and oxidative metabolism in muscle biopsies of female cleaners with and without myalgia. *Clin J Pain.* 2004;20: 440–446.
91. Bruce E. Myofascial pain syndrome: early recognition and comprehensive management. *AAOHN J.* 1995;43(09):469–474.
92. Rubin D. Myofascial trigger point syndromes: an approach to management. *Arch. Phys Med Rehabil.* 1981;62 (3),107e110.
93. Travell JG, Simons DG. Myofascial Pain and Dysfunction: The Trigger Point Manual Vol. 2: The Lower Extremities. Media, PA: Williams & Wilkins; 1992.
94. Baker BA. The muscle trigger: Evidence of overload injury. *J. Neurol. Orthop Med Surg.* 1986; 7:35–44.
95. Schuller E., Eisenmenger W., Brier G.: Whiplash injury in low speed car accidents. *J Musculoskeletal Pain.* 2000; 8(1/2):55–67.
96. Fernández-de-las-Peñas C, Fernández-Camero J, Miangolarra-Page JC. Musculoskeletal disorders in mechanical neck pain: Myofascial trigger points versus cervical joint dysfunction. *J. Musculoskeletal Pain.* 2005;13(1):27–35.
97. Kuan, TS. Current studies on myofascial pain syndrome. *Curr. Pain Headache Rep.* 2009; 13(5), 365e369.
98. Fernández-de-las-Peñas C, Alonso-Blanco C, Alguacil-Diego IM, Miangolarra-Page JC. Myofascial trigger points and posteroanterior joint hypomobility in the mid-cervical spine in subjects presenting with mechanical neck pain: A pilot study. *J Manipulative Ther.* 2006; 14: 88–94.
99. Bogduk N, Simons DC. Neck pain: Joint pain or trigger points. in: Vörfy H, Merskey H. Progress in Fibromyalgia and Myofascial Pain. Elsevier. 1993;267–273.
100. Padamsee M, Mehta N, White G.E: Trigger point injection: A neglected modality in the treatment of TMJ dysfunction *J Pedod.* 1987;12(1):72–92.
101. McNulty WH, Gevirtz RN, Hubbard DR, Berkoff GM. Needle electromyographic evaluation of trigger point response to a psychological stressor. *Psychophysiology.* 1994;31(3), 313e316.
102. Shah JP, Phillips TM, Danoff JV, Gerber LH. A novel microanalytical technique for assaying soft tissue demonstrates significant quantitative biochemical differences in 3 clinically distinct groups: normal, latent, and active. *Arch. Phys Med Rehabil.* 2003;84 (9), E4.
103. Vázquez-Delgado E, Cascos-Romero J, Gay-Escoda C. Myofascial pain associated to trigger points: a literature review. Part 2: differential diagnosis and treatment. *Med Oral Patol Oral Cir Bucal.* 2010 Jul 1;15(4):e639-43
104. Geri T, Gizzi L, Di Marco A, Testa M. Endplate potentials are common to midfiber myofascial trigger points. *Am J Phys Med Rehabil.* 2002;81(3), 212e222.
105. Gerwin RD. A review of myofascial pain and fibromyalgia – factors that promote their persistence. *Acupunct Med.* 2005;3:121-134.
106. Dommerholt J. Muscle pain syndromes. in: Cantu RI, Grodin AJ. Myofascial Mumpiaifion. Gaithersburg MD. Aspen, 2001;93–140.
107. Weiner DK, Schmader KE. Postherpetic pain: More than sensory neuralgia? *Pain Med.* 2006; 7: 243–249.
108. Chen SM, Chen JT, Kuan TS, Hong CZ. Myofascial trigger points in intercostal muscles secondary to herpes zoster infection of the intercostal nerve. *Arch Phys Med Rehabil.* 1998; 79: 336–338.
109. Dommerholt J. Complex regional pain syndrome. Part 1: History, diagnostic criteria and etiology. *J Bodywork Mov Ther.* 2004;8: 167–177.
110. Rashid Q, Galer BS. Proximal myofascial dysfunction in complex regional pain syndrome: A retrospective prevalence study. *Clin J Pain.* 1999;15: 151–153.
111. Prateepavanich P, Kupniratsaikul VC. The relationship between myofascial trigger points of gastrocnemius muscle and nocturnal calf cramps. *J. Med. Assoc. Thailand.* 1999; 82:451–459.
112. Longbottom J. A case report of postulated "Barre Liéou syndrome." *Acupunct. Med.,* 2005; 23:34–38.
113. Stellan A. Neurogenic pruritus: An unrecognized problem? A retrospective case series of treatment by acupuncture. *Acupunct. Med.,* 2002; 20: 186–190
114. Han SC, Harrison P. Myofascial pain syndrome and trigger-point management. *Reg Anesth Pain Med.* 1997;22(1):89e101.
115. Pedro-Botet J, Climent E, Benais D. Muscle and statins: from toxicity to the nocebo effect. *Expert Opin Drug Saf.* 2019 Jul;18(7):573-579
116. Andres E, Loukili NH, Noel E, Kaltenbach G, Perrin AE, Noblet-Dick M, Maloel F, Schlienger JL. Vitamin B12 (cobalamin) deficiency in elderly patients. *CMAJ.* 2004;171: 251–259.
117. Pruthi RK, Tefferi A. Pernicious anemia revisited. *Mayo Clin Proc.* 1994;69: 144–150.
118. Plotnikoff GA, Quigley JM. Prevalence of severe hypovitaminosis D in patients with persistent, nonspecific musculoskeletal pain. *Mayo Clin Proc.* 2003;78:1463–1470.
119. Dommerholt J, Issa T. Differential diagnosis: Myofascial pain. in: Chaitow L. Fibromyalgia Syndrome: A Practitioner's Guide to Treatment. Edinburgh, UK: Churchill Livingstone, 2003;149–177.
120. Gerwin RD, Shannon S, Hong CZ, Hubbard D, Gevirtz R. Interrater reliability in myofascial trigger point examination. *Pain.* 1997;69(1–2):65–73.
121. Sciotti VM, Mittak ML, DiMarco L, Ford LM, Plezbert J, Santipadri E, Wigglesworth J, Ball K. Clinical precision of myofascial trigger point location in the trapezius muscle. *Pain.* 2001; 93(3): 259–266.
122. Partanen JV, Ojala TA, Arokoski JP. Myofascial syndrome and pain: a neurophysiological approach. *Pathophysiology.* 2010;17(1):19-28.
123. Sikdar S, Shah JP, Gebreab T, Yen RH, Gilliams E, Danoff J, Gerber LH. Novel applications of ultrasound technology to visualize and characterize myofascial trigger points and surrounding soft tissue. *Arch Phys Med Rehabil.* 2009;90(11):1829-1838.
124. Dibai-Filho AV, Guirio EC, Ferreira VT, Brandino HE, Vaz MM, Guirio RR. Reliability of different methodologies of infrared image analysis of myofascial trigger points in the upper trapezius muscle. *Braz J Phys Ther.* 2015;19(2):122-128
125. Chen Q, Bensamoun S, Basford JR, Thompson JM, An KN. Identification and quantification of myofascial taut bands with magnetic resonance elastography. *Arch Phys Med Rehabil.* 2007;88(12):1658-1661.
126. Shah JP, Danoff JV, Desai MJ, Parikh S, Nakamura LY, Phillips TM, Gerber LH. Biochemicals associated with pain and inflammation are elevated in sites near to and remote from active myofascial trigger points. *Arch Phys Med Rehabil.* 2008;89(1):16-23.
127. Kuusma M, Karppinen J, Niinimäki J, Kurunlahti M, Haapea M, Vanharanta H, Tervonen O. A three years follow-up of lumbar spine endplate (Modic) changes. *Spine.* 2006;31(15):1714.
128. Lewis CL, Laudicina NM, Khuu A, Loverro KL. The Human Pelvis: Variation in structure and function during gait. *Anat Rec (Hoboken).* 2017 Apr; 300(4): 633–642.
129. Bugané F, Benedetti MG, Casadio G, Attala S, Biagi F, Manca M, Leardini A. Estimation of spatial-temporal gait parameters in level walking base on a single accelerometer: validation on normal subjects by standard gait analysis. *Comput Methods Programs Biomed* 2012;108(1):29–37.
130. Bonnet V, Mazzà C, McCamley J, Cappozzo A. Use of weighted Fourier linear combiner filters to estimate lower trunk 3D orientation from gyroscope sensors data. *J Neuroeng Rehabil* 2013;10:29.
131. Mazzà C, Donati M, McCamley J, Picerno P, Cappozzo A: An optimized Kalman filter for the estimation of trunk orientation from inertial sensors data during treadmill walking. *Gait Posture* 2012;35:138–142.
132. O'Leary CB, Cahill CR, Robinson AW, Barnes MJ, Hong J. A systematic review: The effects of podiatric deviations on nonspecific chronic low back pain. *Journal of Back and Musculoskeletal Rehabilitation.* Vol. 26, no. 2, pp. 2013, 117-123.
133. Akbari M, Karimi H, Farahani H, Faghizadeh S. Balance problems after unilateral lateral ankle sprains; Rehabilitation Research & Development, 2006;43(7): 819-24.

134. Bennett PJ, Patterson C, Wearing S, Baglioni T. Development and Validation of a questionnaire designed to measure foot-health status; American Podiatric Medical Association, 1998; 88: 419-428.
135. Błaszczyk JW, Michalski A. Ageing and postural stability; Studies in Physical Culture and Tourism, 2006; 13: 11-14.
136. Herrington L, Hatcher J, Hatcher A, McNicholas MA. Comparison of Star Excursion Balance Test reach distances between ACL deficient patients and asymptomatic controls. *Knee*. 2009;16(2):149-52.
137. Skopljak A, Mufic M, Sukalo A, Masic I, Zunic L. Pedobarography in Diagnosis and Clinical Application, *Acta Inform Med*. 2014 Dec; 22(6): 374–378.
138. Choi YR, Lee HS, Kim DE, Lee DH, Kim JM, Ahn JY. The Diagnostic Value of Pedobarography, *Orthopedics*. 2014;37(12): e1063-e1067.
139. Bitenc-Jasieko A, Konior K, Lietz-Kijak D. Meta-Analysis of Integrated Therapeutic Methods in Noninvasive Lower Back Pain Therapy (LBP): The Role of Interdisciplinary Functional Diagnostics, *Pain Res Manag*. 2020 Mar 19;2020:3967414.
140. Deng W, Papavasileiou I, Qiao Z, Zhang W, Lam KY, Han S. Advances in Automation Technologies for Lower Extremity Neurorehabilitation: A Review and Future Challenges, *IEEE Rev Biomed Eng*. 2018;11:289-305.
141. Tavares C, Domingues MF, Frizzera-Neto A, Leite T, Leitão C, Alberto N, Marques C, Radwan A, Rocon E, André P, Antunes P. Gait Shear and Plantar Pressure Monitoring: A Non-Invasive OFS Based Solution for e-Health Architectures, *Sensors (Basel)*. 2018 Apr 25;18(5).
142. Ferber R, Osis ST, Hicks JL, Delp SL. Gait biomechanics in the era of data science, *J Biomech*. 2016 Dec 8;49(16):3759-3761.
143. Souchard P. Reeducation posturale globale. Elsevier Masson SAS. Tous droits réservés. 2011; 39-42.
144. Lawson T, Morrison A, Blaxland S, Wenman M, Schmidt CG, Hunt MA. Laboratory-based measurement of standing balance in individuals with knee osteoarthritis: A systematic review. *Clin Biomech*. Elsevier Ltd; 2015;30: 330–342.
145. Geldhof E, Cardon G, De Bourdeaudhuij I, Danneels L, Coorevits P, Vanderstraeten G, De Clercq D. Static and dynamic standing balance: Test-retest reliability and reference values in 9 to 10-year-old children. *Eur J Pediatr*. 2006;165: 779–786.
146. Scoppa F, Capra R, Gallamini M, Shiffer R. Clinical stabilometry standardization. Basic definitions—Acquisition interval—Sampling frequency. *Gait Posture*. Elsevier B.V. 2013;37: 290–292.
147. Fernández-de-las-Peñas C, Alonso-Blanco C, Cuadrado ML, Gerwin RD, Pareja JA. Trigger points in the suboccipital muscles and forward head posture in tension-type headache. *Headache*. 2006; 46: 454–460.
148. Aguilera FJ, Martín DP, Masanet RA, Botella AC, Soler LB, Morell FB. Immediate effect of ultrasound and ischemic compression techniques for the treatment of trapezius latent myofascial trigger points in healthy subjects: a randomized controlled study. *J Manip Physiol Therapeut*. 2009;32 (7), 515e520.
149. Fernández-de-las-Peñas C, Campo, MS, Camero, JF, Page JCM. Manual therapies in myofascial trigger point treatment: a systematic review. *J Bodyw Mov Ther*. 2005;9(1), 27e34.
150. Cotchett MP, Munteanu SE, Landorf KB. Effectiveness of trigger point dry needling for plantar heel pain: a randomized controlled trial. *Phys Ther*. 2014 Aug;94(8):1083-94.
151. De Meulemeester KE, Castellein B, Coppeters I, Barbe T, Cools A, Cagnie B. Comparing trigger point dry needling and manual pressure technique for the management of myofascial neck/shoulder pain: a randomized clinical trial. *J Manip Physiol Therapeut*. 2017;40 (1), 11e20.
152. Martín-Pintado Zugasti A, Rodríguez-Fernández ÁL, García-Muro F, López-López A, Mayoral O, Mesa-Jiménez J, Fernández-Camero J. Effects of spray and stretch on postneedling soreness and sensitivity after dry needling of a latent myofascial trigger point. *Arch Phys Med Rehabil*. 2014;95(10): 1925-1932.
153. Charles D, Hudgins T, MacNaughton J, Newman E, Tan J, Wigger M. A systematic review of manual therapy techniques, dry cupping and dry needling in the reduction of myofascial pain and myofascial trigger points. *J Bodyw Mov Ther*. 2019 Jul;23(3):539-546. *J Altern Complement Med*. 2020 Apr;26(4):291-299.
154. Dayanir IO, Birinci T, Kaya Mutlu E, Akcetin MA, Akdemir AO. Comparison of Three Manual Therapy Techniques as Trigger Point Therapy for Chronic Nonspecific Low Back Pain: A Randomized Controlled Pilot Trial.
155. Dardziński JA, Ostrov BE, Hamann LS. Myofascial pain unresponsive to standard treatment: successful use of a strain and counterstrain technique with physical therapy. *J Clin Rheumatol*. 2000, 6(4): 169-174.
156. Butttagat V, Eungpinichpong W, Chatchawan U, Kharmwan S. The immediate effects of traditional Thai massage on heart rate variability and stress-related parameters in patients with back pain associated with myofascial trigger points. *J Bodyw Mov Ther*. 2011;15(1): 15-23.
157. Moraska A, Chandler C. Changes in clinical parameters in patients with Tension-type Headache Following Massage Therapy: a pilot study. *J Man Manip Ther* 2008, 16(2): 106-112.
158. Vaughan B, Morrison T, Buttigieg D, Macfarlane C, Fryer G. Approach to low back pain - osteopathy. *Aust Fam Physician*. 2014;43: 197-19.
159. Franke H, Fryer G, Ostelo R, Kamper S. Muscle energy technique for non-specific low-back pain. A Cochrane systematic review. *International Journal of Osteopathic Medicine*, Vol. 20. 2016; 41 – 52.
160. Hanten WP, Olson SL, Butts NL, Nowicki AL. Effectiveness of a home program of ischemic pressure followed by sustained stretch for treatment of myofascial trigger points. *Phys Ther*. 2000;80(10): 997-1003.
161. Bindra S, Kumar M, Singh P, Singh J. A study on the efficacy of Muscle Energy Technique as compared to conventional therapy in chronic low back pain due to sacroiliac joint dysfunction. *Indian Journal of Physiotherapy and Occupational Therapy*. 2012;6(1):200 3.
162. Dhinkaran M, Sareen A, Arora T. Comparative analysis of Muscle Energy Technique and conventional physiotherapy in treatment of sacroiliac joint dysfunction. *Indian Journal of Physiotherapy and Occupational Therapy* 2011;5(4):127 30.
163. Furlan AD, Yazdi F, Tsertsvadze A, et al. A systematic review and meta-analysis of efficacy, cost-effectiveness, and safety of selected complementary and alternative medicine for neck and low-back pain. *Evid Based Complement Alternat Med*. 2012; 953139.
164. Bronfort G, Haas M, Evans RL, Bouter LM. Efficacy of spinal manipulation and mobilization for low back pain and neck pain: a systematic review and best evidence synthesis. *Spine J*. 2004; 4:335–56.
165. Kogure A, Kotani K, Katada S, Takagi H, Kamikozuru M, Isaji T, Hakata S. A Randomized, Single-Blind, Placebo-Controlled Study on the Efficacy of the Arthrokinematic Approach-Hakata Method in Patients with Chronic Nonspecific Low Back Pain December 8. 2015; PLoS ONE 10(12):e0144325
166. Hakata S, Sumita K, Katada S. Efficacy of AKA-Hakata method in the treatment of acu-te lumbago. *Manuelle Medizin*. 2005;43: 19–24
167. Moutzouri M, Billis E, Strimpakos N, Kottika P, Oldham JA. The effects of the Mulligan Sustained Natural Apophyseal Glide (SNAG) mobilisation in the lumbar flexion range of asymptomatic subjects as measured by the Zebris CMS20 3-D motion analysis system. *BMC Musculoskeletal Disorders*. 2008; 9:131.
168. May J, Krzyzanowicz R, Nasypany A, Baker R, Seegmiller J. Muligan Concept Use and Clinical Profile From the Perspective of American Certified Muligan Practitioners, *Journal of Sport Rehabilitation*, 2015 Nov.; 24 (4): 337 – 341.
169. Hussien HM, Abdel-Raouf NA, Kattabei OM, Ahmed HH. Effect of Mulligan Concept Lumbar SNAG on Chronic Nonspecific Low Back Pain. *J Chiropr Med*. 2017 Jun; 16(2): 94–102.
170. Konstantinou K, Foster N, Rushton A, Baxter D, Wright C, Breen A. Flexion mobilizations with movement techniques: the immediate effects on range of movement and pain in subjects with low back pain. *J Manipulative Physiol Ther*. 2007 Mar-Apr;30(3):178-85
171. Scrimshaw SV, Maher CG.. Randomized controlled trial of neural mobilization after spinal surgery. *Spine (Phila Pa 1976)* 2001;26(24):2647–2652.
172. Shacklock M. Neural mobilization: a systematic review of randomized controlled trials with an analysis of therapeutic efficacy. *J Man Manip Ther*. 2008;16(1):23–24.
173. Casato G, Stecco C, Busin R. Role of fasciae in nonspecific low back pain, *Eur J Transl Myol*. 2019Aug 6;29(3):8330.
174. Branchini M, Lopopolo F, Andreoli E, Loreti I, Marchand AM, Stecco A. Fascial Manipulation for chronic aspecific low back pain: a single blinded randomized controlled trial, Ver 2. *F1000Res*. 2015 Nov 3;4:1208.
175. Stecco C, Porzionato A, Macchi V, Stecco A, Vígato E, Parenti A, Delmas V, Aldegheri R, De Caro R.. The expansions of the pectoral girdle muscles onto the brachial fascia: morphological aspects and spatial disposition. *Cells Tissues Organs*. 2007;188: 320-9.
176. Masi AT, Hannon JC. Human resting muscle tone (HRMT): narrative introduction and modern concepts; *J Bodyw Mov Ther* 2008;12(4):320-32.
177. Wilke J, Krause F, Vogt L, Banzer W. What is evidence-based about myofascial chains: a systematic review. *Archives of physical medicine and rehabilitation*; *Arch Phys Med Rehabil*. 2016;97(3):454-61.
178. Pardehshenas H, Maroufi N, Sanjari M, Parnianpour M, Levin SM. Lumbopelvic muscle activation patterns in three stances under graded loading conditions: Proposing a tensegrity model for load transfer through the sacroiliac joints; *J Bodyw Mov Ther*. 2014;18(4):633-42.
179. Kassolik K, Jaskólska A, Kisiel-Sajewicz K, Marusiak J, Kawczyński A, Jaskólski A. Tensegrity principle in massage demonstrated by electro- and mechanomyography. *J Bodyw Mov Ther*. 2009;13(2):164-70.
180. Piron A. The tensegrity concept applied to the laryngeal biodynamics. *Rev Laryngol Otol Rhinol*. 2007;128(5):273-8.
181. Stecco C, Stern R, Porzionato A, Macchi V, Masiero S, Stecco A, De Caro R. Hyaluronan within fascia in the etiology of myofascial pain. *Surg Radiol Anat*. 2011;33:891-6.
182. Langevin HM, Fox JR, Koptuch C, Badger GJ, Greenan-Naumann AC, Bouffard NA, Konofagou EE, Lee WN, Triano JJ, Henry SM. Reduced thoracolumbar fascia shear strain in human chronic low back pain. *BMC Musculoskelet Disord*. 2011;12:203.
183. Kassolik K, Andrzejewski W. Tensegration massage. *Physiotherapy*. 2010;18(1): 67-72.
184. Ajimsha MS, Binsu D, Chithra S. Effectiveness of Myofascial release in the management of chronic low back pain in nursing professionals. *J Bodyw Mov Ther*. 2014;18:273-81.
185. Twomey L, Taylor J. Flexion creep deformation and hysteresis in the lumbar vertebral column. *Spine*. 1982;7(2): 116-122.
186. Ghanbari A, Rahimjaberi A, Mohamadi M, et al. The effect of trigger point management by positional release therapy on tension type headache. *NeuroRehabilitation*. 2012;30(4): 333-339.
187. Iaroshkevskiy OA, Morozova OG, Logvinenko AV, Lypynska YV. Non-pharmacological treatment of chronic neck-shoulder myofascial pain in patients with forward head posture, *Wiad Lek*. 2019;72(1):84-88.
188. Kalichman L, Ben David C. Effect of self-myofascial release on myofascial pain, muscle flexibility, and strength: A narrative review. *J Bodyw Mov Ther*. 2017 Apr;21(2):446-451.

189. Castro Sánchez AM, García López H, Fernández Sánchez M, Pérez Mármol JM, Aguilar-Ferrández ME, Luque Suárez A, Matarán Peñarocha GA. Improvement in clinical outcomes after dry needling versus myofascial release on pain pressure thresholds, quality of life, fatigue, pain intensity, quality of sleep, anxiety, and depression in patients with fibromyalgia syndrome. *Disabil Rehabil.* 2019 Sep;41(19):2235-2246.
190. Castro-Sánchez AM, Matarán-Peñarocha GA, Arroyo-Morales M, Saavedra-Hernández M, Fernández-Sola C, Moreno-Lorenzo C. Effects of myofascial release techniques on pain, physical function, and postural stability in patients with fibromyalgia: a randomized controlled trial. *Clin Rehabil.* 2011 Sep;25(9):800-13.
191. Franke H, Franke JD, Fryer G. Osteopathic manipulative treatment for nonspecific low back pain: a systematic review and meta-analysis. *BMC Musculoskelet Disord.* 2014; 15:286.
192. Mejuto-Vázquez MJ, Salom-Moreno J, Ortega-Santiago R, Truyols-Domínguez S, Fernández-de-Las-Peñas C. Short-term changes in neck pain, widespread pressure pain sensitivity, and cervical range of motion after the application of trigger point dry needling in patients with acute mechanical neck pain: a randomized clinical trial. *J Orthop Sport Phys Ther.* 2014;44 (4), 252e260.
193. Dunning J, Butts R, Mourad F, Young I, Flannagan S, Perreault T. Dry needling: a literature review with implications for clinical practice guidelines. *Phys. Ther Rev.* 2014;19 (4), 252e265.
194. Dunning J, Butts R, Henry N, Mourad F, Brannon A, Rodriguez H, Young I, Arias-Buría JL, Fernández-de-Las-Peñas C. Electrical dry needling as an adjunct to exercise, manual therapy and ultrasound for plantar fasciitis: A multi-center randomized clinical trial. *PLoS One.* 2018 Oct 31;13(10):e0205405.
195. Gattie E, Cleland JA, Snodgrass S. The effectiveness of trigger point dry needling for musculoskeletal conditions by physical therapists: a systematic review and meta-analysis. *J. Orthop. Sport Phys Ther.* 2017;47 (3), 133e149.
196. Srbely JZ, Dickey JP, Lee D, Lowerison M. Dry needle stimulation of myofascial trigger points evokes segmental anti-nociceptive effects. *J Rehabil Med.* 2010;42 (5), 463e468.
197. Baldry PE. *Myofascial Pain and Fibromyalgia Syndromes: A Clinical Guide to Diagnosis and Management.* Edinburgh, UK: Churchill Livingstone; 2001
198. Calixtre LB, Moreira RF, Franchini GH, Albuquerque-Sendin F, Oliveira AB. Manual therapy for the management of pain and limited range of motion in subjects with signs and symptoms of temporomandibular disorder: a systematic review of randomised controlled trials.
199. Hong CZ. Myofascial trigger point injection. *Crit Rev Phys Rehabil Med.* 1993; 5: 203– 217.
200. Markowski A, Sanford S, Pikowski J, Fauvell D, Cimino D, Caplan S. A pilot study analyzing the effects of Chinese cupping as an adjunct treatment for patients with subacute low back pain on relieving pain, improving range of motion, and improving function. *J Altern Complement Med.* 2014;20(2),113e117.
201. Kenzo K, Wallis J, Kase T. *Clinical Therapeutic Applications of the Kinesio Taping Method.* 2nd edition. Ken Ikai Co. Ltd., 2003.
202. Abbasi S, Rohani-Shirazi Z, Shokri E, García-Muro San José F. The effect of Kinesio Taping on postural control in subjects with non-specific chronic low back pain. *J Bodyw Mov Ther.* 2018 Apr;22(2):487-492.
203. Neumann, Donald A. *Kinesiology of the Musculoskeletal System: Foundations for Physical Rehabilitation.* Mosby, 2002.
204. Toprak Celenay S, Ozer Kaya D. Immediate effects of kinesio taping on pain and postural stability in patients with chronic low back pain. *J Bodyw Mov Ther.* 2019 Jan;23(1):206-210.
205. Bernardelli RS, Scheeren EM, Fuentes Filho AR, Pereira PA, Gariba MA, Moser ADL, Bichinho GL. Effects of Kinesio Taping on postural balance in patients with low back pain, a randomized controlled trial. *J Bodyw Mov Ther.* 2019 Jul;23(3):508-514.
206. Jeon JH, Jung YJ, Lee JY, Choi JS, Mun JH, Park WY, Seo CH, Jang KU. The effect of extracorporeal shock wave therapy on myofascial pain syndrome. *Ann Rehabil Med.* 2012; 36(5): 665-674.
207. Gleitz M, Hornig K. Trigger points – diagnosis and treatment concepts with special reference to extracorporeal shockwaves (article in German). *Orthopade.* 2012;41(2): 113-125.
208. Smania N, Corato E, Fiaschi A, Pietropoli P, Aglioti SM, Tinazzi M. Therapeutic effects of peripheral repetitive magnetic stimulation on myofascial pain syndrome. *Clin Neurophysiol.* 2003;114(2): 350-358.
209. Thomas AW, Graham K, Prato FS, McKay J, Forster PM, Moulin DE, Chari S. A randomized, double-blind, placebo-controlled clinical trial using a low-frequency magnetic field in the treatment of musculoskeletal chronic pain. *Pain Res Manag.* 2007; 12(4): 249-258.
210. Panagos A, Jensen M, Cardenas DD. Treatment of myofascial shoulder pain in the spinal cord injured population using static magnetic fields: a case series. *J Spinal Cord Med.* 2004; 27(2): 138-142.
211. Cheng N, Van Hoof H, Bockx E, Hoogmartens MJ, Mulier JC, De Dijkster FJ, Sansen WM, De Loecker W. The effect of electric currents on ATP generation, protein synthesis and membrane transport of rat skin. *Clin Orthop Relat Res.* 1982;171: 264-272.
212. Gorenberg M, Schwartz K. Imaging-guided hyperstimulation analgesia in low back pain. *J Pain Res.* 2013, 6: 487-491.
213. Simons DG, Travell J. Myofascial trigger points: A possible explanation. *Pain,* 1981; 10: 106–109.
214. Gam AN, Warming S, Larsen LH, et al. Treatment of myofascial trigger points with ultrasound combined with massage and exercise – a randomised controlled trial. *Pain.* 1998; 77(1): 73-79.
215. Draper DO, Mahaffey C, Kaiser D, et al. Thermal ultrasound decreases tissue stiffness of trigger points in upper trapezius muscles. *Physiother Theory Pract* 2010, 26(3): 167-172
216. Schwagerus L, Dörner K, Bender S, Müller K, Bitterlich N, Rothkötter HJ, Günther O. Myofascial chronic pelvic pain in women: A retrospective evaluation of the effects of specific diagnostics and therapy. *Schmerz.* 2020 Oct;34(5):388-399.
217. Anderson RU, Wise D, Sawyer T, Nathanson BH, Nevin Smith J. Equal Improvement in Men and Women in the Treatment of Urologic Chronic Pelvic Pain Syndrome Using a Multi-modal Protocol with an Internal Myofascial Trigger Point Wand, Appl Psychophysiol Biofeedback. 2016;41:215–218. Cherkin DC, Deyo RA, Battla MC, Street JH, Hund M, Barlow W. A comparison of Physical therapy chiropractic manipulation or an educational booklet for the treatment of low back pain. *New Eng J Med.* 1998;339(15):1021–1029.
219. Machado LA, de Souza MS, Ferreira PH, Ferreira ML. The McKenzie method for low back pain: a systematic review of the literature with a meta-analysis approach. *Spine.* 2006; 31:254–262.
220. Hides JA, Stanton WR, McMahon S, Sims K, Richardson CA. Effect of stabilization training on multifidus muscle cross-sectional area among young elite cricketers with low back pain. *Journal of Orthopaedic & Sports Physical Therapy.* 2008;38(3):101–108.
221. Bussing A, Ostermann T, Ludtke R, Michalsen A. Effects of yoga interventions on pain and pain associated disability: a meta analysis. *Journal of Pain.* 2012;13(1):1–9.
222. Nambi GS, Inbasekaran D, Khuman R, Devi S, Shanmuganathan, Jagannathan K. Changes in pain intensity and health related quality of life with Iyengar yoga in nonspecific chronic low back pain: a randomized controlled study. *Int J Yoga.* 2014; 7:48-53.
223. Cramer H, Lauche R, Haller H, Dobos G. A systematic review and meta-analysis of yoga for low back pain. *Clin J Pain.* 2013;29(5):450–460.
224. Dierckx R, Bron C, Dorrestijn O, et al. Guideline for diagnosis and treatment of subacromial pain syndrome: a multidisciplinary review by the Dutch Orthopaedic Association. *Acta Orthop.* 2014;85(3):314-322.
225. Jayaseelan DJ, Moats N, Ricardo CR. Rehabilitation of proximal hamstring tendinopathy utilizing eccentric training, lumbopelvic stabilization, and trigger point dry needling: 2 case reports. *J Orthop Sports Phys Ther.* 2014;44(3):198-205.
226. Dembowski SC, Westrick RB, Zylstra E, Johnson MR. Treatment of hamstring strain in a collegiate pole-vaulter integrating dry needling with an eccentric training program: a resident's case report. *Int J Sports Phys Ther.* 2013;8(3):328-339.
227. Bahçecioglu Turan G. The effect of yoga on respiratory functions, symptom control and life quality of asthma patients: A randomized controlled study. *Complement Ther Clin Pract.* 2019 Oct 28;101070.
228. Cramer H, Ward L, Steel A, Lauche R, Dobos G, Zhang Y. Prevalence, patterns, and predictors of yoga use: results of a U.S. nationally representative survey. *American Journal of Preventive Medicine.* 2016;50(2):230–5.
229. Jackson JK, Shepherd TR, Kell RT. The influence of periodized resistance training on recreationally active males with chronic nonspecific low back pain. *Journal of Strength and Conditioning Research.* 2011;25(1):242–251.
230. Shamsi M, Sarrafzadeh J, Jamshidi A, Arjmand N, Arjmand F. Comparison of spinal stability following motor control and general exercises in nonspecific chronic low back pain patients. *Clinical Biomechanics.* 2017; 48:42–48.
231. Noormohammadpour P, Kordi M, Mansournia M, Akbari-Fakhrabadi M, Kordi R. Exercise Program in the Treatment of Nurses with Chronic Low Back Pain: A Single-Blinded Randomized Controlled Trial. *Asian Spine J.* 2018 Jun; 12(3): 490–502.
232. Wilson A, Hides JA, Blizzard L, Callisaya M, Cooper A, Srikanth VK, Winzenberg T. Measuring ultrasound images of abdominal and lumbar multifidus muscles in older adults: a reliability study. *Manual Therapy.* 2016; 23:114–119.
233. Wallwork TL, Hides JA, Stanton WR. Intratester and interrater reliability of assessment of lumbar multifidus muscle thickness using rehabilitative ultrasound imaging. *Journal of Orthopaedic & Sports Physical Therapy.* 2007;37(10):608–612.
234. Stokes M, Rankin G, Newham DJ. Ultrasound imaging of lumbar multifidus muscle: normal reference ranges for measurements and practical guidance on the technique. *Manual Therapy.* 2005;10(2):116–126.
235. Jarvik JG, Hollingworth W, Martin B, Emerson SS, Gray DT, Overman S, Robinson D, Staiger T, Wessbecher F, Sullivan SD, Kreuter W, Deyo RA. Rapid magnetic resonance imaging vs radiographs for patients with low back pain: a randomized controlled trial. *JAMA.* 2003;289(21):2810–2818.
236. Hodges PW, Tucker K. Moving differently in pain: a new theory to explain the adaptation to pain. *Pain.* 2011;152(Supplement): S90–S98.
237. Taylor JB, Goode AP, George SZ, Cook CE. Incidence and risk factors for first-time incident low back pain: a systematic review and meta-analysis. *Spine J.* 2014;14: 2299–231.
238. Fillingim RB. Individual differences in pain. *Pain.* 2017;158(suppl 1): S11–S18.
239. Rabey M, Beales D, Slater H, O'Sullivan P. Multidimensional pain profiles in four cases of chronic non-specific axial low back pain: an examination of the limitations of contemporary classification systems. *Man Ther.* 2015;20: 138–147.
240. Pires D, Cruz EB, Caeiro C. Aquatic exercise and pain neurophysiology education versus aquatic exercise alone for patients with chronic low back pain: a randomized controlled trial. *Clin Rehabil.* 2015; 29:538–547.
241. Louw A, Diener I, Butler DS, et al. The effect of neuroscience education on pain, disability, anxiety, and stress in chronic musculoskeletal pain. *Arch Phys Med Rehabil.* 2011; 92:2041–2056.
242. Peek AL, Miller C, Heneghan NR. Thoracic manual therapy in the management of non-specific shoulder pain: a systematic review. *J Man Manip Ther.* 2015;23(4):176–187.
243. Arias-Buría JL, Valero-Alcaide R, Cleland JA, Salom-Moreno J, Ortega-Santiago R, Atin-Arratibel MA, Fernández-de-las-Peñas C. Inclusion of trigger point dry needling in a multimodal physical therapy program for postoperative shoulder pain: a randomized clinical trial. *J Manip Physiol Ther.* 2015;38(3):179–187.

244. Liu L, Huang QM, Liu QG, Ye G, Bo CZ, Chen MJ, Li P. Effectiveness of dry needling for myofascial trigger points associated with neck and shoulder pain: a systematic review and meta-analysis. *Arch Phys Med Rehabil*. 2015;96(5):944-55.
245. Segura-Pérez M, Hernández-Criado MT, Calvo-Lobo C, Vega-Piris L, Fernández-Martín R, Rodríguez-Sanz D., A Multimodal Approach for Myofascial Pain Syndrome: A Prospective Study. *J Manipulative Physiol Ther*. 2017 Jul-Aug;40(6):397-403.
246. Marcos-Martín F, González-Ferrero L, Martín-Alcocer N, Paris-Aleman A, La Touche R. Multimodal physiotherapy treatment based on a biobehavioral approach for patients with chronic cervico-craniofacial pain: a prospective case series. *Physiother Theory Pract*. 2018 Sep;34(9):671-681.
247. Núñez-Cortés R, Cruz-Montecinos C, Vásquez-Rosel Á, Paredes-Molina O, Cuesta-Vargas. Dry Needling Combined With Physical Therapy in Patients With Chronic Postsurgical Pain Following Total Knee Arthroplasty: A Case Series. *J Orthop Sports Phys Ther*. 2017 Mar;47(3):209-216.
248. Moustafa IM, Diab AA, Hegazy F, Harrison DE. Does improvement towards a normal cervical sagittal configuration aid in the management of cervical myofascial pain syndrome: a 1-year randomized controlled trial. *BMC Musculoskelet Disord*. 2018 Nov 12;19(1):396
249. Pérez-Palomares S, Oliván-Blázquez B, Pérez-Palomares A, Gaspar-Calvo E, Pérez-Benito M, López-Lapeña E, de la Torre-Beldarraín ML, Magallón-Botaya R. Contribution of Dry Needling to Individualized Physical Therapy Treatment of Shoulder Pain: A Randomized Clinical Trial *J Orthop Sports Phys Ther*. 2017 Jan;47(1):11-20.
250. de Morton, NA. The PEDro scale is a valid measure of the methodological quality of clinical trials: a demographic study. *Aust J Physiother*. 2009;55 (2), 129e133.
251. Llamas-Ramos R, Pecos-Martín D, Gallego-Izquierdo T, Llamas-Ramos I, Plaza-Manzano G, Ortega-Santiago R, Cleland J, Fernández-de-Las-Peñas C. Comparison of the short-term outcomes between trigger point dry needling and trigger point manual therapy for the management of chronic mechanical neck pain: a randomized clinical trial. *J Orthop Sports Phys Ther*. 2014; 44: 852– 861.
252. Jaeger B. Myofascial trigger point pain. *Alpha Omegan*. 2013 Spring-Summer;106(1-2):14-22.
253. He C, Ma H. Effectiveness of trigger point dry needling for plantar heel pain: a meta-analysis of seven randomized controlled trials. *Journal of Pain Research*. 2017;10:1933–1942.
254. Martín-Pintado-Zugasti A, Rodríguez-Fernández AL, Fernández-Camero J. Postneedling soreness after deep dry needling of a latent myofascial trigger point in the upper trapezius muscle: Characteristics, sex differences and associated factors. *Journal of Back and Musculoskeletal Rehabilitation*. 2016;29(2):301-308.
255. Mason JS, Crowell M, Dolbeer J, Morris J, Terry A, Koppenhaver S, Goss DL. 2016. The effectiveness of dry needling and stretching vs. stretching alone on hamstring flexibility in patients with knee pain: a randomized controlled trial. *International Journal of Sports Physical Therapy*. 2016 Oct;11(5):672-683.
256. Fernández-Camero J, Gilaranz-de-Frutos L, León-Hernández JV, Pecos-Martín D, Alguacil-Diego I, Gallego-Izquierdo T, Martín-Pintado-Zugasti A. Effectiveness of different deep dry needling dosages in the treatment of patients with cervical myofascial pain: a pilot RCT. *American Journal of Physical Medicine and Rehabilitation*, 2017;96(10):726-733.
257. Calvo-Lobo C, Pacheco-Da-Costa S, Martínez-Martínez J, Rodríguez-Sanz D, Cuesta-Alvaro P, Lopez-Lopez D. Dry needling on the infraspinatus latent and active myofascial trigger points in older adults with nonspecific shoulder pain: A randomized clinical trial. *Journal of Geriatric Physical Therapy*. 2018;41(1):1-13.
258. Cerezo-Téllez E, Torres-Lacomba M, Fuentes-Gallardo I, Perez-Muñoz M, MayoralDel-Moral O, Lluch-Girbés E, Prieto-Valiente L, Falla D. Effectiveness of dry needling for chronic nonspecific neck pain: a randomized, single-blinded, clinical trial. *Pain*. 2016;157(9):1905-1917.
259. Barbero M, Falla D, Mafodda L, Cescon C, Gatti R. The location of peak upper trapezius muscle activity during submaximal contractions is not associated with the location of myofascial trigger points: new insights revealed by highdensity surface EMG. *Clinical Journal of Pain*. 2016;32(12):1044-1052.
260. Rathbone ATL, Grosman-Rimon L, Kumbhare DA. Interrater agreement of manual palpation for identification of myofascial trigger points: a systematic review and meta-analysis. *Clinical Journal of Pain*. 2017;33(8):715-729.
261. Poveda-Pagán EJ, Lozano-Quijada C, Segura-Heras JV, Peral-Berna M, Lumbreras B. Referred pain patterns of the infraspinatus muscle elicited by deep dry needling and manual palpation. *Journal of Alternative and Complementary Medicine*. 2017;23(11):890-896.
262. Tsai C-T, Hsieh L-F, Kuan T-S, Kao M-J, Chou L-W, Hong C-Z. Remote effects of dry needling on the irritability of the myofascial trigger point in the upper trapezius muscle. *Am J Phys Med Rehabil*. 2010;89:133–140.
263. Roy JS, MacDermid JC, Woodhouse LJ: Measuring shoulder function: a systematic review of four questionnaires. *Arthritis Rheum*. 2009;61:623-632.
264. Kamper SJ, Ostelo RW, Knol DL, Maher CG, de Vet HC, Hancock MJ: Global Perceived Effect scales provided reliable assessments of health transition in people with musculoskeletal disorders, but ratings are strongly influenced by current status. *J Clin Epidemiol*. 2010;63:760-766.e1.
265. Beaton DE, Katz JN, Fossel AH, Wright JG, Tarasuk V, Bombardier C: Measuring the whole or the parts? validity, reliability, and responsiveness of the Disabilities of the Arm, Shoulder and Hand outcome measure in different regions of the upper extremity. *J Hand Ther*. 2001;14:128-14.
266. Lucas KR, Rich PA, Polus BI. Muscle activation patterns in the scapular positioning muscles during loaded scapular plane elevation: the effects of Latent Myofascial Trigger Points. *Clin Biomech (Bristol, Avon)*. 2010;25:765–770.
267. Celik D, Yeldan I. The relationship between latent trigger point and muscle strength in healthy subjects: a double-blind study. *J Back Musculoskelet Rehabil*. 2011;24:251–256
268. Ge HY, Monterde S, Graven-Nielsen T, Arendt-Nielsen L. Latent myofascial trigger points are associated with an increased intramuscular electromyographic activity during synergistic muscle activation. *J Pain*. 2014;15:181–187.
269. Jinks C, Jordan K, Croft P. Measuring the population impact of knee pain and disability with the Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC). *Pain*. 2002;100:55-64.
270. Conboy VB, Morris RW, Kiss J, Carr AJ. An evaluation of the Constant-Murley shoulder assessment. *J Bone Joint Surg Br*. 1996;78:229-232.
271. Roy JS, MacDermid JC, Goel D, Faber KJ, Athwal GS, Drosdowech DS. What is a successful outcome following reverse total shoulder arthroplasty? *Open Orthop J*. 2010;4:157-163.
272. MacDermid JC, Walton DM, Avery S, Blanchard A, Etruw E, McAlpine C, Goldsmith CH. Measurement properties of the neck disability index: a systematic review. *J Orthop Sports Phys Ther*. 2009;39:400–417.
273. Juul T, Søgaard K, Davis AM, Roos EM. Psychometric properties of the neck outcome score, neck disability index, and short form–36 were evaluated in patients with neck pain. *J Clin Epidemiol*. 2016;79:31–40.
274. Kirkley A, Griffin S, Dainty K. Scoring systems for the functional assessment of the shoulder. *Arthroscopy*. 2003;19:1109-1120.
275. Shumway-Cook A, Woollacott MH. Motor control: theory and practical applications. Lippincott Williams & Wilkins. 2001.
276. Falla D, Jull G, Russell T, Vicenzino B, Hodges P. Effect of neck exercise on sitting posture in patients with chronic neck pain. *Phys Ther*. 2007;87:408–417.
277. Brink Y, Crous LC, Louw GA, Qimmer-Somers K, Schreve K. The association between postural alignment and psychosocial factors to upper quadrant pain in high school students: a prospective study. *Man Ther*. 2009;14:647–653.
278. Lombardi I, Jr., Magri ÁG., Fleury AM., Da Silva AC., Natour J. and Progressive resistance training in patients with shoulder impingement syndrome: a randomized controlled trial. *Arthritis Rheum*. 2008; 59: 615– 622.
279. Samarinos J., Marinko L. and Effectiveness of manual physical therapy for painful shoulder conditions: a systematic review. *J Man Manip Ther*. 2009; 17: 206– 215.
280. Solway S., Brooks D., Lacasse Y., Thomas S. and A qualitative systematic overview of the measurement properties of functional walk tests used in the cardiorespiratory domain. *Chest*. 2001; 119: 256– 270
281. Podsiadlo D, Richardson S. The timed "Up & Go": a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc*. 1991;39:142– 148.
282. Janssen WG, Bussmann HB, Stam HJ. Determinants of the sit-to-stand movement: a review. *Phys Ther*. 2002;82:866– 879.
283. Dobson F, Hinman RS, Roos EM, Abbott JH, Stratford P, Davis AM, Buchbinder R, Snyder-Mackler L, Henrotin Y, Thumboo J, Hansen P, Bennell KL. OARSI recommended performance-based tests to assess physical function in people diagnosed with hip or knee osteoarthritis. *Osteoarthritis Cartilage*. 2013; 21:1042– 1052.
284. Smith WN, Del Rossi G, Adams JB, Abdelrahman KZ, Asfour SA, Roos BA, Signorile JF. Simple equations to predict concentric lower-body muscle power in older adults using the 30-second chair-rise test: a pilot study. *Clin Interv Aging*. 2010;5:173– 180.
285. Puenteadura EJ, Flynn T. Combining manual therapy with pain neuroscience education in the treatment of chronic low back pain: A narrative review of the literature. *Physiother Theory Pract*. 2016 Jul;32(5):408-14.
286. Lluch Girbes E, Mees M, Baert I, Nijs, J. Balancing "hands-on" with "hands-off" physical therapy interventions for the treatment of central sensitization pain in osteoarthritis. *Man Ther*. 2015;20, 349-352.
287. Ludewig PM, Reynolds JF. The association of scapular kinematics and glenohumeral joint pathologies. *J Orthop Sports Phys Ther*. 2009; 39: 90– 104
288. Turp JC, Jokstad A, Motschall E, Schindler HJ, Windecker-Getaz I, Ettlin DA, Is there a superiority of multimodal as opposed to simple therapy in patients with temporomandibular disorders? A qualitative systematic review of the literature. *Clinical Oral Implants Research*. 2007;18 Suppl 3, 138-150
289. Kanas M, Faria RS, Salles LG, Sorpreso ICE, Martins DE, Cunha RAD, Wajchenberg M. Home-based exercise therapy for treating non-specific chronic low back pain. *Rev Assoc Med Bras (1992)*. 2018 Sep;64(9):824-831
290. Myers JB, Wassinger CA, Lephart SM. Sensorimotor contribution to shoulder stability: effect of injury and rehabilitation. *Man Ther*. 2006; 11: 197– 201.
291. Giannakopoulos NN, Rauer AK, Hellmann D, Hugger S, Schmitter M, Hugger A. Comparison of device-supported sensorimotor training and splint intervention for myofascial temporomandibular disorder pain patients. *J Oral Rehabil*. 2018 Sep;45(9):669-676.
292. Hakgüder A, Taştekin N, Birtane M, Uzunca K, Zateri C, Süt N. Comparison of the short-term efficacy of physical therapy in subacromial impingement syndrome patients with stage I and II magnetic resonance imaging findings. *Türk J Rheumatol*. 2011; 26: 127– 134.
293. Sahrmann S. Movement System Impairment Syndromes of the Extremities, Cervical and Thoracic Spines. St Louis, MO: Elsevier, Mosby; 2010.
294. Dinnes J, Loveman E, McIntyre L, Waugh N. The effectiveness of diagnostic tests for the assessment of shoulder pain due to soft tissue disorders: a systematic review. *Health Technol Assess*. 2003;7:iii, 1-166.