

fizjoterapia polska



POLISH JOURNAL OF PHYSIOTHERAPY

OFICJALNE PISMO POLSKIEGO TOWARZYSTWA FIZJOTERAPII

THE OFFICIAL JOURNAL OF THE POLISH SOCIETY OF PHYSIOTHERAPY

NR 3/2022 (22) DWUMIESIĘCZNIK ISSN 1642-

The use of deep friction massage techniques and Kinesiology Taping® (KT) in patients with faulty body posture

Zastosowanie masażu głębokiego i technik Kinesiology Tapingu® (KT) u pacjentów z wadliwą postawą ciała



Post-COVID-19 rehabilitation in the post-hospitalization period

Postępowanie rehabilitacyjne po przebytej chorobie COVID-19 w okresie poszpitalnym

ZAMÓW PRENUMERATE!

SUBSCRIBE!

www.fizjoterapiapolska.pl

www.djstudio.shop.pl

prenumerata@fizjoterapiapolska.pl



ULTRASONOGRAFIA W FIZJOTERAPII



Autoryzowani dystrybutorzy

Mar-Med

+48 22 853 14 11

info@mar-med.pl

Ado-Med

+48 32 770 68 29

adomed@adomed.pl


MAR-MED
OD 1995 ROKU

 **ADO-MED®**
APARATURA MEDYCZNA



Zawód Fizjoterapeuty dobrze chroniony

Poczuj się bezpiecznie



INTER Fizjoterapeuci

Dedykowany Pakiet Ubezpieczeń

Zaufaj rozwiązaniom sprawdzonym w branży medycznej.

Wykup dedykowany pakiet ubezpieczeń INTER Fizjoterapeuci, który zapewni Ci:

- ochronę finansową na wypadek roszczeń pacjentów
— **NOWE UBEZPIECZENIE OBOWIĄZKOWE OC**
- ubezpieczenie wynajmowanego sprzętu fizjoterapeutycznego
- profesjonalną pomoc radców prawnych i zwrot kosztów obsługi prawnej
- odszkodowanie w przypadku fizycznej agresji pacjenta
- ochronę finansową związaną z naruszeniem praw pacjenta
- odszkodowanie w przypadku nieszczęśliwego wypadku

Nasza oferta była konsultowana ze stowarzyszeniami zrzeszającymi fizjoterapeutów tak, aby najskuteczniej chronić i wspierać Ciebie oraz Twoich pacjentów.

► Skontaktuj się ze swoim agentem i skorzystaj z wyjątkowej oferty!

Towarzystwo Ubezpieczeń INTER Polska S.A.

Al. Jerozolimskie 142 B

02-305 Warszawa

www.interpolska.pl

inter
UBEZPIECZENIA



zabezpiecz się przed potencjalnymi **roszczeniami** **pacjentów**

program ubezpieczeń dla fizjoterapeutów
pod patronatem PTF

dla kogo?

Zarówno dla fizjoterapeutów prowadzących własną działalność w formie praktyki zawodowej, podmiotu leczniczego jak również tych, którzy wykonują zawód wyłącznie na podstawie umowy o pracę lub umowy zlecenie.

co obejmuje program ubezpieczeń?

- igłoterapie
- zabiegi manualne (mobilizacje i manipulacje)
- leczenie osteopatyczne
- naruszenie praw pacjenta i szkody w mieniu pacjentów

oraz szereg innych rozszerzeń ukierunkowanych na zawód fizjoterapeuty



kontakt w sprawie ubezpieczeń:

Piotr Gnat

+48 663 480 698

piotr.gnat@mentor.pl

[linkedin.com/in/piotrgnat](https://www.linkedin.com/in/piotrgnat)

ubezpiecz się **on-line** na **PTFubezpieczenia.pl**



THUASNE

SpryStep®

Dynamiczne ortezy
stawu skokowego i stopy

UŁATWIA
PORUSZANIE SIĘ
Z KAŻDYM KROKIEM



© Makenzie Hollar - © Studio Caterin - © Jennifer Williams - © M-1 STUDIOS

SpryStep® flex i SpryStep® mają na celu zapewnienie wsparcia pacjentom cierpiącym z powodu opadającej stopy pochodzenia neurologicznego, pourazowego lub mięśniowego. SpryStep® plus i SpryStep® max mają na celu zapewnienie podprarcia stopy oraz dolnej kończynie pacjenta z osłabieniem dolnych kończyn pochodzenia neurologicznego, urazowego lub mięśniowego. Produkty z grupy SpryStep® są wyrobami medycznymi, posiadającymi oznakowanie CE. Przed użyciem należy zapoznać się z instrukcją oraz zasięgnąć porady specjalisty. SpryStep®, SpryStep® flex, SpryStep® plus i SpryStep® max : Thuasne Deutschland GmbH, Germany

TANITA

Healthy Habits for Happiness



JAPANESE
TECHNOLOGY

WORLD NO. 1
FOR NEW HEALTH INSIGHTS

Analizatory Składu Ciała

Mierz, monitoruj, motywuj.

Daj swoim pacjentom informacje, których potrzebują do osiągnięcia sukcesu!

Analiza składu ciała wykonywana jest w około 30 sekund, a wyniki przedstawiane są na przejrzystym raporcie.



Produkty profesjonalne TANITA wykorzystywane są przez szpitale, ośrodki badawcze, centra diagnostyczne, placówki rehabilitacyjne, kluby sportowe, osoby pracujące ze sportowcami różnych dyscyplin na całym świecie.

Więcej na tanitapolska.pl

MICROGATE



OPTOGAIT

INNOWACYJNA DIAGNOSTYKA ZDOLNOŚCI MOTORYCZNYCH I ANALIZA CHODU

Systemy MICROGATE wspierają diagnozę, ocenę postępów oraz proces rehabilitacji.

Modelowanie programów rehabilitacyjnych i kontrola procesu rehabilitacji są ułatwione dzięki obiektywnej ocenie sposobu ruchu, wykrywaniu problematycznych obszarów, ocenie biomechanicznych braków oraz ocenie asymetrii.

Możliwe parametry pomiarowe:

- fazy chodu lub biegu
- długość kroku
- prędkość i przyspieszenie
- równowaga i symetria ruchu
- video Full HD

... i wiele innych, w zależności od przeprowadzonych testów.

W połączeniu z GYKO, możliwa jest ocena stabilności dynamicznej tułowia podczas chodu/biegu, analiza skoku, analiza stabilności posturalnej, analiza zakresu ruchomości stawów (ROM), ocena siły mięśniowej.

Więcej na microgatepolska.pl

EXXENTRIC

FLYWHEEL TRAINING

Trening siłowy i rehabilitacja z użyciem zmiennej bezwładności kół zamachowych.

Exxentric wykorzystuje moment bezwładności koła zamachowego zamiast zwykłej grawitacji. To daje możliwość wykonywania ćwiczeń standardowych oraz zaawansowanych metod treningu ekscentrycznego, koncentrycznego i izometrycznego.

Jako skuteczna metoda poprawy siły i stabilności, trening ekscentryczny ułatwi pacjentom osiągnięcie zamierzonych efektów – poprawy ogólnego stanu zdrowia, wyników sportowych, rehabilitacji, czy zapobieganiu urazom.

Szeroki wybór akcesoriów i dodatków do treningu z kołem zamachowym pomoże w stworzeniu idealnego rozwiązania dla Ciebie.

Więcej na treningekscentryczny.pl



POWRÓT DO SPRAWNOŚCI PO AMPUTACJI

Czym jest program po amputacji?

Po Amputacji to unikalne w skali kraju rozwiązanie dla osób, u których konieczna była operacja odjęcia kończyn. Celem programu jest kompleksowe wsparcie w procesie odzyskania sprawności po amputacji, niezależnie od jej przyczyny. Pomagamy pokonywać granice, osiągać kolejne cele, kreować bardziej przyjazną rzeczywistość.



W ramach programu oferujemy:

- Opieka Menadżera Rehabilitacji
- Dofinansowanie do zakupu protez
- Wsparcie psychologiczne
- Bezpłatne konsultacje protetyczne
- Rehabilitacja w ośrodkach na terenie kraju
- Pomoc Asystentów Wsparcia

**Skontaktuj się z nami i zapytaj
o bezpłatne egzemplarze
Poradnika dla
osób po
amputacji
do Twojej
placówki.**



**Masz pytanie odnośnie programu.
Napisz do nas lub skontaktuj się z nami
telefonicznie:**

+48 793 003 695

biuro@poamputacji.pl www.poamputacji.pl



POAMPUTACJI
Razem możemy więcej

SPRZEDAŻ I WYPOŻYCZALNIA ZMOTORYZOWANYCH SZYN CPM ARTROMOT®

Nowoczesna rehabilitacja **CPM** stawu kolanowego, biodrowego, łokciowego, barkowego, skokowego, nadgarstka oraz stawów palców dłoni i kciuka.



ARTROMOT-H



ARTROMOT-F



ARTROSTIM
FOCUS PLUS

ARTROMOT-K1 ARTROMOT-SP3 ARTROMOT-S3 ARTROMOT-E2



Najnowsze konstrukcje ARTROMOT zapewniają ruch bierny stawów w zgodzie z koncepcją **PNF** (Proprioceptive Neuromuscular Facilitation).

KALMED Iwona Renz
 ul. Wilczak 3
 61-623 Poznań
 www.kalmed.com.pl

tel. 61 828 06 86
 faks 61 828 06 87
 kom. 601 64 02 23, 601 647 877
 kalmed@kalmed.com.pl

Serwis i całodobowa
 pomoc techniczna:
 tel. 501 483 637
 service@kalmed.com.pl

ULTRASONOGRAFIA W FIZJOTERAPII



Autoryzowani dystrybutorzy

Mar-Med

+48 22 853 14 11

info@mar-med.pl

Ado-Med

+48 32 770 68 29

adomed@adomed.pl


MAR-MED
OD 1995 ROKU

 **ADO-MED**
APARATURA MEDYCZNA

NOWOŚĆ W OFERCIE

ASTAR.

PhysioGo.Lite SONO

**NIEWIELKIE URZĄDZENIE
EFEKTYWNA TERAPIA ULTRADŹWIEKOWA**

Zaawansowana technologia firmy Astar to gwarancja niezawodności i precyzji parametrów. Urządzenie, dzięki gotowym programom terapeutycznym, pomaga osiągać fizjoterapeucie możliwie najlepsze efekty działania fal ultradźwiękowych.

Głowica SnG to bezobrotowe akcesorium o dużej powierzchni czola (17,3 cm² lub 34,5 cm² w zależności od wybranego trybu działania). Znajduje zastosowanie w klasycznej terapii ultradźwiękami, fonoforezie, terapii LIPUS i zabiegach skojarzonych (w połączeniu z elektroterapią).



wsparcie merytoryczne
www.fizjotechnologia.com



ul. Świt 33
43-382 Bielsko-Biała

t +48 33 829 24 40
astarmed@astar.eu

**POLSKI
PRODUKT**  **WYBIERASZ
I WSPIERASZ**

www.astar.pl

Dr. Comifort®

Nowy wymiar wygody.

Obuwie profilaktyczno-zdrowotne
o atrakcyjnym wzornictwie



APROBATA
AMERYKAŃSKIEGO
MEDYCZNEGO
STOWARZYSZENIA
PODIATRYCZNEGO



WYRÓB
MEDYCZNY

**Stabilny, wzmocniony
i wyściełany zapętek**
Zapewnia silniejsze
wsparcie łuku
podłużnego stopy

**Miękki, wyściełany
kołnierz cholewki**
Minimalizuje podrażnienia

Wyściełany język
Zmniejsza tarcie
i ulepsza dopasowanie

Lekka konstrukcja
Zmniejsza codzienne
zmęczenie

**Antypoślizgowa,
wytrzymała podeszwa
o lekkiej konstrukcji**
Zwiększa przyczepność,
amortyzuje i odciąża stopy

**Wysoka jakość materiałów
- oddychające siatki i naturalne skóry**
Dostosowują się do stopy,
utrzymują je w suchości
i zapobiegają przegrzewaniu

**Zwiększona
szerokość i głębokość
w obrębie palców
i przodostopia**
Minimalizuje ucisk
i zapobiega urazom

Trzy
rozmiary
szerokości

Podwyższona
tęgłość

Zwiększona
przestrzeń
na palce

**Ochronna przestrzeń
na palce - brak szwów
w rejonie przodostopia**
Minimalizuje możliwość zranień

WSKAZANIA

- haluksy • wkładki specjalistyczne • palce młotkowate, szponiaste • cukrzyca (stopa cukrzycowa) • reumatoidalne zapalenie stawów
- bóle pięty i podeszwy stopy (zapalenie rozciągniętej podeszwy - ostroga piętowa) • płaskostopie (stopa poprzecznie płaska)
- bóle pleców • wysokie podbicie • praca stojąca • nerwiak Mortona • obrzęk limfatyczny • opatrunki • ortezy i bandaże • obrzęki
- modzele • protezy • odciski • urazy wpływające na ścięgna, mięśnie i kości (np. ścięgno Achillesa) • wrastające paznokcie



ul. Wilczak 3
61-623 Poznań
tel. 61 828 06 86
fax. 61 828 06 87
kom. 601 640 223, 601 647 877
e-mail: kalmed@kalmed.com.pl
www.kalmed.com.pl



www.butyladzrowia.pl

www.dr-comifort.pl


fizjoterapia polska



sklep internetowy:
www.djstudio.shop.pl

w sklepie dostępne między innymi:

- archiwalne numery Fizjoterapii Polskiej w wersji papierowej
- artykuły w wersji elektronicznej
- książki poświęcone fizjoterapii
- prenumerata Fizjoterapii Polskiej





Polisa**Med**

program
**ubezpieczeń
dla studentów**
kierunków medycznych



Drodzy Studenci

szukający artykułów do pracy naukowej.

**Przypominamy o dobrowolnym ubezpieczeniu
OC studentów kierunków medycznych!**

dlatego warto je mieć?

- ponieważ bywa wymagane w trakcie praktyk, staży czy wolontariatu
- niektóre Uczelnie wymagają je do udziału w zajęciach praktycznych
- działa na całym świecie, a dodatkowo otrzymasz certyfikat w języku angielskim w razie wyjazdu na ERASMUS-a
- wywołuje uśmiech na twarzy Pań z dziekanatów – sami sprawdziliśmy!



**posiadamy również w ofercie
ubezpieczenia dla masażystów
i techników masażystów.**



Polisa**Med**

**kontakt w sprawie
ubezpieczeń:**


+48 56 642 41 82

kontakt@polisa.med.pl

Ubezpiecz się **on-line** na

polisa.med.pl

NOWOŚCI



FUNKCYJNA BIELIZNA LECZNICZA

PRZECIWŻYLAKOWA

Przeciwwżylakowe wyroby pończosznice włoskich producentów, bardzo skuteczne i niezwykle eleganckie. Dostępne **w I, II oraz III klasie kompresji** w wielu modelach, w różnym stopniu przezroczystości (m. in. wyjątkowo przezroczyste w II kl. ucisku), w szerokiej gamie kolorystycznej, w różnych wersjach długości, z palcami zamkniętymi lub otwartymi

● podkolanówki ● pończochy ● legginsy ● rajstopy ● rękawy kompresyjne

ANTYCELLULITOWA, NA LIMFODEMIĘ I LIPODEMIĘ

Bielizna i odzież wykonana jest z mikrofibry. Unikalny splot nawet przy najmniejszym ruchu wywołuje **efekt masażu**. Działanie stymuluje cyrkulację podskórną i drenaż limfatyczny. Prowadzi to do poprawy jakości skóry

- z włókna emana®
- z kofeiną i wit. E
- z nanosrebrem

NA NIETRZYMANIE MOCZU

Wyroby medyczne **wielokrotnego użytku** z dyskretną stałą wszywką o właściwościach chłonnych. Polecane jako codzienna bielizna gwarantująca ochronę przed przemakaniem - 100% absorpcji cieczy, zapewniająca całkowitą suchość warstw: zewnętrznej i wewnętrznej

- do wielokrotnego prania (min. 100 prań)

ART & COLL
MEDICALE

artcoll.pl

e-sklep@artcoll.pl

tel. 22 720 35 96

+48 510 160 100

Effect of low level laser versus electrical stimulation on facial nerve regeneration in patients with Bell's palsy: a randomized controlled trial

Wpływ laseroterapii niskiego poziomu i stymulacji elektrycznej na regenerację nerwu twarzowego u pacjentów z porażeniem Bella: randomizowane badanie kontrolowane

**Ahmed Mohammed Fouad Shoman^{1(A,B,C,D,E,F,G)}, Ashraf Hassan Saleh^{2(B,C,D,E)},
Ahmed Mohamed Abdelhalim^{3(A,B,E,F)}, Ahmed Nazmy Kassab^{4(A,C,D,E,F)}**

¹Burn and Surgery Department, Faculty of Physical Therapy, Cairo University, Egypt

²Burn and Surgery Department, Faculty of Physical Therapy, Badr University, Cairo, Egypt

³Internal Medicine and Geriatrics Department, Faculty of Physical Therapy, Badr University, Cairo, Egypt

⁴Department of Medical Applications of Lasers, National Institute of Laser Enhanced Sciences, Cairo University, Egypt

Abstract

Purpose. This study was done to compare between the effect of low level laser and electrical stimulation on Bell's palsy. **Methods.** A randomized controlled trial was done on 45 patients of both genders with Bell's palsy; their ages ranged from 18–45 years. Patients were randomly divided into three equal groups. All groups (A, B & C) have received medical treatment. In addition, group (A) (Low level laser therapy), was treated with low level laser therapy, two sessions/ week for 6 weeks, while group (B) (ES group) was treated with electrical stimulation for the same time and group (C) (control) followed the traditional treatment, which consisted of massage and facial exercises. The primary outcome for the study was the degree of facial nerve regeneration assessed by facial nerve electromyographic sharp wave presence, while the secondary outcomes were facial motor function, assessed by House-Brackmann scale and the quality of life, measured by facial disability index. The outcomes were evaluated pre and after 6 weeks of the interventions. **Results.** There was a statistically significant difference ($P = 0.0001$) between the three groups regarding facial disability index score and House-Brackmann score in favor of laser group. **Conclusion.** Low level laser treatment is more effective than electrical stimulation and traditional treatment in achieving functional recovery in Bell's palsy patients.

Key words:

facial palsy, low level laser therapy, electrical stimulation therapy

Streszczenie

Cel. Niniejsze badanie przeprowadzono w celu porównania wpływu laseroterapii niskiego poziomu i stymulacji elektrycznej na porażenie Bella. **Metody.** Randomizowane kontrolowane badanie przeprowadzono na 45 pacjentach obu płci z porażeniem Bella; ich wiek wahał się od 18 do 45 lat. Pacjenci zostali losowo podzieleni na trzy równe grupy. Wszystkie grupy (A, B i C) zostały poddane leczeniu. Dodatkowo grupa (A) (laseroterapia niskiego poziomu) była poddawana laseroterapii niskiego poziomu, dwie sesje/tydzień przez 6 tygodni, natomiast grupa (B) (grupa ES) była poddawana stymulacji elektrycznej przez ten sam czas, a grupa (C) (kontrolna) była poddawana leczeniu tradycyjnemu, które obejmowało masaż i ćwiczenia twarzy. Najważniejszym skutkiem przeprowadzanego badania był stopień regeneracji nerwu twarzowego oceniany na podstawie obecności ostrych fal nerwu twarzowego widocznych w badaniu elektromiograficznym, natomiast drugorzędnymi skutkami były funkcje motoryczne twarzy oceniane w skali House-Brackmanna oraz jakość życia mierzona wskaźnikiem niepełnosprawności twarzy. Wyniki zostały ocenione przed i po 6 tygodniach leczenia. **Wyniki.** Wystąpiła statystycznie istotna różnica ($P=0,0001$) między trzema grupami w zakresie wskaźnika niepełnosprawności twarzy i wyniku House-Brackmann na korzyść grupy poddawanej laseroterapii. **Wniosek.** Laseroterapia niskiego poziomu jest skuteczniejsza niż stymulacja elektryczna i leczenie tradycyjne w osiąganiu powrotu do zdrowia u pacjentów z porażeniem Bella.

Słowa kluczowe

porażenie nerwu twarzowego, laseroterapia niskiego poziomu, elektrostymulacja

Introduction

Bell's palsy, also known as idiopathic facial palsy, is regarded as the most common form of one-sided facial paralysis. It is an acute face paralysis that can occur because of facial nerve inflammation close to the stylomastoid foramen [1]. The paralysis of facial nerve is related to many various functional and cosmetic issues that could greatly impact the social and psychological state of the patients [2]. Bell's palsy is a lower motor neuron condition that is thought to cause around 60–75% of acute one-sided facial paralysis cases [1]. Bell's palsy can result into not only a disturbing facial deformity, but also an impaired capability to communicate using facial articulation and expression [3]. Physical therapy modalities that are mostly used for Bell's palsy include electrotherapy, massage, facial exercise, bio-feedback, and electrical stimulation [1, 4].

Electrical stimulation, as a therapeutic modality, has been widely accepted and commonly used for its ability to improve the performance of healthy and the diseased muscles skeletal muscles in both research studies and in clinical settings [5, 6].

Laser stands for light amplification of the stimulated emission of radiation, a form of phototherapy that includes applying monochromatic light upon biological tissues to provoke a biologically modulated response within those tissues [7].

Using laser light in therapeutic applications with low intensity is referred to as low level laser therapy (LLLT). LLLT augments cellular metabolism leading to accelerated energy production along with increased protein and collagen synthesis. Additionally, laser helps form new blood vessels, leads to vascular relaxation and enhances microcirculation of the skin [8].

Many studies have approved the efficacy of physical therapy modalities on patients with Bell's palsy, but in our knowledge, the comparison between the effect of electrical stimulation and low level laser on Bell's palsy was not studied. So, the aim of the study was to compare between the effect of low level laser and electrical stimulation on Bell's palsy.

Materials and methods

Design

A randomized control trial was conducted to compare between the effect of low level laser and electrical stimulation on Bell's palsy. Data were collected pre and post treatment from June 2018 to March 2019. The research was approved by the institutional review board at National Institute of LASER Enhanced Sciences, Cairo University [Cu-NILES/15/21], following the Declaration of Helsinki Guidelines on conducting the human research. Each patient was educated about the study assessment and treatment procedures and informed consent was obtained from each one of them.

Participants

Forty-five patients of both genders with Bell's palsy were selected from the outpatient clinic of faculty of physical therapy & the national institute of laser enhanced sciences, with their ages ranging from 18–45 years. They were randomly assigned into three equal groups (A, B & C).

Exclusion criteria included Pregnancy or breastfeeding, patients who received medical treatment or acupuncture therapy in other hospitals before this study, diabetes course ≥ 5 years,

hypertension (3rd grade) / hyperlipidemia present for 3 or more years. Also, a patient with a history of malignant tumor, sexually transmitted disease, renal or hepatic disease, gastric or duodenal ulcer, or glaucoma was excluded. Moreover, acute otitis, chronic otitis on the affected side, tuberculosis, immunodeficiency syndromes, recent head injury, psychiatric disease, infectious disease, or any other condition that puts the person at risk of being influenced by the study treatment or that may affect the completion of the study (e.g., participation in other clinical trials within 3 months) were also exclusion criteria.

Randomization

The recruited patients were randomly assigned, after signing consent form, into three equal groups by using computerized randomization. Serially numbered opaque envelopes were used and opened by an investigator who has no direct contact with the participants. Following randomization, there was no dropping out of subjects from the study, figure 1.

Interventions

Patients with Bell's palsy were assigned into three equal groups. Group (A) involved 15 patients, treated by LLLT in addition to medical treatment, while group (B) involved 15 patients, treated by electrical stimulation in addition to the same to medical treatment and group (C) involved 15 patients who received the traditional facial palsy treatment that consisted of facial massage, facial muscles' exercises and the same medical treatment. All patients received the interventions for 6 weeks.

Medical treatment in all groups involved administration of a daily dose of 60 mg of corticosteroids (prednisolone) for 10 days, with a tapered course, in addition to 400 mg of antiviral (acyclovir), five times a day for a week [9]. Also, topical ocular lubricant was applied to prevent corneal drying.

Group (A) (Low level laser therapy)

In addition to the medical treatment, group (A) received LLLT using Galium AL. Arsenide diode laser (chattanooga intellect mobile laser, USA), with a wavelength of 850 nm, and power density of 1 W/cm². The treatment was given twice a week for 6 weeks for a total of 12 sessions. Each session involved applying laser on 8 points (1 min/ point). Using direct skin contact, facial nerve upper, middle, and lower branches as well as the nerve trunk, orbicularis oris muscle, muscles of the nose, levators of the upper lip and depressors of the lower lip were irradiated [10].

Group (B) (electrical stimulation group)

In addition to the traditional treatment, group (B) received electrical stimulation using electrical stimulator unit (faradic current- Glopus multicurrent, Italy), with a pulse duration ranging from 0.1ms to 1 ms and a frequency of 50 HZ. Intensity level was increased till having a visible muscle contraction. Electrical stimulation was applied by placing one electrode in front of the ear where the nerve trunk is present, while the other electrode was placed on the stimulated muscle. Stimulation was applied on eleven facial muscles, each for 10 min, achieving 30 minimal muscle contractions divided to 3 sets, 5 times/ week for 6 weeks [11].




Figure 1. Flow chart of participants

Group (C) (control)

This group was treated with medical treatment added to massage and facial muscles' exercises, as the traditional facial palsy treatment. Patients in group (C) were taught to perform massage and facial exercises correctly by a physiotherapist and they were instructed to do the massage and exercises three times a day throughout treatment period for 6 weeks. Patients were also recommended to blow a balloon and to chew a gum on the paralyzed side. The given face and neck massage were done to relax the facial musculature and promote circulation. Also, they performed active assistive resistive exercises in front of a mirror and underwent proprioceptive neuromuscular facilitation [8].

Outcome measures

Electromyography (EMG) for facial nerve

Using EMG in monitoring facial nerve regeneration is helpful in predicting forthcoming recovery, as identifying muscle re-innervation using EMG can precede obvious clinical movements by 2–3 months [12]. Before application, the face was

washed with water and soap to remove sweat or any other interference. According to Schumann et., [13] the electrode placement for specific muscle movement recording followed a typical pattern, where the stimulating electrode was put on facial nerve trunk on the stylomastoid foramen, and the recording electrodes were put on the facial muscles. Comparison between the unaffected and affected side was done to calculate the percentage of degeneration. This test was done pre and post treatment program for all groups.

Facial disability index (FDI)

The FDI is a widely used tool that aims to evaluate specific features related to quality of life, which can be affected due to the alterations in facial impression. It is a brief, self-reported questionnaire that involves ten questions assigned to physical impairment and psychosocial subscales, where having a higher score indicated a better quality of life. It has an internal consistency (Cronbach's alpha coefficient) of 0.83 for the whole scale, making it an effective and reliable assessment tool [14].

House-Brackmann scale (HB)

The House-Brackmann scale is a universally-validated simple grading method that can be utilized to assess the function of facial nerve [15], monitor its regeneration, and evaluates its recovery post treatment. The scale categorizes the paresis degree into six levels, where the first level represents no paresis and the sixth represents total paresis. The net score for the scale is attained by including the score for each of the facial nerve branches, thus the total scale values range from 6 up to 36 [14].

Statistical analysis

The study statistical analysis was carried out using the statistical Package program SPSS, version 25 for Windows (SPSS, Inc., Chicago, IL). Patients' descriptive statistics, involving demographic data and other study variables (EMG findings, FDI, and House- Brackmann scale) were represented by the mean and standard deviation or the median and range whenever applicable. To compare the study variables, a multivariate

analysis of variance (MANOVA) 3 x 2 – mixed design was utilized, where the first independent variable was represented by the three tested groups and the second independent variable was before and after treatment assessment periods. Paired t-test was utilized to compare results before and after treatment within each group of the three groups for facial EMG positive sharp wave presence, FDI, and House- Brackmann scale, while unpaired t-test was utilized to compare the results between LLLT group, electrical stimulation group and the control group for the same variables. Significant results were indicated when the level of probability was at 0.05 ($P \leq 0.05$).

Results

Forty-five patients, diagnosed as having facial palsy, participated in the study and they were randomly assigned into three equal groups of 15 patients each. At baseline, there were no significant differences between the three groups regarding age, gender, or side of affection ($p > 0.05$), (Table 1).

Table 1. Descriptive statistics and one-way analysis of variance (ANOVA) for age, gender, and side of affection of the three tested groups

Age (years)	Group A (N = 20)	Group B (N = 20)	Group C (N = 20)	F-value	P-value	Significance
Mean $\pm$ SD	34.66 $\pm$ 8.44	32.8 $\pm$ 8.09	32.66 $\pm$ 8.86	0.261	0.772	N.S
Gender distribution	Group A (N = 20)	Group B (N = 20)	Group C (N = 20)	χ^2 –value	P-value	Significance
Male, N (%)	10 (66.7%)	10 (66.7%)	8 (53.3%)	0.756	0.685	N.S
Female, N (%)	5 (33.3%)	5 (33.3%)	7 (46.7%)			
Side distribution	Group A (N = 20)	Group B (N = 20)	Group C (N = 20)	χ^2 –value	P-value	Significance
Left side, N (%)	8 (53.3%)	6 (40%)	6 (40%)	0.72	0.698	N.S
Right side, N (%)	7 (46.7%)	9 (60%)	9 (60%)			

N, number of cases N.S, P value > 0.05

Regarding the overall effect, statistical analysis using MANOVA 3x2 mixed design showed significant effects of the tested group (the first independent variable) on the all tested dependent variables, which are FDI ($F = 12.237$, $P = 0.0001^*$).

Moreover, there were significant effects of the measuring

periods (the second independent variable) on the tested dependent variables ($F = 444.894$, $P = 0.0001^*$). Additionally, the interaction between the two independent variable was significant, showing that the effect of the tested groups on the dependent variables was influenced by the measuring periods ($F = 61.57$, $P = 0.0001^*$), (Table 2).

Table 2. The 3x2 mixed design MANOVA for dependent variable at different measuring periods between all groups

Source of variation	F- value	P-value
Groups	12.237	0.0001*
Measuring periods	444.894	0.0001*
Interaction effect	61.57	0.0001*

P-value: probability value * Significant at P-value < 0.05

For the House-Brackmann scale, within group's comparison of "pre treatment" versus "post treatment" median values showed significant difference in the three groups (A, B & C). Likewise, post hoc tests indicated a significant increase in House-Brackmann scale scoring at post treatment in comparison to pre-treatment in group (A) ($Z = -3.482$, $P = 0.0001^*$), group (B) ($Z = -3.482$, $P = 0.0001^*$) and group (C) ($Z = -3.276$, $P = 0.0001^*$), (Table 3).

Considering the effect of the tested group on House-Brackmann

scale, the Kruskal-Wallis H test showed a significant difference in the post-treatment median values of House-Brackmann scale among three groups ($P = 0.0001^*$). Using Mann-Whitney U test, results indicated significant differences between the groups (A & B) with $P = 0.0001^*$, the groups (A & C), with $P = 0.0001^*$ and the groups (B & C), with $P = 0.023^*$. This significant increase was in favor of group (A), compared to the groups (B & C) and in favor of group (B), compared to the group (C), (Table 4).

Table 3. Descriptive statistics and post hoc tests of the House -Brackmann scale among different groups

House-Brackmann scale	Group A, Median (IQR)	Group B, Median (IQR)	Group C, Median (IQR)
Pre treatment	2 (1)	2 (1)	2 (1)
Post treatment	6 (2)	4 (2)	3 (1)
Wilcoxon Signed Rank test	Group A, Median (IQR)	Group B, Median (IQR)	Group C, Median (IQR)
Z-value	-3.482	-3.482	-3.276
p-value	0.0001*	0.0001*	0.0001*

*Significant at alpha level < 0.05 , IQR: Interquartile Range

Table 4. Kruskal-Wallis H and Mann-Whitney U post hoc tests of the House -Brackmann scale among different groups

Kruskal-Wallis H test		χ 2-value		P-value		
Pre treatment		0.183		0.913		
Post treatment		26.318		0.0001*		
Mann-Whitney U test	Group (A) Vs. (B)		Group (A) Vs. (C)		Group (B) Vs. (C)	
	U-value	P-value	U-value	P-value	U-value	P-value
Pre treatment	103	0.713	106.5	0.806	109.5	0.902
Post treatment	23	0.0001*	7	0.0001*	58.5	0.023*

*Significant at alpha level < 0.05

For the FDI, within group's comparison of the "pre-treatment" versus "post treatment" mean values indicated significant increase in the three groups (A, B & C) ($P = 0.0001^*$). On comparing the three groups, post hoc tests revealed a significant difference among the groups (A & B)

and (group & C) with $P = 0.0001^*$ for both. This significant increase was in favor of group (A), compared to the group (B & C). While, there was no significant difference between (group B & C), with $P = 0.078$, there was a clinical difference and more improvement in favor of group B, (Table 5).

Table 5. Descriptive statistics and 3×2 mixed design MANOVA for Facial disability index at different measuring periods among different groups

FDI	Group A, (Mean ± SD)	Group B, (Mean ± SD)	Group C, (Mean ± SD)
Pre- treatment	24 ± 8.28	22.66 ± 7.03	23.33 ± 8.16
Post treatment	60 ± 11.33	40.66 ± 8.83	32.66 ± 7.98
MD	-36	-18	-9.33
% of change	150%	79.43%	39.99%

Multiple pairwise comparisons between pre and post treatment values for FDI at different groups			
Pre Vs. post	Group A	Group B	Group C
P-value	0.0001*	0.0001*	0.0001*
Multiple pairwise comparison tests (Post hoc tests) for the FDI among different groups at different measuring periods			
	Group (A) Vs. group (B)	Group (A) Vs. group (C)	Group (B) Vs. group (C)
Pre	0.99	0.98	0.97
Post	0.0001*	0.0001*	0.078

MD, mean difference *Significant at alpha level < 0.05

Regarding the presence of positive sharp wave in EMG, within group's comparison, using McNemar's test, showed a statistically significant increase in the reported percentage of patients having EMG positive sharp wave post treatment in the group (A), ($P = 0.016^*$), while there was no statistically significant increase on the percentage of those patients in the groups (B) & (C) ($P = 0.125$ and $P = 0.99$) respectively, (Table 6).

On comparing the three groups pre-treatment, Chi square tests (post hoc tests) revealed that there was no significant difference ($\chi^2 = 0.556$, $P = 0.757$) in the presence of EMG positive sharp wave between any of the three groups (A, B or C). Also, post hoc tests showed that there was no significant difference in the presence of EMG positive sharp wave among three groups post treatment ($\chi^2 = 4.2$, $P = 0.122$), (Table 6).

Table 6. Chi-squares tests and frequency distribution of the positive sharp wave presence in EMG at different measuring periods at different groups

Frequency distribution						
Group A		Group B		Group C		
Absent	Present	Absent	Present	Absent	Present	
Pre treatment	9 (60%)	6 (40%)	10 (66.7%)	5 (33.3%)	8 (53.3%)	7 (46.7%)
Post treatment	2 (13.3%)	13 (86.7%)	6 (40%)	9 (60%)	7 (46.7%)	8 (53.3%)
McNemar's tests						
Group A		Group B		Group C		
P-value	0.016*		0.125		0.99	
Chi square tests (among groups)						
χ ² -value			P-value			
Pre treatment			0.556			0.757
Post treatment			4.2			0.122
Chi square tests as Multiple pairwise comparison tests (post hoc tests) for the positive sharp wave presence in EMG among different groups at different measuring periods						
Group (A) Vs. group (B)		Group (A) Vs. group (C)		Group (B) Vs. group (C)		
χ ² -value	p-value	χ ² -value	p-value	χ ² -value	P-value	
Pre treatment	0.144	0.705	0.136	0.713	0.556	0.456
Post treatment	2.727	0.099	3.968	0.051	0.136	0.713

*Significant at alpha level < 0.05

Discussion

Bell's palsy is identified as the most frequent cause of one-sided peripheral facial nerve palsy. It commonly occurs at the ages between 15 to 45 years, with a relative high annual incident. Symptoms, involving weakness of facial muscles, retro-auricular pain pattern, abnormal noise tolerance level, and unilateral disturbed taste, are normally sudden at onset, ranging from mild to severe degree and they often peak within a day to 3 days. Symptoms include [16].

This Comparative study investigated effect of LLLT and electrical stimulation on 45 patients, divided into three equal groups of both genders, with Bell's palsy and their ages ranged from 18–45 years. In the current study there was no significant difference between the groups in the age, gender distribution or side of affection.

In agreement with that result, Ordahan and Karahan [17] showed that a total of 46 patients with unilateral facial paralysis had no statistically significant difference in mean age between the groups. Similarly, a study by Abdrabou et al. [18] showed insignificant differences between study groups in the duration of paresis.

The current study findings also revealed that there was no significant difference between the three groups regarding the presence of positive sharp wave in the EMG, while within groups, there was a significant difference in only group (A) that received LLLT with medication after 6 weeks of treatment.

Results from a study by Low et al. [19] agreed with these results as it showed that LLLT has positively affected normal peripheral nerve parameters in form of both nerve excitability and action potential. Also, Abdrabou et al. [18] indicated a significant improvement in latency, amplitude, and axonal loss in patients with Bell's palsy when treated by LLLT group than those treated by electric stimulation. While, on the other hand, Quinn et al. [20] disagree as they stated no hard evidence on whether the LLLT was effective in managing twenty one patients having Bell's palsy.

The present study also found a significant improvement in all groups post treatment regarding both House-Brackmann scale, and FDI, with that improvement was in favor of LLLT group. These findings were congruent to Khullar et al., [21] who stated that LLLT had improved nerve functional recovery, possibly due to direct augmentation of the axonal regeneration process. Additionally, Ordahan and Karahan [17] stated a greater improvement in FDI scores in the group receiving LLLT at 3 and 6 weeks post treatment, compared to the exercise group. Likewise, Alayat et al. [10] used FDI to evaluate the effect of LLLT on Bell's palsy outcomes, and observed functional improvement in patients who received LLLT. Moreover, Fontana and Bagnato [22] reported

a complete enhancement in a child aged 3 years having Bell's palsy following LLLT for 3 weeks. At the same time, a study of 228 children Bell's palsy, done by Prescott [23], discovered that a high-dosage steroid treatment did not significantly affect recovery times. In addition, Ladalardo et al. [24] reported that Bell's palsy patients, who were treated by gallium arsenide laser diodes, demonstrated a significant functional improvement.

Many studies [have explained the reason behind the improvement seen in patients with Bell's palsy undergoing LLLT. Generally, LLLT was found to stimulate the photoreceptors in the mitochondrial and cell membrane, resulting into enhanced cellular functions, and cell proliferation rate, as it increases DNA, and RNA synthesis as well as collagen production [25, 26]. Also, Kneebone [27] found that applying laser could enhance neuronal regeneration through both local and systematic actions. Another mechanism is that LLLT reduces neuronal retrograde degeneration in the neurons connected to the spinal cord and increases the damaged peripheral nerves healing [28].

Moreover, Chow et al. [29] determined that LLLT has a direct influence on nerve structure, leading to enhanced healing rate of communication blocks. Additional studies also researchers showed that LLLT dilates arterial and capillary blood vessels, thus it helps increase microcirculation, activate angiogenesis, and stimulate nerve regeneration [30].

Although few studies have discussed the efficacy of LLLT in the treatment of Bell's palsy cases, Bernal [31] revealed that it is a painless adjunctive treatment method, with no adverse effects for patients having facial paralysis, especially when they are contraindicated to the steroid treatments, as with diabetes and hypertension.

The current study has many strength points. First, it was randomized and involving patients of both males and females. Also, it used objective methods to assess outcomes. The study has a point of limitation, though. Further studies are needed to be conducted on patients of one gender who have Bell's palsy and to include a larger sample for more conclusive results.

Conclusion

Low level laser treatment is more effective than electrical stimulation and traditional treatment in managing Bell's palsy patients, and achieves better functional recovery and more improved quality of life.

Adres do korespondencji / Corresponding author

Ahmed Mohammed Fouad Shoman

E-mail:shomanpt2010@yahoo.com

Acknowledgement

The authors would like to thank all individuals who participated in this study.

Piśmiennictwo/ References

1. Teixeira LJ, Valbuza JS, Prado GF. Physical therapy for Bell's palsy (idiopathic facial paralysis). *Cochrane Database Syst Rev*.2011; 7(12):CD006283.
2. Pereira LM, Obara K, Dias JM, et al. Facial exercise therapy for facial palsy: systematic review and meta-analysis. *Clin Rehabil*. 2011;25(7):649-58.
3. Ho AL, Scott AM, Klassen AF, et al. Measuring quality of life and patient satisfaction in facial paralysis patients: a systematic review of patient-reported outcome measures. *Plast Reconstr Surg*. 2012;130(1):91-99.
4. Kahn JB, Gliklich RE, Boyev KP, et al. Validation of a patient-graded instrument for facial nerve paralysis: the FaCE scale. *Laryngoscope*. 2001;111(3):387-98.
5. Manikandan N. Effect of facial neuromuscular re-education on facial symmetry in patients with Bell's palsy: a randomized controlled trial. *Clin Rehab*.2007;21:338–343.
6. Kjartansson AW, Lundeberg EE. Effects of electrical nerve stimulation (ENS) in CPPS" *Scand J plastReconstr Hand Surg*.2006;16:119-25.
7. Kitchen S. Scientific background. *Electrotherapy: evidence-based practice*. 11th ed., Churchill Livingstone, 2002, 3-30.
8. Karu IT. Low power laser therapy. *Biomedical Photonics handbook*, CRC press; florida, USA, 2003, 1233-1258.
9. Tiemstra, J D, Khatkhate N. Bell's Palsy: Diagnosis and Management. *Am Fam Physician*. 2007;76(7):997-1002.
10. Alayat MS, Elsodany AM, El Fiky AA. Efficacy of high and low level laser therapy in the treatment of Bell's palsy: a randomized double blind placebo-controlled trial. *Lasers Med Sci*. 2014;29(1):335-42.
11. Tuncay F, Borman P, Taşer B, et al. Role of electrical stimulation added to conventional therapy in patients with idiopathic facial (Bell) palsy. *Am J Phys Med Rehabil*. 2015;94(3):222-8.
12. Guntinas-Lichius O, Volk GF, Olsen KD, et al. Facial nerve electrodiagnostics for patients with facial palsy: a clinical practice guideline. *Eur Arch Otorhinolaryngol*. 2020;277(7):1855-1874.
13. Schumann NP, Bongers K, Guntinas-Lichius O, Scholle HC. Facial muscle activation patterns in healthy male humans: a multi-channel surface EMG study. *J Neurosci Methods*. 2010;187(1):120-8.
14. Gonzalez-Cardero E, Infante-Cossio P, Cayuela A, et al. Facial disability index (FDI): adaptation to Spanish, reliability and validity. *Med Oral Patol Oral Cir Bucal*. 2012;17(6):e1006-e1012.
15. Evans RA, Harries ML, Baguley DM, Moffat DA. Reliability of the House and Brackmann grading system for facial palsy. *J Laryngol Otol*. 1989;103(11):1045-6.
16. Ton G, Lee LW, Ng HP, et al. Efficacy of laser acupuncture for patients with chronic Bell's palsy: A study protocol for a randomized, double-blind, sham-controlled pilot trial. *Medicine (Baltimore)*. 2019;98(15):e15120.
17. Ordahan B, Karahan A. Role of low-level laser therapy added to facial expression exercises in patients with idiopathic facial (Bell's) palsy. *Lasers Med Sci*.2017; 32(4): 931-936.
18. Abdrabou A, Mustafa Y, Amer K, Eldeep A. The effect of laser acupuncture vs electrical stimulation on patients with idiopathic Bells palsy. *IJRMR*. 2017; 4(10): 2942-2945.
19. Low J, Reed A. *Electrotherapy explained: principles and practice*, 2nd Ed, Butterworth Heinemann; London,1994, 117-133.
20. Quinn R, Cramp F. The efficacy of electrotherapy for Bell's Palsy, a systematic review". *Physical Therapy Reviews*.2003; 8(3):151-164
21. Khullar SM, Brodin P, Messelt EB, Haanaes HR. The effects of low level laser treatment on recovery of nerve conduction and motor function after compression injury in the rat sciatic nerve. *Eur J Oral Sci*. 1995;103(5):299-305.
22. Fontana CR and Bagnato VS. Low-level laser therapy in pediatric Bell's palsy: case report in a three-year-old child. *J Altern Complement Med*.2013;19:376–382.
23. Prescott CA. Idiopathic facial nerve palsy in children and the effect of treatment with steroid. *Int J Pediatr Otorhinolaryngol*.1987;13:257–264.
24. Ladalarido TC, Brugnara A Jr, Takamoto M, et al. Functional and electrophysiological evaluation of the effects of laser therapy in the treatment of peripheral facial paralysis. *Lasers in Dentistry VII*. 2001;4249:134–138
25. Kujawa J, Zavodnik L, Zavodnik I, et al. Effect of low-intensity (3.75-25 J/cm2) near-infrared (810 nm) laser radiation on red blood cell ATPase activities and membrane structure. *J Clin Laser Med Surg*. 2004;22(2):111-7.
26. AlGhamdi KM, Kumar A, Moussa NA. Low-level laser therapy: a useful technique for enhancing the proliferation of various cultured cells. *Lasers Med Sci*.2012; 27:237–249
27. Kneebone WJ. Enhancement of nerve regeneration by therapeutic laser. *Pract Pain Manag*.2010; 10:70–72
28. Barbosa RI, Marcolino AM, de Jesus Guirro RR, et al. Comparative effects of wavelengths of low-power laser in regeneration of sciatic nerve in rats following crushing lesion. *Lasers Med Sci*. 2010;25(3):423-30.
29. Chow R, Armati P, Laakso EL, et al. Inhibitory effects of laser irradiation on peripheral mammalian nerves and relevance to analgesic effects: a systematic review. *Photomed Laser Surg*. 2011;29(6):365-81.
30. Tam G. Low power laser therapy and analgesic action. *J Clin Laser Med Surg*.1999; 17:29–33.
31. Bernal G. Helium neon and diode laser therapy is an effective adjunctive therapy for facial paralysis. *Laser Ther*.1993;5:79–87.