

fizjoterapia polska

POLISH JOURNAL OF PHYSIOTHERAPY

OFICJALNE PISMO POLSKIEGO TOWARZYSTWA FIZJOTERAPII

THE OFFICIAL JOURNAL OF THE POLISH SOCIETY OF PHYSIOTHERAPY

NR 1/2023 (23) KWARTALNIK ISSN 1642-0136

Restytucja wielkości cech postawy w płaszczyźnie czołowej po obciążeniu masą transportowanych przyborów szkolnych w trybie ciagu lewą i prawą ręką uczniów obojga płci w wieku 7 lat

Restitution of the size of postural features in the frontal plane after loading with the weight of school items carried with the right and left hand in 7-year-old pupils of both sexes

Ocena postawy ciała gimnazjalistów w kontekście ich aktywności fizycznej

Assessment of body posture of junior high school students in the context of their physical activity

ZAMÓW PRENUMERATĘ!

SUBSCRIBE!


www.fizjoterapiapolska.pl

www.djstudio.shop.pl

prenumerata@fizjoterapiapolska.pl



ULTRASONOGRAFIA W FIZJOTERAPII



Autoryzowani dystrybutorzy

Mar-Med

+48 22 853 14 11

info@mar-med.pl

Ado-Med

+48 32 770 68 29

adomed@adomed.pl



MAR-MED

OD 1995 ROKU



ADO-MED

APARATURA MEDYCZNA



Zawód Fizjoterapeuty dobrze chroniony

Poczuj się bezpiecznie



INTER Fizjoterapeuci

Dedykowany Pakiet Ubezpieczeń

Zaufaj rozwiązaniom sprawdzonym w branży medycznej.

Wykup dedykowany pakiet ubezpieczeń INTER Fizjoterapeuci, który zapewni Ci:

- ochronę finansową na wypadek roszczeń pacjentów
— **NOWE UBEZPIECZENIE OBOWIĄZKOWE OC**
- ubezpieczenie wynajmowanego sprzętu fizjoterapeutycznego
- profesjonalną pomoc radców prawnych i zwrot kosztów obsługi prawnej
- odszkodowanie w przypadku fizycznej agresji pacjenta
- ochronę finansową związaną z naruszeniem praw pacjenta
- odszkodowanie w przypadku nieszczęśliwego wypadku

Nasza oferta była konsultowana ze stowarzyszeniami zrzeszającymi fizjoterapeutów tak, aby najskuteczniej chronić i wspierać Ciebie oraz Twoich pacjentów.

► Skontaktuj się ze swoim agentem i skorzystaj z wyjątkowej oferty!

Towarzystwo Ubezpieczeń INTER Polska S.A.

Al. Jerozolimskie 142 B

02-305 Warszawa

www.interpolska.pl

inter
UBEZPIECZENIA

fizjoterapia polska



sklep internetowy:
www.djstudio.shop.pl

w sklepie dostępne między innymi:

- *archiwalne numery Fizjoterapii Polskiej w wersji papierowej*
- *artykuły w wersji elektronicznej*
- *książki poświęcone fizjoterapii*
- *prenumerata Fizjoterapii Polskiej*

PATRONAT MERYTORYCZNY
Komitet Rehabilitacji, Kultury Fizycznej
i Integracji Społecznej PAN

Sławomir JANDZIŚ, Mariusz MIGAŁA

Rys historyczny rozwoju rehabilitacji w Polsce i na świecie



Opole 2015



*Międzynarodowy Dzień Inwalidy
„Życie bez bólu” (1991–2019)*

Who's Who in the World in Physiotherapy

pod redakcją
Zbigniewa Śliwińskiego i Grzegorza Śliwińskiego
przy współpracy
Zofii Śliwińskiej i Lecha Karbowskiego

fizjoterapia polska

POLISH JOURNAL OF PHYSIOTHERAPY

OFICJALNE PISMO POLSKIEGO TOWARZYSTWA FIZJOTERAPII
THE OFFICIAL JOURNAL OF THE POLISH SOCIETY OF PHYSIOTHERAPY

NR 3(202) (2) KWARTALNIK ISSN 1642-8136

Physiotherapeutic procedure in a patient after the first artificial heart implantation in Poland – SynCardia Total Artificial Heart (TAH)



Postępowanie
fizjoterapeutyczne
u pacjenta po pierwszej
w Polsce implantacji
sztucznego serca – SynCardia
Total Artificial Heart
(TAH)

The effect of hippotherapy on children with autism – physical and psychological factors
Wpływ hipoterapii na wybrane czynniki fizyczne i psychologiczne u dzieci z autyzmem

ZAMÓW **PRENUMERATĘ!**
SUBSCRIBE!

www.fizjoterapiapolska.pl
prenumerata@fizjoterapiapolska.pl





KALMED
Iwona Renz, Poznań

ARTROMOT®
WYŁĄCZNY PRZEDSTAWICIEL
WWW.KALMED.COM.PL



SPRZEDAŻ I WYPOŻYCZALNIA ZMOTORYZOWANYCH SZYN CPM ARTROMOT®

Nowoczesna rehabilitacja CPM stawu kolanowego, biodrowego, łokciowego, barkowego, skokowego, nadgarstka oraz stawów palców dłoni i kciuka.



ARTROMOT-H



ARTROMOT-F



ARTROSTIM
FOCUS PLUS

ARTROMOT-K1 ARTROMOT-SP3 ARTROMOT-S3 ARTROMOT-E2

Najnowsze konstrukcje ARTROMOT zapewniają ruch bierny stawów w zgodzie z koncepcją PNF (Proprioceptive Neuromuscular Facilitation).

KALMED Iwona Renz tel. 61 828 06 86
ul. Wilczak 3 faks 61 828 06 87
61-623 Poznań kom. 601 64 02 23, 601 647 877
www.kalmed.com.pl kalmed@kalmed.com.pl

Serwis i całodobowa
pomoc techniczna:
tel. 501 483 637
service@kalmed.com.pl

NOWOŚĆ W OFERCIE

ASTAR.

PhysioGo.Lite SONO

**NIEWIELKIE URZĄDZENIE
EFEKTYWNA TERAPIA ULTRADŹWIĘKOWA**

Zaawansowana technologia firmy Astar to gwarancja niezawodności i precyzji parametrów. Urządzenie, dzięki gotowym programom terapeutycznym, pomaga osiągać fizjoterapeucie możliwie najlepsze efekty działania fal ultradźwiękowych.

Głowica SnG to bezobrotowe akcesorium o dużej powierzchni czopa (17,3 cm² lub 34,5 cm² w zależności od wybranego trybu działania). Znajduje zastosowanie w klasycznej terapii ultradźwiękami, fonoforezie, terapii LIPUS i zabiegach skojarzonych (w połączeniu z elektroterapią).



wsparcie merytoryczne
www.fizjotechnologia.com



ul. Świt 33
43-382 Bielsko-Biała

t +48 33 829 24 40
astarmed@astar.eu

**POLSKI
PRODUKT**  **WYBIERASZ
I WSPIERASZ**

www.astar.pl

B.A.D



B.A.D. - Boundaries Are Dumb - the first Polish clothing brand dedicated primarily to all amputees but also to all open-minded people who do not recognise the limitations in their lives. In order to meet the expectations of our clients, we are creating the highest quality products with special reinforcements and systems to facilitate the removal. Stylish and comfortable - our modern, minimalist design emphasises the versatility of B.A.D. All products are made in Poland from first-class cotton because customer satisfaction is crucial to our brand. Say **“No”** to limitations and answer the question, “Are you ready to be **B.A.D.?**”



**DESIGNED FOR PEOPLE
WHO CAN'T RESPECT
BOUNDARIES**



MEN'S FABRIC TROUSERS FOR RIGHT AND LEFT-SIDED FEMORAL PROSTHESIS USERS

Comfortable fabric trousers that look ordinary on the surface. However, thanks to the **use of a zipper** placed under the flap on the outside of the left or right leg will allow you to unfasten and **freely adjust the prosthetics** without having to remove your pants

They are fitted with an elastic waistband supported by string to tie. The product is made of high-quality fabric with the addition of elastane, which improves the comfort of use.

There is a short zipper on the left leg in the inner seam at thigh level, **allowing easy access to the valve.** An additional improvement is a **reinforcement** in the area of the bend in the knee and on the back of the pants, the place that wears off more quickly from sitting.



bad_clth_
www.instagram.com/bad_clth_/



BAD - Boundaries Are Dumb
www.fb.com/BoundariesAreDumb

www.bad-clth.com





PZWL



Pomoc w optymalizacji procesu rehabilitacji **chorych z COVID-19**

w różnych okresach choroby i zdrowienia z uwzględnieniem zachowania **zasad bezpieczeństwa** związanych z potencjalnym zakażeniem terapeutów

**SPRAWDŹ
TUTAJ**



**TYLKO
W FORMIE
e-booka**



**POBIERZ
DARMOWY E-BOOK**
od PZWL w prezencie!



mindray

healthcare within reach

ULTRASONOGRAFIA

W FIZJOTERAPII



Autoryzowani dystrybutorzy

Mar-Med

+48 22 853 14 11

info@mar-med.pl

Ado-Med

+48 32 770 68 29

adomed@adomed.pl



MAR-MED

OD 1995 ROKU



ADO-MED

APARATURA MEDYCZNA

The 2nd
International Conference On Multidiscipline Education



Kampus
Merdeka
INDONESIA JAYA



Empowering Minds: Navigating the Future of Education

Keynote Speakers



Prof. Kerry J. Kennedy
The Then Hong Kong Institute of Education
Hongkong



Prof. Madya Ts. Dr. Arumugam Raman
Universiti Utara Malaysia
Malaysia



Prof. George Arief D Liem
Nanyang Technological University
Singapore



Dr. Erick Burhaein, M.Pd., AIFO
Universitas Ma'arif Nahdlatul Ulama Kebumen
Indonesia

Sub Theme

- Blended Learning
- Curricula
- Early Childhood Education
- Educational Leadership
- Educational Psychology
- Education Programs and Teaching
- Foreign Languages Education
- Health
- Health Education
- Higher Education
- Innovative Methodologies in Learning
- Learners of Qualitative Research
- Learning Environments
- Methodology of Sport and History of Physical Culture and Sport
- Multimedia in Digital Learning
- Physical Activity and Health
- Physical Education
- Public Health
- Sport Sciences
- Teaching and Assessment
- Teaching Disability
- Virtual and Augmented Reality Learning Environments

Opening Remarks



H. Herman Suherman, S.T., M.A.P.
Regent of Cianjur
Indonesia



Deden Nasihin, S.Sos.I., M.K.P.
Deputy Chairman of DPRD
Cianjur, Indonesia



Prof. Dr. H. Dwidja Priyatno, SH., MH., Sp.N.
Rector of Suryakencana University
Cianjur



Dr. H. Munawar Rois, M.Pd.
Dean of Faculty Of Education Teacher Training
Suryakencana University Cianjur

Publication Options

- International Journal of Learning Teaching and Educational Research (Scopus Q3)
- Hong Kong Journal of Social Sciences (Scopus Q4 dan WOS)
- International Journal of Disabilitas and Health Sciences (Scopus Q4)
- Polish Journal of Physiotherapy (Scopus Q4)
- The International Society for Technology Educational and Sciences (ISTES) (Proceedings are indexed in Scopus)
- Al-Ishlah : Jurnal Pendidikan (Terindeks SINTA 2)
- Edu Sportivo: Indonesian Journal of Physical Education (Terindeks SINTA 2)
- Journal Elemen (Terindeks SINTA 2)
- Journal Sport Area (Terindeks SINTA 2)
- Journal Teori dan Aplikasi Matematika (Terindeks SINTA 2)

Conference Registration Fees



General Participants 50 K
Host Student Presenters 400 K
General Presenters 500 K



Bank Mandiri | BNI | BRI
1820006898530 | 0622468257 | 010501070965501
a.n D. Nurfaejrin Ningsih

Important Dates

18 March 2023 - 30 May 2023 Abstract Arrangement | 1 June 2023 - 20 July 2023 Full Paper Acceptance | 1 - 20 July 2023 Payment Due | 20 July 2023 - 2 August 2023 Full Paper Review | 27 July 2023 - 2 August 2023 Announcement of Full Paper Accepted | 5 August 2023 Conference Day | September - December 2023 Full Paper Publication

Contact Person +62 877-7879-4797 Rani Sugiarni

Organized by:

Faculty of Education Teacher Training
Suryakencana University Cianjur, Indonesia

Jl. Pasir Gede Raya, Bojongherang, Cianjur, Indonesia



Dr. Comfort®

Nowy wymiar wygody.

Obuwie profilaktyczno-zdrowotne
o atrakcyjnym wzornictwie



APROBATA
AMERYKAŃSKIEGO
MEDYCZNEGO
STOWARZYSZENIA
PODIATRYCZNEGO



WYRÓB
MEDYCZNY

**Stabilny, wzmocniony
i wyściełany zapiętek**
Zapewnia silniejsze
wsparcie łuku
podłużnego stopy

**Miękki, wyściełany
kołnierz cholewki**
Minimalizuje podrażnienia

Wyściełany język
Zmniejsza tarcie
i ulepsza dopasowanie

Lekka konstrukcja
Zmniejsza codzienne
zmęczenie

**Antypoślizgowa,
wytrzymała podeszwa
o lekkiej konstrukcji**
Zwiększa przyczepność,
amortyzuje i odciąża stopy

**Zwiększona
szerokość i głębokość
w obrębie palców
i przodostopia**
Minimalizuje ucisk
i zapobiega urazom

**Wysoka jakość materiałów
- oddychające siatki i naturalne skóry**
Dostosowują się do stopy,
utrzymują je w suchości
i zapobiegają przegrzewaniu

Trzy
rozmiary
szerokości

Podwyższona
tęgość

Zwiększona
przestrzeń
na palce

**Ochronna przestrzeń
na palce - brak szwów
w rejonie przodostopia**
Minimalizuje możliwość zranień

WSKAZANIA

- haluksy • wkładki specjalistyczne • palce młotkowate, szponiaste • cukrzyca (stopa cukrzycowa) • reumatoidalne zapalenie stawów
- bóle pięty i podeszwy stopy (zapalenie rozciągniętej podeszwy - ostroga piętowa) • płaskostopie (stopa poprzecznie płaska)
- bóle pleców • wysokie podbicie • praca stojąca • nerwiak Mortona • obrzęk limfatyczny • opatrunki • ortezy i bandaże • obrzęki
- modzele • protezy • odciski • urazy wpływające na ścięgna, mięśnie i kości (np. ścięgno Achillesa) • wrastające paznokcie



ul. Wilczak 3
61-623 Poznań
tel. 61 828 06 86
fax. 61 828 06 87
kom. 601 640 223, 601 647 877
e-mail: kalmed@kalmed.com.pl
www.kalmed.com.pl



www.butydlazdrowia.pl

www.dr-comfort.pl



26. Sympozjum Sekcji Rehabilitacji Kardiologicznej i Fizjologii Wysiłku Polskiego Towarzystwa Kardiologicznego

11-13 maja 2023, Wisła, Hotel STOK

www.rehabilitacja2023ptk.pl



Rehabilitacja kardiologiczna i fizjologia wysiłku – zapraszamy do rejestracji na wyjątkową konferencję w Wiśle

W dniach 11–13 maja w Hotelu Stok Wiśle odbędzie się wyjątkowe i interdyscyplinarne spotkanie specjalistów z całej Polski – 26. Sympozjum Sekcji Rehabilitacji Kardiologicznej i Fizjologii Wysiłku Polskiego Towarzystwa Kardiologicznego. Serdecznie zapraszamy do rejestracji.

26. Sympozjum Sekcji Rehabilitacji Kardiologicznej i Fizjologii Wysiłku Polskiego Towarzystwa Kardiologicznego to coroczne spotkanie specjalistów, zajmujących się rehabilitacją kardiologiczną, prewencją chorób układu krążenia i innymi formami aktywności fizycznej, która ma prowadzić do poprawy stanu naszego zdrowia.

Ta trzydniowa konferencja przeznaczona jest dla lekarzy kardiologów, specjalistów rehabilitacji medycznej oraz innych specjalności, którzy w swojej codziennej praktyce zajmują się rehabilitacją i fizjologią wysiłku, ale także dla fizjoterapeutów, pielęgniarek, techników i przedstawicieli innych zawodów medycznych, zainteresowanych tematyką spotkania, oraz studentów.

Jakie tematy zostaną poruszone podczas konferencji?

26. Sympozjum Sekcji Rehabilitacji Kardiologicznej i Fizjologii Wysiłku to konferencja, na którą zaproszeni zostali wybitni specjaliści z dziedziny kardiologii i nie tylko. Podczas wydarzenia wygłoszonych zostanie prawie 100 wykładów merytorycznych w ciągu aż 20 sesji. Uczestnicy będą mieli również szansę na udział w sesjach przypadków klinicznych, intensywnych warsztatach, a także panelach dyskusyjnych. To wydarzenie cechujące się dużą interdyscyplinarnością, dlatego z pewnością każdy znajdzie coś dla siebie.

Podczas wydarzenia kompleksowo pochylimy się nad dziedziną rehabilitacji kardiologicznej i fizjologii wysiłku. Wśród tematów wiodących znajdują się:

- rehabilitacja w dobie pandemii i po pandemii COVID-19;
- telerehabilitacja i rehabilitacja hybrydowa;
- rehabilitacja kardiologiczna w specyficznych grupach pacjentów;
- programy KOS-zawał i KONS;
- nowe standardy ESC, PTK i SRKiFW;
- Testy wysiłkowe i testy spiroergometryczne
- monitorowanie wysiłku fizycznego;
- prewencja pierwotna i wtórna chorób sercowo-naczyniowych;
- farmakoterapia pacjentów rehabilitowanych kardiologicznie i nie tylko;
- sport i aktywność sportowa w kardiologii;
- czynniki ryzyka chorób układu krążenia.

Program merytoryczny wydarzenia jest niezwykle bogaty i angażujący. Warto podkreślić także, iż na konferencji pojawią się specjalne sesje wykładów prowadzone przez zaproszone sekcje i ассоjacje Polskiego Towarzystwa Kardiologicznego, m.in. Sekcję Kardiologii Sportowej, Ассоjację Niewydolności Serca, Ассоjację Elektrokardiologii Nieinwazyjnej i Telemedycyny, Sekcję Pielęgniarstwa Kardiologicznego i Pokrewnych Zawodów Medycznych, „Klub 30”, Sekcję Farmakoterapii Sercowo-Naczyniowej, Sekcję Prewencji i Epidemiologii, a także Polskie Towarzystwo Medycyny Sportowej.

„Pandemia wymusiła na nas zmianę paradygmatu rehabilitacji kardiologicznej”

Organizatorami wydarzenia są wydawnictwo naukowe Evereth Publishing oraz Sekcja Rehabilitacji Kardiologicznej i Fizjologii Wysiłku Polskiego Towarzystwa Kardiologicznego (SRKiFW). Przewodniczącą Komitetu Naukowego jest prof. dr hab. n. med. Małgorzata Kurpesa, Wiceprzewodniczącymi – prof. dr hab. n. med. Anna Jagier, dr hab. n. med. Dominika Szalewska, a Komitetu Organizacyjnego – dr n. med. Bartosz Szafran.

Dr n. med. Agnieszka Mawlichanów, Przewodnicząca SRKiFW, podkreśla, iż ostatnie Sympozjum miało miejsce w 2019 r. w Wiśle. W tym czasie udało się zorganizować wydarzenie w formule online, jednak zdaniem Przewodniczącej obecnie „wszyscy spragnieni jesteśmy spotkania osobistego, wymiany doświadczeń i bezpośrednich rozmów, nie tylko na sali wykładowej, ale i w kulturalach”.

– Cztery lata w sporcie to pełna olimpiada, a w naszej dziedzinie kardiologii można powiedzieć – cała wieczność. Pandemia wymusiła na nas zmianę paradygmatu rehabilitacji kardiologicznej, między innymi stworzyła pole dla rozwoju modelu hybrydowego i monitorowanego telemedycznie. W tym czasie ukazało się wiele ważnych dokumentów, stworzonych przez polskie i europejskie towarzystwa kardiologiczne, dotyczące rehabilitacji, prewencji i aktywności fizycznej. Dynamicznie w naszym kraju rozwija się też program KOS-zawał, przynoszący liczne korzyści, ale też budzący kontrowersje. O tym wszystkim i jeszcze wielu innych sprawach pragniemy podyskutować w czasie naszego majowego spotkania – zapowiedziała dr Mawlichanów.

Rejestracja na 26. Sympozjum Sekcji Rehabilitacji Kardiologicznej i Fizjologii Wysiłku możliwa jest na stronie internetowej konferencji rehabilitacja2023ptk.pl/rejestracja/. Informacje na temat opłaty zjazdowej i wydarzeń towarzyszących znajdują się tutaj: rehabilitacja2023ptk.pl/oplata-konferencyjna/.

Informujemy jednocześnie, iż liczba miejsc na konferencji jest ograniczona, dlatego warto zarejestrować się już dzisiaj.
Serdecznie zapraszamy do Hotelu Stok w Wiśle!

**Dołącz do najstarszego polskiego
towarzystwa naukowego
zrzeszającego fizjoterapeutów.**

Polskie Towarzystwo Fizjoterapii

od 1962 roku jako sekcja PTWzK

od 1987 roku jako samodzielne stowarzyszenie



- członek WCPT 1967-2019
- członek ER-WCPT 1998-2019
- projektodawca ustawy o zawodzie fizjoterapeuty (lipiec 2014)

Pracujemy w:

- 15 oddziałach wojewódzkich
- 10 sekcjach tematycznych

Odwiedź nas na stronie:

www.fizjoterapia.org.pl

i rozwijaj z nami polską fizjoterapię

Deklaracja członkowska dostępna jest w zakładce w menu strony

Biotensegration of malocclusion and temporomandibular disorders with diseases of the musculoskeletal system and posture defects – a literature review, supported by own observations

Biotensegracja wad zgryzu i zaburzeń czynnościowych stawów skroniowo-żuchwowych z chorobami narządu ruchu i wadami postawy – przegląd piśmiennictwa poparty obserwacjami własnymi

Danuta Lietz- Kijak^{1(A,B,E,G)}, Aleksandra Bitenc-Jasiejko^{1(B,E,G)}, Piotr Skomro^{1(B,E,G)}, Krzysztof Konior^{2(B,E,G)}, Konrad Kijak^{3(E,F,G)}, Małgorzata Kowacka^{4(E,F,G)}, Zbigniew Śliwiński^{5(A,B,E)}

¹Department of Propaedeutic, Physical Diagnostics and Dental Physiotherapy, Faculty of Medicine and Dentistry Pomeranian Medical University, Szczecin, Poland

²Doctoral Study at the Department at the Propaedeutic, Physical Diagnostics and Dental Physiotherapy, Faculty of Medicine and Dentistry, Pomeranian Medical University, Szczecin; Medical Center in Nowogard, Poland

³Student Scientific Society at the Department and Clinic of Internal Diseases, Angiology and Physical Medicine, Medical University of Silesia in Katowice, Poland

⁴Independent Public Health Care Complex in Żarki, Poland

⁵Department of Physiotherapy, Collegium Medicum UJK in Kielce, Poland

Abstract

Although it enables a quick evaluation, medical diagnostics of the human myofascial-skeletal system is not always used to its full capabilities. This often hampers the objective assessment of a dysfunction and limits treatment options. Dysfunctions of the stomatognathic system, of different aetiology, are becoming more prevalent. Many scientific studies are in to relationships between organs and systems of the human body that can cause dysfunctions of the temporomandibular disorders. Studies investigating correlations between a myofascial-skeletal disorder and temporomandibular disorders are to be found in the literature. To achieve a good understanding of aetiology of these dysfunctions, a holistic view of a patient is advisable, which takes into consideration the fact that the temporomandibular joint is an integral part of the whole body. This study is a review of the literature, supported by own examples on investigating the relationship between myofascial-skeletal disorders and a temporomandibular disorder, which, in the light of current medical knowledge, is attributed to the phenomenon of biotensegrity. A conclusion has been put forward on the basis of 136 scientific reports that a dependency exists between the said dysfunctions.

Keywords

biotensegrity, disorders of the musculoskeletal system, posture defects, dysfunctions of the masticatory system, temporomandibular joint disorder

Streszczenie

Diagnostyka medyczna układu mięśniowo-powięziowo-szkieletowego człowieka, choć umożliwia szybką ocenę, nie zawsze jest w pełni wykorzystywana. Utrudnia to często obiektywną ocenę dysfunkcji i ogranicza możliwości leczenia. Coraz powszechniejsze stają się dysfunkcje układu stomatognatycznego o różnej etiologii. Wiele badań naukowych poświęconych jest związkowi między narządami i układami organizmu człowieka, które mogą powodować dysfunkcje w obrębie stawów skroniowo-żuchwowych. W literaturze można znaleźć prace badające korelacje pomiędzy zaburzeniem mięśniowo-powięziowo-szkieletowym a zaburzeniami skroniowo-żuchwowymi. Dla dobrego zrozumienia etiologii tych dysfunkcji wskazane jest całościowe spojrzenie na pacjenta, uwzględniające fakt, że staw skroniowo-żuchwowy jest integralną częścią całego organizmu. Niniejsza praca jest przeglądem piśmiennictwa, popartym własnymi przykładami, dotyczącym badania związku zaburzeń mięśniowo-powięziowo-szkieletowych z zaburzeniem skroniowo-żuchwowym, które w świetle aktualnej wiedzy medycznej przypisuje się zjawisku biotensegracji. Na podstawie 136 doniesień naukowych wysunięto wniosek, że istnieje zależność między tymi dysfunkcjami.

Słowa kluczowe

biotensegracja, zaburzenia narządu ruchu, wady postawy, dysfunkcje narządu żucia, zaburzenia stawu skroniowo-żuchwowego

Introduction

Disorders of the musculoskeletal system

Musculoskeletal complaints are one of the commonest health issues, which comprises a vast range of medical problems and thus they affect millions of workers reporting pain in the joints, spine and muscles [1–4]. Pains resulting from musculoskeletal disorders (MSD) are caused by are caused by tension disorders in myofascial structures, changes in osteoarticular structures and neurological problems, etc. Consequently, lead to the loss of overall ability and a deterioration in the quality of life. MSD usually affect the neck (the cervical section of the spine), back of the neck, shoulders and lower limbs [5–13]. They can also relate to the pain of the back (especially the lower back), lower limbs, feet, caused by chronic strain of the organism [14–22]. They can be provoked by injuries (including chronic overload injuries) and be the result of systemic diseases affecting the condition of the musculoskeletal system [23–26]. Starting as slight pains, these health problems transform into severe dysfunctions that often lead to partial disability. External factors contributing to MSD may be: the type of work performed (both sedentary and heavy physical activity), physical activity and a number of individual factors (weight, posture defects, improper lifestyle) [27–46]. Most MSD develop over time. It is often difficult to find one particular cause of the disorder, and the inducing factors may be varied [47]. For example, there are many indications that deformations and degeneration of the spine, and consequently the significant pain caused, are a consequence of undetected posture defects in young people [48,49], whose only symptom is the visual asymmetry of the body, with an absence of visible symptoms [50] and motor activity [51, 52].

Periodic and screening diagnostics should be carried out on an individualized manner, especially in relation to the patient's age [53–55]. The total development of the posture function takes place at the age of about 11–12 years, and remains stable until approximately 65 years of age [56]. Consequently, the conclusion should be based on the assessment of the entire posture, taking into account, in particular, the structural integration (the so-called tensegration) of all the elements of the locomotor system, without neglecting the visceral tissue. An important aspect of postural evaluation is also the examination of balance, which is carried out with a pedobarograph [57]. The essence of postural defect prophylaxis is periodical and requires the full evaluation of the posture, walking pattern and compensation mechanisms [58–60]. The detailed diagnosis in clinical conditions of spinal curvature in the three-plane view, assessment of shoulder asymmetry in relation to the hips, the examination of mobility ranges (bend, straightness, etc.), and evaluation of gait are recommended [61]. In addition to the assessment of global patterns, the entire body posture should also be assessed segmentally, i.e. lower limbs, pelvis, trunk (including the spine), shoulder position, head (including craniofacial), upper limbs. [62]. However, the assessment of the causes of defects and deformations should not be neglected, differentiating the problems originating from the osteoarticular, muscular and nervous systems [63, 64], taking into account systemic diseases and the patient's weight. The human posture consists of a position in space and aims to keep the bo-

dy in balance, both while standing and while locomotion. It is an involuntary process, maintained by skeletal muscle contraction, coordinated by stimuli and neuromuscular regulation. It is functional when all systems are in harmony with the so-called kinetic chains, which are painless, and with a correct relationship of the individual body segments in three planes. Thus, a body that is not in structural equilibrium would be characterized by imbalance and consequent pain [65, 66].

The stomatognathic system, together with the maxilla and mandible, take an active role in maintaining the correct position of the skull with the help of the neck flexors and extensors, the hyoid, masticatory, as well as shoulder girdle muscles. It has been demonstrated that mandible activity changes the electrical capacity of the postural muscles, i.e. vertebral, temporomandibular, cervical spine and shoulder girdle [67–69]. Malocclusion and weakening of the oral cavity muscles may lead to changes in the frontal and sagittal position, which may result in scoliosis or apparent asymmetry of limb length. An imbalance of masticatory muscles disrupts the kinematic chain of the oral cavity and changes the posture, affecting pain in the lower back. The relationship between dental occlusion is justified by functional and anatomical compounds of the masticatory system and the process of body balance regulation [70, 71]. Unconscious postural reactions are associated with nociceptive reflexes resulting from contact between tooth surfaces [72, 73]. It has been frequently indicated that dental braces do not prevent the progress of spinal curvature in adolescents as well as deformities in adults [74–79]. The direct relationship between the masticatory organ and the cervical segment of the spine is not questionable. Cervical and occipital pain is considered to be the next major challenge for the global civilization [80–86]. Positioning the head in protraction (FHP – Forward Head Posture) was defined as an epidemic concern of modern times. The forward movement of the head through biokinematic compounds causes a change in the center of gravity of the body. The shoulders are positioned in protraction, thoracic kyphosis and lumbar lordosis will deepen. The pelvis is tilted anteriorly. As a consequence, the position of the lower limbs changes [87–97]. The curve of the spine stabilizes about 5–6 years after the proprioceptive maturation of the feet. The spinal-foot system maintains balance and provides support and resistance to pressure forces [98–101]. Considering civilization's growing problem of spinal defects and pain, its relating consequences on the feet should also be considered. The problems of scoliosis effects on lower extremities and gait have been repeatedly analysed by means of pedobarography, which showed significant effects on the distribution of pressure within the feet. [102–104]. The axial imbalance of the spine affects the asymmetric distribution of the body's equilibrium during standing and walking, which may affect the development of malformations in the lower parts of the body through long-lasting compensating and destabilizing processes [105–107]. In the screening diagnostics of body posture, the use of pedobarography is justified, particularly in the evaluation of posture in terms of tensegrity (examination of balance, gait, etc.) [108]. It would be deemed logical to conclude that the assessment of posture should also take into account the ascending problems, which in this case would be

the impact of foot defects and dysfunctions on the parts of the upper body.

Diseases of the musculoskeletal system often are chronic and stable. Musculoskeletal pains, depending on their localisation, are divided into generalised and local, depending on their duration, into acute and chronic, and depending on the cause, into primary and secondary. Clinically, it is important to differentiate disorders manifested by musculoskeletal pain into inflammatory and non-inflammatory [109]. This allows for a correct diagnosis of the nature of the basic pathological process causing the ailments [110, 111]. The risk of the development of musculoskeletal disorders also increases in the circumstances of operating psychosocial factors, such as time pressure, low level of job satisfaction, lack of [the feeling] of being in control under the condition of high requirements, or inadequate social support [112]. Many authors assume modifying the influence of the psychosocial factors of work on the development of overuse syndromes caused by physical factors. Meanwhile, many occupational factors (e.g. repetitiveness of work tasks, work pace, no rest breaks, etc.) serve as a proof of poor control over working conditions and may be themselves an independent factor of overuse syndromes' development. Stress caused by psychosocial factors may affect human health in two ways that differ in the mechanism and pace of change [113]. The instant path that comprises biological changes taking place in the organism while under stress, whereas the delayed path relates to all fixed changes in the psyche and behaviour of an individual due to cumulative effects of the instant changes in somatic, psychological and behavioural processes. The instant path plays a major role in the development of musculoskeletal disorders. It comprises important changes in the organism such as an increase of catecholamines and cortisol secretion, increase of heart rate activity and skeletal muscles tension, including the neck and back of the neck muscles [114–117]. The following individual characteristics such as age, female gender (number of pregnancies, taking oral contraceptives, menopause, hormone replacement therapy), obesity, low overall physical activity, smoking and alcohol abuse, as well as the following diseases: diabetes mellitus, thyroid insufficiency and rheumatoid arthritis should be taken into consideration while discussing risk factors of musculoskeletal disorders [118–120]. Non-occupational overuse, a type of sport or a hobby constitute another group of factors. Fibromyalgia, whose main characteristic is generalised pain of the musculoskeletal system, should be taken into consideration in differential diagnoses of overuse syndromes. In accordance with the criteria set out by the American College of Rheumatology, fibromyalgia is diagnosed when the pain in many parts of the body lasts at least 3 months and the pain upon compression force is at least 11 out of 18 of the so-called trigger points [121].

The complexity of the problem of myofascial pain has a direct impact on the use of therapeutic methods. Only in the initial phase of pain, releasing techniques are used to relax the tissues, focused on the local problem. A number of physiotherapeutic treatments are implemented, applicable in the course of inflammation, neuralgia, increased tension, i.e. electrotherapy, magnetotherapy, cryotherapy, sonotherapy, thermotherapy, etc.

[122–128]. However, taking into account the tensegrity relationship and the dependence in the course of bio-kinematic chains, many research and scientific works indicate the need to use integrated therapy methods. They mainly include manual techniques to increase functionality of the musculoskeletal structures as well as the whole organism (structural integration). One of the methods of holistic therapy of the musculoskeletal system is (OMT), which includes a range of manipulations, muscle energy techniques (MET), visceral techniques and exercising. These methods are applicable both in acute pain and later stabilization of body posture. [129–135]. Manipulation techniques within the joints are widely used in the therapy of acute conditions (such as HVLA-High Velocity Low Amplitude) [136–139], in the treatment of pain the importance of comprehensive therapy in order to unblock joint blocks to restore proper muscle tone has been repeatedly emphasized. Other methods of integrated manual therapy and rehabilitation exercises are: e.g. Kaltenborn-Evjenth method, McKenzie method, S-E-T method, the Mulligan's concept [140–147]. Kinesio Taping (KT) used in the treatment of diseases of the musculoskeletal system, serves sensory and proprioceptive effects, enables the regeneration of places affected by the disease process, giving the effect of improving microcirculation. It has an extremely beneficial effect on microcirculation and the lymphatic system. Improving the functionality of structures [148–166].

Research and scientific works also indicate that the methods of tensegrity therapy take into account neuromobilization. They allow to restore the movability of the nerve in relation to the structures surrounding the nervous tissue. They reactivate the stretching and tensing capabilities of the nervous tissue itself. They should include tissues with pain as well as those that are functionally related to pathology [167–169]. In many countries, acupuncture is one of the methods of integrated therapy of the musculoskeletal system, however, its use still raises many controversies [170–180]. The final therapeutic stabilization of the patient also includes whole-body therapy in most cases. In addition to the previously indicated methods, Yoga, also based on the posture tensegrity model, is often used. In addition to stabilizing the central posture, it also takes into account the effect supporting the psychoemotional state, breathing functions, etc. [181–185]. Stability training for patients with a musculoskeletal disorder reduces pain, instability and prevents setbacks [186–190]. Experiencing pain of the skeletal or muscular system as well as the joints, patients may repeatedly use pharmaceuticals: paracetamol or tramadol, or substances of the non-steroidal anti-inflammatory drugs class (NSAIDs) such as ibuprofen, diclofenac, ketoprofen, metamizole or dexketoprofen [191–194]. The above study of the literature indicates that the key therapeutic approach in the problems of pain in the musculoskeletal system is a holistic therapy that takes into account the biotension of the body posture. Therapy based on structural integration is referred to as tensegrity therapy.

Objectives

The aim of this study is to highlight an essential issue, which is a comprehensive diagnosis of functional disorders of the locomotor system of the masticatory system. Additionally, this would entail the need for the inclusion in the clinical assessment of

diagnostic procedures of the whole-body posture, the examination of balance and gait and the process of therapeutic planning, as well as significant elements of the assessment of structural integration. The aim is to also provide the reader with a comprehensive overview, supported by literature and individual clinical cases.

Methodology of meta-analysis

An extensive search of the materials used in this manuscript was conducted online using PubMed, the Cochrane, Medline, Web of Science and EMBASE databases. The results obtained were evaluated and verified for correct qualification by key words. Out of 967 manuscripts, 330 were used in the final concept, including 194 studies which were used in Introduction. The remaining 136 publications were included in the analysis, after assessing their quality.

Main Issue

The term **tensegrity** refers to manual therapy based on structural integration. The first mention of the role of deep fascia in low back pain was made in 1939 [195], while modern medicine has identified it to be one of the main causes on non-injury back pain [196–199]. When tension in the tissue becomes excessive, a damaging stimulus arises. Then, the stimulus is propagated linearly in the human body [200–202]. Hence the conclusion that pain can appear in a place distant from where the pain stimulus first arose [203–207]. This points to the fact that restoring tension of anatomical bands is an indispensable process of rehabilitation and pain therapy, as well as of the restoring of structural integration. This enables the generation of greater force and handling overloading, which results in increased amortisation and help for the muscles that stabilise the spine and are closely linked with a specific fascial band [208, 209]. The concept of fascial bands allows the connections of musculo-fascial lines that influence a dysfunction within the whole posture to be traced. Thereby, including these lines in therapeutic procedures heavily influences the efficacy of the therapy. This holistic approach to the fascial system is based on the still ongoing research into fascial anatomy. A literature study in the area clearly indicates that only a few authors treat the fascia as a three-dimensional system [210]. Nevertheless, it has been repeatedly indicated that manual therapy based on the tensegrity concept (structural integration) is one of the most effective methods of balancing tension within the movement system [211, 212].

Tensegrity (tension-integrity), i.e. structural relationship, where forces are not localised but transferred, which is a theory that attempts to describe experimentally the relationship between musculoskeletal disorders and a dysfunction of the temporomandibular joint. This is a term invented by Buckminster Fuller and Kenneth Snelson. The term explains the way the human organism acts as a whole and how the bones of our skeleton, thanks to the tensions of tendons, ligaments and muscles, allow the body to maintain a vertical posture against the force of gravity. In the tensegrity model of the human body, the bones constitute a compression system suspended in the system of constant tensions comprised by the myofascial system. The

fascia, which is connective tissue, has a key role. It surrounds the body's organs as a scaffolding, transferring tensions along the fibres [213–216]. In the case of the temporomandibular joint (TMJ), just the joint capsule and capsular ligament transfer signals through the proprioceptive system because the surface of the joint and the mandibular disc per se do not have pain receptors [217, 218]. According to the theory of tensegrity, in the case of excessive tension on the tissue, a damaging stimulus arises that is then propagated linearly along the whole organism. As a result, ailments can appear in a place distant from where the stimulus is operating. According to Still, the fascia constitutes the place in a human body where the causes of diseases can be found, and that the proper fascia functioning guarantees life, whereas its dysfunction leads to the death of the body [219]. Ultrasound examinations revealed that during oscillatory moves e.g. during walking, the fascia often acts as a mechanical energy store [220].

A good body posture is closely linked to an exemplary and healthy musculoskeletal system. It is due to the fact that the posture is a system of points and segments of the body set against each other without any pathological lesions, which guarantee its stabilization and balance at a minimum physical effort and providing optimal static-dynamic efficiency, and ensures the proper development and functioning of the internal organs. Any deflection from a good body posture is called a faulty posture. Faulty postures can be congenital or acquired [221–223]. Congenital body postures may result from metabolic diseases, genetic defects or disorders caused by mechanical, toxic or specific intrauterine factors during foetal development. On the other hand, the acquired result from the body being exposed to the influence of external factors after foetal life, include various diseases of the musculoskeletal or nervous system. The latter include such factors as a sitting lifestyle, unhealthy eating habits and minimal physical activity [224–226]. Implications of bad postures can be clinical or economic, as well as social and societal [227]. Importantly, it should be expected that musculoskeletal disorders will become more prevalent along with a growing number of overweight and obese people in the overall population [228]. Obesity strongly correlates with a higher percentage of bad postures. Unfortunately, according to the WHO data, this is still an upward trend [229–230]. Faulty postures is a broad concept, which includes defects of the skeletal, ligament and muscular system. They may concern the spine, shoulder girdle, chest, pelvis, thigh joints and lower extremities including knee joints, ankle joints and metatarsus [231, 232]. Furthermore, defects per se can be singular or multiple [233–238]. An early diagnosis and prompt adoption of remedial measures in the form of a corrective surgery, kinaesthetic rehabilitation or other physiotherapeutic methods may contribute to the restoration and then strengthening of right patterns of physical activity [239].

Dysfunctions of the masticatory system

The stomatognathic system is a morphological and functional set of tissues and organs of the oral cavity and facial skeleton, which, in its entirety, is the biofunctional system responsible for functions such as masticating, articulating sounds, breathing and expressing emotions non-verbally [240].

It can be divided into the dental and alveolar structure, comprising teeth and the periodontium; a dental and dental syndrome, determining the interrelation between the teeth of the upper and lower dental arch (occlusion, articular states of the mandible); and a musculo-articular structure, comprising the right and left temporomandibular joints (TMJ) and the masticatory muscles. The stomatognathic system is also comprised of the facial skeleton bones, suprahyoid muscles, mimic muscles, the muscles of the tongue and palate, blood and lymph vessels, cranium nerves (i.e. trigeminal, facial, glossopharyngeal, hypoglossal) and salivary glands (including large paired salivary glands: parotid, sublingual and submandibular) [241]. While examining the stomatognathic system of a patient, each dentist takes their history, and performs a physical and additional examination. Particular attention should be devoted to an extra-oral palpation examination, during which anatomical structures such as the muscles: superficial temporal, masticator, sternocleidomastoid, trapezius, deltoid, suprahyoid and suboccipital muscles, as well as the temporomandibular joint should be thoroughly examined. Also, the face symmetry should be analysed, as well as lips' competence, upper airways' patency, extent of the movement of the mandible (including abduction trajectory), acoustics of the temporomandibular joint and the outlet of the trigeminal nerve for neuropathy [242]. Dysfunctions of the stomatognathic system are more prevalent and their aetiology is diverse. Despite being quick, the diagnostics of the masticatory organ dysfunctions is not always used up to its full capability limiting the possibility of obtaining an objective assessment of the dysfunction and relevant treatment options [243]. Many medical specialisations focusing on the treatment of specific dysfunctions have developed, yet a holistic approach to a patient provides an opportunity to choose the right treatment for the patient individually, depending on the pathological condition diagnosed in their stomatognathic system [244]. With regard to the diversity of malocclusions, more and more questions and dilemmas arise about the anatomical structures contributing to the development of these defects.

The specialists should examine the patient's posture, their gait, the way they move, and other individual biomechanical parameters of the musculoskeletal system [245–250]. Many studies are available that have been investigating relationships between organs and systems in the human body that could affect the functioning of the masticatory system. Bringing in the concepts of Anatomy Trains and tensegrity gives even the possibility of linking a dysfunction of the temporomandibular joint to defects in the foot architecture, linking the beginning with the end of the myofascial chain [251]. Patients with dysfunctions of the masticatory system often report ailments in other parts of their bodies, even including those in the joints of the lower limbs and feet [252, 253]. According to the concept of Anatomy Trains, the following anatomical bands can be identified: anterior band, posterior band, spiral and lateral band as well as the deep band. The deep band has a direct connection between the aspect and architecture of the foot and the temporomandibular joint through the following anatomical structures: the tibialis posterior that stabilises the longitudinal and transverse foot arches; the long flexor of the toe that strengthens the foot arches and, additionally, supination (valgity prevention), and

their linear connection with the suprahyoid muscles, the mandible and the manubrium, in which the insertion of the sternocleidomastoid muscles is localised [254, 255].

Studies into the relationship between a musculoskeletal dysfunction and a malfunctioning of the TMJ are available in the literature. However, a holistic view of the patient, and paying attention to the fact that the TMJ is an integral part of the whole human body would allow a better understanding of the aetiology of these disorders to be gained [256]. Due to the multiple aetiology of the temporomandibular joint dysfunctions (TMD), the issue remains complex and thus providing the right diagnosis and treatment options is problematic [257, 258]. From the medical point of view, diagnosing patients with the said disorders lies in describing the relationships between distant anatomic structures through transferring forces using soft tissues such as tendons, ligaments, muscles, fascia and the commissure formed by the osseous tissue. The fascia is the most important element in the above system, being connective tissue that passes strains along the fibres [259, 260]. Learning about these relationships will enable a better informed selection of the therapy to provide a long-lasting therapeutic effect.

Patients are more and more often complaining about problems with the TMD to doctors of various specialisations. After a long-lasting period of neurological, laryngological, orthopaedic and maxillofacial diagnostics, the patients turn to dentists, where a diagnosis is provided, and the attempts of treatment are undertaken to decrease the patients' discomfort [261–263]. Unfortunately, the effects of the treatment do not last long or they rely on using orthopaedic repositioning or relaxing splints. Splints are becoming part and parcel of the therapy that often takes many years [264–269]. The effect of malocclusion on the TMJ has been comprehensively described. However, there remain about 30% of patients with proper indications for the canine teeth and skeleton according to Angle's classification as well as normal occlusion height where problems with the joint are manifested as pain, discomfort, clicking or cracking [270, 271].

It is more often the case that a dentist seeks a physiotherapist's help to complement dental treatment. Integrated dental and physiotherapeutic care may be very effective in the diagnosing process and rehabilitation of patient with the aforementioned disorders and provide long-lasting therapeutic effects. It is possible thanks to the elimination of diagnostic errors as well as influencing all anatomical structures that may be responsible for the problems reported by the patient [272].

Substance of the issue – Evidence for the considerations above

All figures and images have been prepared by the authors and are their property.

Temporomandibular joint disorders (TMD) are not a single diagnosis. They are more a set of diseases that encompass various conditions, and, consequently, various clinical symptoms. TMD affect 15% of adults, with the prevalence peak between 20 and 40 years of age. The problem is twice as common in women. It is also worth mentioning that smoking increases the risk of the disorder in women under 30 years of age [273, 274]. Malfunctioning of the TMJ can be divided into intra- and extra-articular. Adaptive mechanisms are responsible for the fact that most dysfunctions in a joint develop imperceptibly. Such impercepti-

ble development was described as a painless dysfunction of the masticatory organ. Only 5% of patients seeking medical attention complain about specific symptoms. In the light of this fact, it is possible to confirm the development of the symptomatic syndrome of the masticatory organ that can be referred to as a pain masticatory dysfunction, myofascial pain or maxillofacial pain [275, 276]. With regard to the source of the pain, there is a muscular and articular form of the syndrome. Myalgia is constant, dull, stabbing, and difficult to localise. Arthralgia is also dull, diffused, and increases with a change of body position.

The aetiology of TMD is multifarious and not fully understood. It is known that occlusion conditions, injuries of the mandible, parafunctions, dysfunctions, and caries influence the

development of the disorders in the joint concerned [277]. Most recent studies point to muscular tension as one of the more important causes of pathology development in a joint [278]. Long-term muscle overload leads to an improper functioning pattern that is transferred onto the joint, disordering adaptive mechanisms. The pattern of anterior or posterior posture also determines the type of occlusion.

Some cranio-mandibular dysfunctions also cause abnormalities: lateral flexion in the atlanto-occipital joint, secondary compensation in the atlantoaxial joint or restrictions within the hyoid bone. It was shown that this abnormality may cause scoliosis [279], deepening of spinal curvatures and pelvic asymmetry. The image of the stomatognathic system shows the presence of cross-bite, open bite or posterior bite (Fig. 1A and B and Fig. 2).



Fig. 1. A and B. An intraoral photo of a patient who was orthodontically treated due to a posterior bite
All graphics and images were prepared by and remain the property of the authors.



Fig. 2. A patient with shortening of the right limb (43 mm) and lowering of the right iliac ala. In the sagittal plane, the shoulders in protraction, hyperextension of the right knee. In the transverse plane, rotation of the pelvis to the right

Retrusion of the mandible is another example of cranio-mandibular dysfunction causing a shift of the projection of the centre of

pressure to the front. It will correlate with increased lordosis of the cervical and lumbar spine, and anterior pelvic tilt (Fig. 3, 4, 5, 6).



Fig. 3. An intraoral photo of a patient with total supraocclusion.



Fig. 4. In the sagittal plane, increased cervical lordosis, protraction of the shoulders, increased lumbar lordosis – anterior pelvic tilt – hyperextension of the knees, hallux-valgus in both feet




Fig. 5. The result of projection of the centre of pressure onto the ground and foot barycenter, with a shift of the centre of pressure to the front




Fig. 6. The result of the body balance test, with a clear predominance of posterior oscillations as a result of balancing the anterior tilt of the silhouette

On the other hand, shifting the projection of the centre of pressure to the back will cause flattening of cervical and lumbar lordosis, and protrusion of the mandible (Fig. 7, 8 A and B) [280, 281].

While diagnosing TMD, it is important to determine whether this is an ascending or descending dysfunction. The reason for this lies in the fact that the symptoms will manifest themselves in the TMJ, whereas the cause of the dysfunction can be local-

sed apart from the joint. Barr, Zink, Meesserman, Fukuda tests are good tools to assess ascending or descending dysfunctions [282]. In the case of the TMD, it is important to determine a posture pattern, as described by Hall, Wernham and Litteljohn [283]. Anterior and posterior posture patterns can be distinguished causing not only malocclusions but also dysfunctions in the following systems: osteomuscular, respiratory and circulatory, and visceral (Tab. 1).



Fig. 7. An intraoral photo of a patient with inferior prognathism



Fig. 8 A and B. In the sagittal plane, flattened cervical and lumbar lordosis, posterior pelvic tilt. In the transverse plane, external rotation of both lower limbs, the feet are hollowed, adduction of the left forefoot. In the frontal plane, asymmetry of the pectoral muscles resulting from protraction of the left shoulder and, consequently, internal rotation of the upper limb

Table 1. Posture patterns according to Hall

Area	Anterior pattern according to Hall	Posterior pattern according to Hall
the osteomuscular system	<ul style="list-style-type: none"> • a deepened cervical lordosis • increased tension in the cervicothoracic transition • increased tension of back muscles and spinal ligaments • ankylosis and increased tension at Th₁₁ and Th₁₂ • increased tension of the lumbosacral transition 	<ul style="list-style-type: none"> • occipital bone in extension and compression • increased tension in the cervicothoracic transition • a deepened thoracic kyphosis and decreased tension of lower thoracic segment • compression of the sternal articulations • an enlarged lumbar lordosis • an overload of the sacroiliac joint
the circulatory and respiratory system	<ul style="list-style-type: none"> • tension of the diaphragm, often during inspiration • weakened and overly stretched abdominal muscles 	<ul style="list-style-type: none"> • tension of the diaphragm, often during expiration • upset balance between the abdominal cavity and chest pressure • an increase in the tension of the abdominal cavity walls
the visceral system	<ul style="list-style-type: none"> • ptosis of internal organs • loosening of the parietal peritoneum • proneness to hernias and irritation of the true pelvis area 	<ul style="list-style-type: none"> • increased pressure on the abdominal cavity and pelvis organs • proneness to disorders of the circulatory system • proneness to problems with the respiratory system • proneness to constipations

Many clinical studies have been carried out that, based mainly on an interdependence of skull and mandibular dysfunctions on the cervicothoracic and lumbar spine, describe the

relationship between TMD and human posture (Fig. 9 A, B, C, and Fig. 10) [247, 284].



Figures 9 A, B, C. An intraoral photo of a patient with right-sided partial lateral cross-bite and crowded mandibular incisors



Figure 10. A patient with multifaceted posture disorders visible in the sagittal plane (protraction of the head and shoulders, increased cervical and lumbar lordosis, thoracic kyphosis – with simultaneous anterior pelvic tilt and pelvic protrusion (in the pelvis-head line); hyperextension of the knees, external rotation of the limbs with broad-based balance spacing of the feet)



Figure 11. A and B. An intraoral photo of a patient with distocclusion and protrusion of the maxillary incisors



Figure 12. A patient with protraction of the head and shoulders, increased cervical and lumbar lordosis, thoracic kyphosis – with simultaneous anterior pelvic tilt and pelvic protrusion (in the pelvis-head line); hyperextension of the knees, external rotation of the limbs with broad-based balance spacing of the feet



Figure 13. A and B. An intraoral photo of a patient with right-sided partial lateral cross-bite




Figure 14. A and B. In the sagittal plane: protraction of the head, increased cervical and lumbar lordosis, asymmetrical planovalgus feet. In the frontal plane, in the lateral right-sided position, asymmetry of the shoulders, increased internal rotation of the left limb, significant left planovalgus foot, both flat feet

It should also be mentioned that a specific posture pattern of a patient with TMD does not only affect the position of the cervical spine, but also significantly influences the possibility of the trapezius and levator scapulae muscle hyperactivity deve-

lopment and deepens lumbar lordosis. This is directly linked with a pathological and excessive pelvis anteversion and bending of the iliac joints, contributing in this way to the hyperextension of the knees (Fig. 15, 16) [288, 289].



Fig. 15. A patient with mixed dentition, distocclusion and protrusion of the maxillary incisors



Fig. 16. In the frontal plane, the torso flexed to the right, valgus knees. In the transverse plane: increased rotation of the left upper limb, asymmetrical lower limb rotation (increased internal rotation of the right lower limb), adduction of the right foot

Discussion

Studies by Maeda were carried out to reveal a correlation between the deviations of the length of a lower limb and the patient's occlusion. To investigate this correlation a range of tests were done in which patients were alternately raising their heels while motions in their TMJ were being observed. The results confirmed the correlation between the raising of the right heel and shifting occlusion forces to the right [290, 291]. On the

other hand, studies by Bonato were focused on a relationship between the pain in the TMJ and pain in other distant joints: hip, knee, ankle, or wrist. Bonato concluded that the risk of pain transferred into another joint was 5.5 times higher in patients with TMD. The complaints of the patients examined mainly concerned knee joints [292].

With tensegrity theory in mind, Souza et al. assessed the pressure exerted on a foot and its spreading in patients with a dia-

gnosed TMD. The study demonstrated a statistically significant correlation between the spreading of the pressure exerted on the planta and a pathology within the TMJ [293]. The study initiated a discussion on the interdependence between the foot and the stomatognathic system; an interdependence whose existence has been denied by Saito [294]. Valentino is of the opposite opinion and describes planta stimulation that increases tension in the masseter and temporalis muscles [295]. According to Gonzalez and Manns, a head in protraction (when the head is moved forward) is characterised by hyperextension in the upper part of the cervical spine C_1-C_3 as well as increased curvature in its lower segment C_4-C_7 , which leads to hyperlordosis [296]. Correlations between the position of the head and cervical spine and temporomandibular joint dysfunctions were investigated in many studies. It should be stated here that opinions on the issue are divided. Many researchers have observed the dependence [247, 285–291]. Others have not found any significant correlation [296–300]. The studies into the conditions of the joint under investigation should be conducted in accordance with the Research Diagnostic Criteria for TMD's (RDC/TMD) [275]. The head's position is assessed on the basis of cephalometric or photographic measurements of the C_0-C_1 distance as well as a craniocervical angle (CVA). The craniocervical angle serves as a tool for the measurement of the angle between a line marked horizontally through the spinous process of the seventh cervical vertebra (C_7) and a line in the middle of the antilobium [301]. Visscher et al. did not note any important changes in the head's position of the patients with the TMD in comparison to the control group [296].

In the studies by Shweta Channavir Saddu, an additional study group of the TMD patients was created. The group included patients with muscular and articular dysfunctions with an articular disc dislocation. No relationship was confirmed between the head posture and the TMD. Interestingly however, a cervical lordosis was observed in patients with a temporomandibular joint dysfunction of muscular origin [302]. Sonnensen has drawn a similar conclusion. In studies conducted on a group of children, he proved that the head in protraction with concomitant reduced motility and cracking in the joint [299]. Pedroni et al. also confirmed the relationship between joint disorders and the position of the head and the cervical segment [275]. 68% of the subjects in the group of students with TMD had an extension in the cervical segment as well as the arms positioned to the front. A dysfunction of the cervical segment usually manifests itself with pain as well as abnormal motility in the musculoskeletal structure of the cervical spine. Concomitance of a cervical spine dysfunction and TMD has been confirmed by the findings of clinical studies [250, 251, 301]. The researchers coined the term 'the triad of dysfunction', which suggests a relationship between myofascial pain, internal pathologies in the TMJ as well as a musculoskeletal dysfunction of the cervical segment of the spine [259, 285]. Examining this relationship, Bragatto confirmed a higher percentage of the incidence of TMD among women working in the sitting position (at a computer) who reported chronic pain and neck instability [251]. De Wijer points to the necessity of more comprehensive diagnostics for patients with chronic symptoms from the head and neck area [303]. It seems reasonable not to

treat TMD as a local, separated problem, but to also extend diagnostics for the cervical segment of the spine and the shoulder girdle. Studies by Wijer indicate that symptoms from the TMJ and a dysfunctional cervical segment are concomitant. An orthopaedic test for the cervical segment is applied to recognize the source of the pain [303, 304]. In her studies, Weber did not find a relationship between the craniocervical position and TMD. The researcher emphasized the fact that the concomitance of the cervical spine dysfunctions and TMD is related with the common nervous track originating from the trigeminal nerve. Remarkably, in this study, patients with a concomitant condition of a joint experienced pain under palpation in the neck muscles and along with its movements [250, 285]. A study conducted at the Medical University in Zabrze, Poland, is also worth mentioning. Patients experiencing pain and with reduced motility of the cervical segment were the subjects of this study [305]. The subjects underwent the therapy with the relaxing SVED splint. A significant pain reduction in the TMJ, but also in the cervical segment, whose motility increased, was achieved after 3 months of treatment. A motility improvement as regards the cervical segment of the spine substantiates the tensegrity theory in the context of the relationship between the TMJ and the osteo-myo-fascial track of the lower segments of the spine.

A patient's posture pattern, wherein a temporomandibular dysfunction occurred, means more than only the aforementioned head protraction and cervical hyperlordosis, but also hyperactivity of the upper part of the trapezium and the levator scapulae muscle, rounded neck, deepened lumbar lordosis, compensatory bending of the iliac joints, pelvis anteversion and hyperextension of the knee joints. A few theories explain such a distant range of interdependent disorders. One of them was put forward by H. Diers. The researcher divided disorders into ascending (originating in the lower parts of the body) and descending ones (from the TMJ) wherein the tension is transferred by the proprioceptive system.

The myofascial theory, presented by Myers, points to the existence of fascial bands along which information is being transferred. Types of information include pathologies that initially ignite one dysfunction and then abnormalities in even distant organs [306]. Myers divides fascial bands into the posterior surface band that stretches from the planta to the top of the head; the anterior surface band running from the instep to the lateral part of the skull; and the anterior deep band. The bands' role is to maintain the body posture and the centre of gravity. They act as transmitters for stimuli that confirm the existence of the tensegrity model. Pressure or increased tension in one location results in heightened pressure in other tissues, which proves the integrity and interrelation of the myofascial chains with surrounding tissues. Manheim defined myofascial relaxation as facilitation of the mechanical, neuronal and psychophysical adaptive potential that cooperates and manifests itself through the myofascial system [307, 308]. Many researchers proved the effectiveness of myofascial relaxation in the reduction of pain, including the pain of the TMJ, as well as in restoring a good body posture. Both factors increase the level of the patient's quality of life [309–315]. Many research papers were published in which it was evidenced that postural changes may cause or may just predispose a person to the TMD [275, 316–318]. However,

it should be mentioned that these contradicting the aforementioned stance are also present in the literature [319]. Some studies point to the improvement in the TMJ following physiotherapy or postural re-education [318, 320]. In his studies, Saito confirmed the existence of postural changes in the form of pelvis rotation towards the back (pro-posterial), lumbar hyperlordosis, extension in the thoracic segment and pathological leaning of the head to the right, whilst the mandible to the left, with the mouth open. All these symptoms were present in patients with the TMD (with the previous articular disc dislocation) [321]. Wright evaluated a relative risk of the occurrence of the TMD in patients with an asymmetry of the shoulders of about 5.9% [318]. According to Park, performing appropriate exercises could restore normal movement extent in the TMJ in a significant number of patients suffering from mild scoliosis (10 degrees) [322].

Himiko Ikemitsu et al. investigated a correlation between the direction of the movement of the mandible and scoliosis in patients with TMD. Their study indicated only a relationship between TMD and scoliosis. No correlation was found between the direction of the lateral movement of the mandible [323]. Pain in the lower segment of the spine is common. The system of pelvic joints significantly affects the development of the human body's posture. Their lateral relocation results in a lateral curvature of the spine, when the curvature of the pelvis to the back increases lumbar lordosis, whereas to the front it increases thoracic kyphosis. Numerous studies have been conducted to investigate whether such distant structures as the pelvis, namely its disorders, and the TMJ can affect each other. The findings confirm their interdependence [324, 325]. The work by Zonnenberg et al. is a valuable example of such a study. The main objective of their study was to indicate differences in the setting of the pelvis and the neck in a group of patients with a dysfunction of the TMJ in comparison to the control group (patient without pathologies in the TMJ) [317]. A setting of the aforementioned bony structures was evaluated by marking topographic lines between the acromions, and the anterior and posterior pelvis spurs. Zonnenberg showed that patients with TMD have, in comparison to the control group, a statistically disturbed system of the topographical lines. Body weight should be spread evenly over the largest possible area [317]. Carrying body weight relies on the feet. They are also the location where muscle chains end, whose role, according to the tensegrity theory is to transfer the load to the upper anatomical structures, which causes anomalies in the

correct body posture. Tension of the fascia does not depend on muscle tension. This is due to the existence of autonomous nervous fibres that regulate initial fascial tension with the use of the contractibility of the non-striated muscles [326-328].

Summary

The human body may be termed as a tensegrity structure. This means that a change in the tension of the fascia, tendons and muscles, the structures distant from the head, leads to the TMD, using the transferring mechanism [329]. A proper body posture is characterised when particular segments are placed harmoniously against one another ensuring the flexibility of movements and stability of the support, with the lowest possible energy cost. Every change in the system, of one segment against another, causes changes in the other segments [330]. On the basis of the findings in many publications it was proved that the change in the statics and the function of joints distant to the TMJ is responsible for various dysfunctions of the aforementioned structure. The contrary dependence has also been confirmed. Therapy of particular limb joints or the spine led to a considerable reduction of problems in the TMJ.

Conclusions

Studies on tensegrity provide a lot of information, which changes the way of diagnosing and the treatment of the patient with dysfunctions. Specific structural elements should not be treated and analysed separately, but a holistic approach should be applied. It is worth realizing that disorders of the masticatory system should be treated by an interdisciplinary team, and not by one professional. The interdisciplinary team should include a dentist, physiotherapist, orthopaedist, logopedist, laryngologist and psychologist. Only through a joint and compatible effort of such a team, can the relative benefits be achieved in the context of the improvement in the functioning of the TMJ, as well as the organism as a whole. It should be mentioned that further studies are advisable in order to complement the currently available knowledge on tensegrity.

Adres do korespondencji / Corresponding author

Danuta Lietz-Kijak

E-mail: danuta.lietzkijak@gmail.com

Piśmiennictwo/ References

1. Häkkinen M., Viikari-Juntura E., Martikainen R.: Job experience, work load, and risk of musculoskeletal disorders. *Occup. Environ. Med.* 2001; 58:129–135.
2. Coggon D, Ntani G, Vargas-Prada S, Martinez JM, Serra C, Benavides FG, et al. International variation in absence from work attributed to musculoskeletal illness: findings from the CUPID study. *Occup Environ Med.* 2013; 70(8): 575–84.
3. Westgaard R.H.: Work related musculoskeletal complaints: some ergonomic challenges upon the start of a New century. *Appl. Ergon.* 2000; 31:569–580.
4. Theorell T.: Possible mechanism behind the relationship between the demand-control-support model and disorders of the locomotors system. W: Moon S.D., Sauter S.L. [red.]. *Beyond biomechanism- psychosocial aspects of musculoskeletal disorders in office work.* Taylor and Francis, London 1999, 65–73.
5. Devereux J.J., Vlachonikolis I.G., Buckle P.W.: Epidemiological study to investigate potential interaction between physical and psychosocial factors at work that may increase the risk of symptoms of musculoskeletal disorders of the neck and upper limb. *J. Occup. Environ. Med.* 2002; 59:269–277.
6. Bongers P.M., Ijmker S., van den Heuvel S., Blatter B.M.: Epidemiology of work-related neck and upper limb problems: Psychosocial and personal factors (Part I) and Effective interventions from a bio behavioural perspective (Part II). *J. Occup. Rehabil.* 2006; 16:279–302.
7. De Zwart B.C.H., Frings-Dresen M.H.W., Kilbom A.: Gender differences in upper extremity musculoskeletal complaints in the working population. *Int. Arch. Occup. Environ. Health* 2001; 74:21–30.
8. Roquelaure Y., Ha C., Leclerc A., Touranchet A., Sauteron M., Melchior M. i wsp.: Epidemiologic surveillance of upper-extremity musculoskeletal disorders in the working population. *Arthritis Rheum.* 2006; 55(5):765–778.
9. Holte K.A., Westgaard R.H.: Further studies of shoulder and neck pain and exposures in customer service work with low biomechanical demands. *Ergonomics* 2002; 45(8):887–909.
10. Bongers P.M., Kremer A.M., ter Laak J.: Are psychosocial factors, risks factors for symptoms and signs of the shoulder, elbow, or hand/wrist? A review of the epidemiological literature. *Am. J. Ind. Med.* 2002; 41:315–42.

11. Walker-Bone K., Reading I., Coggon D., Cooper C., Palmer K.T.: Risk factors for specific upper limb disorders as compared with non-specific upper limb pain: Assessing the utility of a structured examination schedule. *Occup. Med. (Lond.)* 2006;56(4):243–250.
12. Miranda H., Viikari-Juntura E., Heistaro S., Heliövaara M., Riihimäki H.: A population study on differences in the determinants of specific shoulder disorder versus nonspecific shoulder pain without clinical findings. *Am. J. Epidemiol.* 2005;161(9):847–855.
13. Bovenzi M., Zadini A., Franzinelli A., Borgogni F.: Occupational musculoskeletal disorders in the neck and upper limbs of forestry workers exposed to hand-arm vibration. *Ergonomics* 1991;34(5):547–562.
14. Patrick N., Emanski E., Knaub MA. Acute and chronic low back pain. *Med Clin North Am.* 2014;98(4):777–789.
15. Balagué F, Mannion AF, Pellisé F, Cedraschi C. Non-specific low back pain. *Lancet.* 2012;379(9814):482–491.
16. Hoy D, Brooks P, Blyth F, Buchbinder R. The epidemiology of low back pain. *Best Pract Res Clin Rheumatol.* 2010;24(6):769–781.
17. Stanton TR, Latimer J, Maher CG, Hancock M. Definitions of recurrence of an episode of low back pain: a systematic review. *Spine* 2009;34: E316–22.
18. Axen I, Leboeuf-Yde C. Trajectories of low back pain. *Best Pract Res Clin Rheumatol.* 2013; 27(5): 601–12.
19. Stanton TR, Henschke N, Maher CG, Refshauge KM, Latimer J, McAuley JH. After an episode of acute low back pain, recurrence is unpredictable and not as common as previously thought. *Spine (Phila Pa 1976).* 2008;33(26):2923–2928.
20. Carey TS, Garrett JM, Jackman A, Hadler N. Recurrence and care seeking after acute back pain: results of a long-term follow-up study. North Carolina Back Pain Project. *Med Care.* 1999;37(2):157–164.
21. Taylor JB, Goode AP, George SZ, Cook CE. Incidence and risk factors for first-time incident low back pain: a systematic review and meta-analysis. *Spine J.* 2014;14: 2299–231.
22. O'Sullivan P, Waller R, Wright A et al. Sensory characteristics of chronic non-specific low back pain: a subgroup investigation. *Man Ther* 2014; 19:311–318.
23. White W.W., Panjabi M.M.: *Clinical Biomechanics of the spine.* Lippincott, Philadelphia 1990; 10–12.
24. Adams M.A., Hutton W.C.: Prolaps intervertebral disc. Hyper flexion injury. *Spine* 1982;7(3):184–191.
25. Sowers L.: Epidemiology of risk factors of osteoarthritis: Systemic factors. *Curr. Opin. Rheumatol.* 2001; 13:447–451.
26. Lawrence R.C., Helmick C.C., Arnett F.C., Deyo R.A., Felson D.T., Giannini E.H. *isp.*: Estimates of the prevalence of arthritis and selected musculoskeletal disorders in the United States. *Arthritis Rheum.* 1998; 41:778–799.
27. Hagen K.B., Magnus P., Vetlesen K.: Neck/shoulder and low-back disorders in the forestry industry: Relationship to work tasks and perceived psychosocial job stress. *Ergonomics* 1998;41(10):1510–1518.
28. Sillanpää J., Huikko S., Nyberg M., Kivi P., Laippala P., Uitti J.: Effect of work with display units on musculoskeletal disorders in the office environment. *Occup. Med.* 2003;53(7):443–451.
29. Lorusso A., Bruno S., L'abbate N.: A review of low back pain and musculoskeletal disorders among Italian nursing personnel. *Ind. Health* 2007; 45:637–644.
30. Holmberg S., Thelin A., Stiemstrom E.-L., Svardsudd K.: The impact of physical work exposure on musculoskeletal symptoms among farmers and rural non-farmers. A population-based study. *Ann. Agric. Environ. Med.* 2003; 10:179–184.
31. Cromie J.E., Robertson V.J., Best M.O.: Work-related musculoskeletal disorders in physical therapists: Prevalence, severity, risks, and responses. *Phys. Ther.* 2000;80(4):336–351.
32. Leggat P.A., Smith D.R.: Musculoskeletal disorders self-reported by dentists in Queensland, Australia. *Aust. Dent. J.* 2006;51(4):324–332.
33. Hayes M, Cockrell D, Smith DR. A systematic review of musculoskeletal disorders among dental professionals. *Int J Dent Hyg.* 2009 Aug;7(3):159-65. doi: 10.1111/j.1601-5037.2009.00395.x.
34. Smith DR, Mihashi M, Adachi Y, Koga H, Ishtake T. A detailed analysis of musculoskeletal disorder risk factors among Japanese nurses. *J Safety Res* 2006; 37: 195–200.
35. Gupta A, Ankola AV, Hebbal M. Dental ergonomics to combat musculoskeletal disorders: a review. *Int J Occup Saf Ergon.* 2013;19(4):561-71.
36. Alyahya F, Algarzaie K, Alsuhay Y, Khounganion R. Awareness of ergonomics & work-related musculoskeletal disorders among dental professionals and students in Riyadh, Saudi Arabia. *J Phys Ther Sci.* 2018 Jun; 30(6): 770–776. doi: 10.1589/jpts.30.770.
37. Finkbeiner BL: Four-handed dentistry: instrument transfer. *J Contemp Dent Pract.* 2001, 2: 57–76.
38. Sarkar PA, Shigji AL: Ergonomics in general dental practice. *People's J Sci Res.* 2012, 5: 56–60.
39. Morse T, Bruneau H, Dussetschleger J.: Musculoskeletal disorders of the neck and shoulder in the dental professions. *Work.* 2010, 35, 4, 419-429.
40. Diaz-Caballero A.J., Gomez-Palencia I.P., Diaz-Cardenas S.: Ergonomic factors that causa the presence of pain muscle in students of dentistry. *Medicina Oral, Patologia Oral Y Cirurgia Bucal.* 2010, 15, 6: 906-911.
41. Kierklo A., Kobus A., Jaworska M., Botuliński B.: Work-related musculoskeletal disorders among dentists – a questionnaire survey. *Annals of Agricultural and Environmental medicine.* 2011, 18, 1: 79-84.
42. Valachi B, Valachi K. Preventing musculoskeletal disorders in clinical dentistry: strategies to address the mechanisms leading to musculoskeletal disorders. *J Am Dent Assoc.* 2003; 134:1604-12.
43. Lindfors P, von Thiele U, Lundberg U. Work characteristics and upper extremity disorders in female dental health workers. *J Occup Health.* 2006; 48:192-7.
44. Harutunian K, Gargallo-Albiol J, Figueiredo R, Gay-Escoda C. Ergonomics and musculoskeletal pain among postgraduate students and faculty members of the School of Dentistry of the University of Barcelona (Spain). A cross-sectional study. *Med Oral Patol Oral Cir Bucal.* 2011 May 1;16(3): e425-9.
45. Hodacova L, Sustova Z, Cermakova E, Kapitan M, Smejkalova J. Self-reported risk factors related to the most frequent musculoskeletal complaints among Czech dentists. *Ind Health.* 2015;53(1):48-55. doi: 10.2486/inhealth.2013-0141.
46. Shrestha BP, Singh GK, Niraula SR. Work related complaints among dentists. *JNMA J Nepal Med Assoc.* 2008 Apr-Jun;47(170):77-81.
47. Gupta A, Bhat M, Mohammed T, Bansal N, Gupta G. Ergonomics in dentistry. *Int J Clin Pediatr Dent.* 2014 Jan-Apr; 7(1): 30–34. doi: 10.5005/jp-journals-10005-1229.
48. Youssef JA, Orndorff DO, Patty CA, et al. Current status of adult spinal deformity. *Glob Spine J* 2013; 3: 51–62.
49. Ailon T, Smith JS, Shaffrey CI, et al. Degenerative spinal deformity. *Neurosurgery* 2015; 77 (suppl 4): S75–91.
50. Bess S, Boachie-Adjei O, Burton D, et al. Pain and disability determine treatment modality for older patients with adult scoliosis, while deformity guides treatment for younger patients. *Spine (Phila Pa 1976)* 2009; 34: 2186–90.
51. Ahrens W, Pigeot I, Pohlmann H, De Henauw S, Lissner L, Molnár D, et al. Prevalence of overweight and obesity in European children below the age of 10. *Int J Obes.* 2014;38: S99–107. <https://doi.org/10.1038/ijo.2014.140>.
52. Bibiloni MDM, Pons A, Tur JA. Prevalence of overweight and obesity in adolescents: a systematic review. *ISRN Obes.* 2013; 2013:392747. <https://doi.org/10.1155/2013/392747>.
53. Lafage R, Schwab F, Glassman S, et al. Age-adjusted alignment goals have the potential to reduce PJK. *Spine (Phila Pa 1976)* 2017; 42: 1275–82.
54. Diebo BG, Oren JH, Challier V, et al. Global sagittal axis: a step toward full-body assessment of sagittal plane deformity in the human body. *J Neurosurg Spine* 2016; 25: 494–99.
55. Lafage R, Schwab F, Challier V, et al. Defining spino-pelvic alignment thresholds: should operative goals in adult spinal deformity surgery account for age? *Spine (Phila Pa 1976)* 2016; 41: 62–68.
56. Barker V. *Postura, Posizione e Movimento.* Ed. Mediterranee, 1998.
57. Gori L, Firenzuoli F. Posturology. Methodological problems and scientific evidence. *Recenti Prog Med* 2005; 96(2): 89-91.
58. Shiba Y, Taneichi H, Inami S, Moridaira H, Takeuchi D, Nohara Y. Dynamic global sagittal alignment evaluated by three-dimensional gait analysis in patients with degenerative lumbar kyphoscoliosis. *Eur Spine J* 2016; 25: 2572–79.
59. Schwab F, Lafage V, Farcy JP, et al. Surgical rates and operative outcome analysis in thoracolumbar and lumbar major adult scoliosis. *Spine (Phila Pa 1976)* 2007; 32: 2723–30.
60. Yagi M, Kaneko S, Yato Y, Asazuma T, Machida M. Walking sagittal balance correction by pedicle subtraction osteotomy in adults with fixed sagittal imbalance. *Eur Spine J* 2016; 25: 2488–96.
61. Smith JS, Shaffrey CI, Fu K-MG, et al. Clinical and radiographic evaluation of the adult spinal deformity patient. *Neurosurg Clin N Am* 2013; 24: 143–56.
62. Kapo S, Rado I, Smajlović N, Kovač S, Talović M, Doder I, et al. Increasing postural deformity trends and body mass index analysis in school-age children. *Zdr Varst.* 2018; 57:25–32. <https://doi.org/10.2478/sjph-2018-0004>.
63. Salenius P, Vanikka E. The development of the tibiofemoral angle in children. *J Bone Joint Surg Am.* 1975; 57:259–61 <http://www.ncbi.nlm.nih.gov/pubmed/1112851>. Accessed 26 Jul 2017.
64. Janssen I, LeBlanc AG, Kannus P, Rimpela A, Legg C, Lumb A, et al. Systematic review of the health benefits of physical activity and fitness in school-aged children and youth. *Int J Behav Nutr Phys Act.* 2010; 7:40. <https://doi.org/10.1186/1479-5868-7-40>.
65. Scoppa F. Posturology: from nonlinear dynamics to transdisciplinarity. *Otoneurologia* 2000; 15: 28-48.
66. Oravičan M. Posturology-fundamental concepts and practical applications. *Analele UVV- Seria EFS* 2009; 11: 61-69.
67. Scoppa F. Posturology: the neurophysiological model, the biomechanical model, the model psychosomatic. *Otoneurologia* 2002; 9: 3-13.
68. Ambrosi F. *Fondamenti di Posturologia.* Ed. 2012.
69. Baldini A. Clinical and instrumental treatment of a patient with dysfunction of the stomatognathic system: a case report. *Ann Stomatol (Roma)* 2010; 1(2): 2-5.
70. Manfredini D, Segù M, Brady Buccì M, Castorfoli T, Di Giosia M, Peretta R, et al. Occlusione, postura e disordini temporomandibolari. Approccio evidence-based alla pratica clinica. *Tagete Archives of Legal Medicine and Dentistry* 2011; 2: 298-301.
71. Ciancaglini R, Gelmetti R, Lazzari E. Evoluzione degli studi sulla relazione tra occlusione e postura. *Mondo Ortodontico* 1/2008, pp. 59-65.
72. Ciancaglini R, Cerni C, Saggini R, Bellomo RG, Ridi R, Pisciella V, et al. On the Symposium: Consensus Conference Posture and Occlusion: Hypothesis of Correlation. *J. Stomat Occ Med* 2009: 87-96.
73. Ober WC, Garrison CW, Silverthorn AC. *Fisiologia, un approccio integrato.* Casa Editrice Ambrosiana 2000.
74. Ciancaglini R, Cerni C, Saggini R, Bellomo RG, Ridi R, Pisciella V, et al. On the Symposium: Consensus Conference Posture and Occlusion: Hypothesis of Correlation. *J. Stomat Occ Med* 2009: 87-96.
75. Ailon T, Smith JS, Shaffrey CI, et al. Degenerative spinal deformity. *Neurosurgery* 2015; 77 (suppl 4): S75–91.
76. Russo A, Bransford R, Wagner T, Lee MJ, Chapman JR. Adult degenerative scoliosis insights, challenges, and treatment outlook. *Curr Orthop Pract* 2008; 19: 357–65.
77. Weinstein SL, Dolan LA, Cheng JCY, Danielsson A, Morcuende JA. Adolescent idiopathic scoliosis. *Lancet* 2008; 371: 1527–37.

78. Silva FE, Lenke LG. Adult degenerative scoliosis: evaluation and management. *Neurosurg Focus* 2010; 28: E1.
79. Liu S, Diebo BG, Henry JK, et al. The benefit of nonoperative treatment for adult spinal deformity: identifying predictors for reaching a minimal clinically important difference. *Spine J* 2016; 16: 210–18.
80. DePalma MJ, Slipman CW. Evidence-informed management of chronic low back pain with epidural steroid injections. *Spine J* 2008; 8: 45–55.
81. Fejer R, Kyvik KO, Hartvigsen J. The prevalence of neck pain in the world population: A systematic critical review of the literature. *Eur Spine J*. 2006;15(6):834–48, <http://dx.doi.org/10.1007/s00586-004-0864-4>.
82. Côté P, Cassidy JD, Carroll L. The factors associated with neck pain and its related disability in the Saskatchewan population. *Spine*. 2000;25(9):1109–17.
83. Chiu TTW, Leung ASL. Neck pain in Hong Kong: A telephone survey on prevalence, consequences, and risk groups. *Spine*. 2006;31(16): E540–4, <http://dx.doi.org/10.1097/01.brs.0000225999.02326.ad>.
84. Davatchi F. Rheumatic diseases in the APLAR region. *J Rheum*. 2006; 9:5–10, <http://dx.doi.org/10.1111/j.1479-8077.2006.00177.x>.
85. Davatchi F, Jamshidi AR, Banihashemi AT, Gholami J, Forouzanfar MH, Akhlaghi M, et al. WHO-ILAR COPCORD study (stage 1, urban study) in Iran. *J Rheumatol*. 2008;35(7):1384–90.
86. Aarabi A, Zamiri B, Mohammadinezhad C, Rahmani F, Mahmoudi H. Musculoskeletal disorders in dentists in Shiraz, southern Iran. *Iranian Red Crescent Med J*. 2009; 11(4):464–5.
87. Chamani G, Zarei MR, Momenzadeh A, Safizadeh H, Rad M, Alahyari A. Prevalence of musculoskeletal disorders among dentists in Kerman, Iran. *J Musculoskelet Pain*. 2012;20(3):202–7.
88. Kang JH, Park RY, Lee SJ, Kim JY, Yoon SR, Jung KI. The effect of the forward head posture on postural balance in long time computer based worker. *Ann Rehabil Med*. 2012;36(1):98–104, <http://dx.doi.org/10.5535/arm.2012.36.1.98>.
89. Yp CHT, Chiu TTW, Poon ATK. The relationship between head posture and severity and disability of patients with neck pain. *Man Ther*. 2008;13(2):148–54, <http://dx.doi.org/10.1016/j.math.2006.11.002>.
90. Lau KT, Cheung KY, Chan MH, Chan KB, Lo KY, Chiu TT. Relationships between sagittal postures of thoracic and cervical spine, presence of neck pain, neck pain severity and disability. *Man Ther*. 2010;15(5):457–62, <http://dx.doi.org/10.1016/j.math.2010.03.009>.
91. Schwab FJ, Blondel B, Bess S, et al. Radiographical spinopelvic parameters and disability in the setting of adult spinal deformity: a prospective multicenter analysis. *Spine (Phila Pa 1976)* 2013; 38: e803–12.
92. Diebo BG, Ferrero E, Lafage R, et al. Recruitment of compensatory mechanisms in sagittal spinal malalignment is age and regional deformity dependent: a full-standing axis analysis of key radiographical parameters. *Spine (Phila Pa 1976)* 2015; 40: 642–49.
93. Day LM, DeSole EM, Beaubrun BM, et al. Radiological severity of hip osteoarthritis in patients with adult spinal deformity: the effect on spinopelvic and lower extremity compensatory mechanisms. *Eur Spine J* 2018; 27: 2294–302.
94. Davatchi F, Banihashemi AT, Gholami J, Faezi ST, Forouzanfar MH, Salehi M, et al. The prevalence of musculoskeletal complaints in a rural area in Iran: A WHO-ILAR COPCORD study (stage 1, rural study) in Iran. *Clin Rheumatol*. 2009;28(11):1267–74, <http://dx.doi.org/10.1007/s10067009-1234-8>.
95. Pargali N, Jowkar N. Prevalence of musculoskeletal pain among dentists in Shiraz, Southern Iran. *Int J Occup Environ Med*. 2010;1(2):69–7.
96. Glassman SD, Bridwell K, Dimar JR, Horton W, Berven S, Schwab F (2005) The impact of positive sagittal balance in adult spinal deformity. *Spine* 30:2024–2029.
97. Schwab F, Patel A, Ungar B, Farcy JP, Lafage V (2010) Adult spinal deformity-postoperative standing imbalance: how much can you tolerate? An overview of key parameters in assessing alignment and planning corrective surgery. *Spine* 35:2224–2231.
98. Martini, Timmison, Talltsch; Anatomia umana. EdiSES, 2003.
99. Wirhed R. Anatomia del movimento e abilità. Edi. Ermes, 2002.
100. Kapandji IA. Fisiologia articolare, volume III, Monduzzi Editore, 2004.
101. Autori vari, Fisiologia dell'uomo, Edi. Ermes, 2002.
102. J.-H. Park, S. Noh, H. Jang, W. Yu, M. Park, H. Choi, The study of correlation between foot-pressure distribution and scoliosis 13th International Conference on Biomedical Engineering, Springer (2009), pp. 974-978.
103. J.-U. Lee, M.-Y. Kim, J. Kim Comparison of static plantar foot pressure between healthy subjects and patients with adolescent idiopathic scoliosis *Toxicol. Environ. Health Sci.*, 6 (2014), pp. 127-132.
104. P. Szulc, P. Bartkowiak, J. Lewandowski, J. Markuszewski, The influence of idiopathic scoliosis on load distribution in the foot, *Chirurgia narządów ruchu i ortopedia polska*, 73 (2008), pp. 187-191.
105. M. Yagi, H. Ohne, T. Konomi, K. Fujiyoshi, S. Kaneko, M. Takemitsu, et al., Walking balance and compensatory gait mechanisms in surgically treated patients with adult spinal deformity *Spine J.*, 17 (2017), pp. 409-417.
106. G.C. Gauchard, P. Lascombes, M. Kuhnast, P.P. Perrin Influence of different types of progressive idiopathic scoliosis on static and dynamic postural control *Spine*, 26 (2001), pp. 1052-1058.
107. T. Haumont, G.C. Gauchard, P. Lascombes, P.P. Perrin Postural instability in early-stage idiopathic scoliosis in adolescent girls *Spine*, 36 (2011), pp. E847-E854.
108. Ma Q, Lin H, Wang L, Zhao L, Chen M, Wang S, Rao Z, Luo Y, Correlation between spinal coronal balance and static baropodometry in children with adolescent idiopathic scoliosis, *Gait Posture*. 2020 Jan; 75:93-97. doi: 10.1016/j.gaitpost.2019.10.003. Epub 2019 Oct 9.
109. Klineberg E, Mazanec D, Orr D, Demicco R. Masquerade: medical causes of back pain. *Cleve Clin J Med*. 2007; 74:905–13. doi: 10.3949/ccjm.74.12.905.
110. Fallentin N.: Regulatory actions to prevent work-related musculoskeletal disorders — The use of research based exposure limits. *Scand. J. Work Environ. Health* 2003;29(4):247–250.
111. Punnett L., Wegman D.H.: Work-related musculoskeletal disorders: The epidemiological evidence and debate. *J. Electromyogr. Kinesiol*. 2004; 14:13–23.
112. Hoogedoom W., Poppel M. van, Bongers P., Koes B., Bouter L.: Systematic review of psychosocial factors at work, private life as risk factor for back pain. *Spine* 2000; 25:2114–2125.
113. Barty S., Burton K., Main C.: A prospective study of psychosocial factors and absence due to musculoskeletal disorders — implications for occupational screening. *Occup. Med*. 2005; 55:375–379.
114. Camacho-Soto A, Sowa GA, Perera S, Weiner DK. Fear avoidance beliefs predict disability in older adults with chronic low back pain. *PM R*. 2012; 4:493–7.
115. Basler H-D, Luckmann J, Wolf U, Quint S. Fear-avoidance beliefs, physical activity, and disability in elderly individuals with chronic low back pain and healthy controls. *Clin J Pain*. 2008, 24(7):604-10. doi: 10.1097/AJP.0b013e31816b54f6
116. Cherkin DC, Sherman KJ, Balderson BH, et al. Effect of mindfulness-based stress reduction vs cognitive behavioral therapy or usual care on back pain and functional limitations in adults with chronic low back pain: a randomized clinical trial. *JAMA*. 2016;315(12):1240–1249.
117. Kamper SJ, Apeldoorn AT, Chiarotto A, Smeets RJ, Ostelo RW, Guzman J, et al. Multidisciplinary biopsychosocial rehabilitation for chronic low back pain. *Cochrane Database Syst Rev* 2014;9 CD000963.
118. Massey E.W.: Carpal tunnel syndrome in pregnancy. *Obstet. Gynecol. Surv*. 1978; 33:145–148.
119. Lam N., Thurstone A.: Association of obesity, tender, age and occupation with carpal tunnel syndrome. *Aust. N.Z.J. Surg*. 1998; 68:190–193.
120. De Zwart B.C.H., Frings-Dresen M.H.W., Kilbom A.: Gender differences in upper extremity musculoskeletal complaints in the working population. *Int. Arch. Occup. Environ. Health* 2001; 74:21–30.
121. National Fibromyalgia Research Association: ACR Fibromyalgia Diagnostic Criteria [2010]. Address: <http://www.nfra.net/Diagnost.htm>
122. Linton SJ, Boersma K, Jansson M, Svärd L, Botvalde M. The effects of cognitive-behavioral and physical therapy preventive interventions on pain-related sick leave: a randomized controlled trial. *Clin J Pain* 2005; 21:109-19.
123. Thomas AW, Graham K, Prato FS, McKay J, Forster PM, Moulin DE, Chari SA randomized, double-blind, placebo-controlled clinical trial using a low-frequency magnetic field in the treatment of musculoskeletal chronic pain. *Pain Res. Manag*. 2007;12(4):249–58.
124. Aréja AS, Kotowich A, Staley D, Summers R, Tappia PS. Electromagnetic fields in the treatment of chronic lower back pain in patients with degenerative disc disease. *Future Science OA*. 2016; 2(1): FSO105.
125. Pall ML. Electromagnetic fields act via activation of voltage-gated calcium channels to produce beneficial or adverse effects. *Journal Cell Mol. Med*. 2013; 17(8): 958–965.
126. Iannitti T, Fisetto G, Esposito A, Rottigni V, Palmieri B. Pulsed electromagnetic field therapy for management of osteoarthritis-related pain, stiffness and physical function: clinical experience in the elderly. *Clin. Interv. Aging*. 2013; 8:1289–93.
127. Nelson FR, Zvirbulis R, Pilla AA. Non-invasive electromagnetic field therapy produces rapid and substantial pain reduction in early knee osteoarthritis: a randomized double-blind pilot study. *Rheumatol. Int*. 2013; 33(8):2169-73.
128. Calderhead RG, Won-Serk K, Ohshiro T, Trelles MA, Vasily DB. Adjunctive 830 nm light-emitting diode therapy can improve the results following aesthetic procedures. *Laser Ther*. 2015; 24(4):277–89.
129. Vaughan B, Morrison T, Buttigieg D, Macfarlane C, Fryer G: Approach to low back pain - osteopathy. *Aust Fam Physician*. 2014, 43: 197-19.
130. Franke H, Fryer G, Ostelo R, Kamper S. Muscle energy technique for non-specific low-back pain. A Cochrane systematic review. *International Journal of Osteopathic Medicine*, 2016, Volume 20, 41 – 52.
131. Wilson E, Payton O, Donegan Shoaf L, Dec K. Muscle energy technique in patients with acute low back pain: a pilot clinical trial. *The Journal of Orthopaedic and Sports Physical Therapy* 2003;33(9):502–12.
132. Patil P, Chandu B, Metgud S, Khatri S. Effectiveness of Muscle Energy Technique on quadratuslumborum in acute low back pain randomized controlled trial. *Indian Journal of Physiotherapy and Occupational Therapy* 2010;4(1):54–8.
133. Selkow N, Grindstaff T, Cross K, Pugh K, Hertel J, Saliba S. Short term effect of Muscle Energy Technique on pain in individuals with non-specific lumbopelvic pain: A pilot study. *The Journal of Manual & Manipulative Therapy* 2009;17(1): E14–8.
134. Bindra S, Kumar M, Singh P, Singh J. A study on the efficacy of Muscle Energy Technique as compared to conventional therapy in chronic low back pain due to sacroiliac joint dysfunction. *Indian Journal of Physiotherapy and Occupational Therapy* 2012;6(1):200–3.
135. Dhinkaran M, Sareen A, Arora T. Comparative analysis of Muscle Energy Technique and conventional physiotherapy in treatment of sacroiliac joint dysfunction. *Indian Journal of Physiotherapy and Occupational Therapy* 2011;5(4):127–30.
136. Assendelft WJ, Morton SC, Yu EI, Suttrop MJ, Shekelle PG. Spinal manipulative therapy for low back pain. A meta-analysis of effectiveness relative to other therapies. *Ann Intern Med*. 2003; 138:871–81.
137. Pasquier M, Daneau C, Marchand AA, Lardon A, Descarreaux M. Spinal manipulation frequency and dosage effects on clinical and physiological outcomes: a scoping review. *Chiropr Man Therap*. 2019 May 22; 27:23. doi: 10.1186/s12998-019-0244-0.
138. Furlan AD, Yazdi F, Tsertsvadze A, et al. A systematic review and meta-analysis of efficacy, cost-effectiveness, and safety of selected complementary and alternative medicine for neck and low-back pain. *Evid Based Complement Alternat Med*. 2012;2012: 953139.
139. Bronfort G, Haas M, Evans RL, Bouter LM. Efficacy of spinal manipulation and mobilization for low back pain and neck pain: a systematic review and best evidence synthesis. *Spine J*. 2004; 4:335–56.

140. Moutzouri M, Billis E, Strimpakos N, Kottika P, Oldham JA. The effects of the Mulligan Sustained Natural Apophyseal Glide (SNAG) mobilisation in the lumbar flexion range of asymptomatic subjects as measured by the Zebris CMS20 3-D motion analysis system. *BMC Musculoskeletal Disorders*. 2008; 9:131.
141. Waqqar S., Shakil-ur-Rehman S., Ahmad S. McKenzie treatment versus mulligan sustained natural apophyseal glides for chronic mechanical low back pain. *Pak J Med Sci*. 2016 Mar-Apr; 32(2): 476–479.
142. Hisham Mohamed Hussien, Neveen Abdellatif Abdel-Raouf, Ormaima Mohamed Kattabei, Hassan Hussien Ahmed. Effect of Mulligan Concept Lumbar SNAG on Chronic Nonspecific Low Back Pain. *J Chiropr Med*. 2017 Jun; 16(2): 94–102.
143. May J., Krzyzanowicz R., Nasypany A., Baker R. Seegmiller J. Mulligan Concept Use and Clinical Profile From the Perspective of American Certified Mulligan Practitioners. *Journal of Sport Rehabilitation*, Nov. 2015, 24 (4): 337 – 341.
144. Dunsford A, Kumar S, Clarke S. Integrating evidence into practice: use of McKenzie-based treatment for mechanical low back pain. *J Multidiscip Healthc*. 2011; 4:393–40.
145. Delaney PM, Hubka MJ. The diagnostics utility of McKenzie clinical assessment for Lower bak pain. *J Manip & Physio Therapeutics* 1999; 22; 628-630.
146. Cherkin DC, Deyo RA, Battla MC, Street JH, Hund M, Barlow W. A comparison of Physical therapy chiropractice manipulation or an educational booklet for the treatment of low back pain. *New Eng J Med*. 1998;339(15):1021–1029.
147. Machado LA, de Souza MS, Ferreira PH, Ferreira ML. The McKenzie method for low back pain: a systematic review of the literature with a meta-analysis approach. *Spine*. 2006; 31:254–262.
148. Kase K, Wallis J., Kase T. *Clinical Thetapeutic Applications of the Kinesio Taping Method*. 2nd editio. Ken Ikai Co. Ltd., 2003.
149. Neumann, Donald A. *Kinesiology of the Musculoskeletal System: Foundations for Physical Rehabilitation*. Mosby, 2002.
150. Kase K, Hashimoto T, Tomoki O. *Development of kinesio taping perfect manual*. Kinesio Taping Association (1996).
151. Kase K, Wallis J, Kase T. *Clinical therapeutic applications of the kinesio taping method*. Tokyo: Ken Ikai Co., Ltd; 2003.
152. Williams S, Whatman C, Hume PA, Sheerin K. Kinesio taping in treatment and prevention of sports injuries: a meta-analysis of the evidence for its effectiveness. *Sports Med*. 2012; 42:153–164.
153. Breitenbach S. Schwangerschaft und Kinesio-Tape. *Phys Ther* 2005; 5: 724-727.
154. Sayed A, Tantawy, Dalia M, Kamel. The effect of kinesio taping with exercise compared with exercise alone on pain, range of motion, and disability of the shoulder in postmastectomy females: a randomized control trial. *J Phys Ther Sci*. 2016; 28(12): 3300–3305.
155. Chen S. M, Alexander R, Kai Lo S, Cook J. Effects of Functional Fascial Taping on pain and function in patients with non-specific low back pain: a pilto randomized controlled trial. *Clin Rehabil*. 2012; 26: 924-933.
156. Added MA, Costa LO, Fukuda TY, Freitas D, Salomao E, Monteiro E, Costa L. Efficacy of adding the kinesio taping method to guideline-endorsed convetional physiotherapy in patients with chronic nonspecific low back pain: a randomised controlled trial. *BMC Musculoskeletal Disorders*. 2013; 14: 301.
157. Paoloni M, Bemetti A, Fratocchi G, Mangone M, Parrinello L, Cooper M, et al. Kinesio Taping applied to lumbar muscles influences clinical and electromyographic characteristics in chronic low back pain patients. *Eur J Phys Rehab Med*. 2011; 2 (47): 237-244.
158. Ay S, Konak HE, Evcik D, Kibar S. The effectiveness of Kinesio Taping on pain and disability in cervical myofascial pain syndrome. *Rev Bras Reumatol Engl Ed*. 2017, 57(2):93-99. doi: 10.1016/j.rbr.2016.03.012.
159. Lee JH. The Kinesio Taping technique may affect therapeutic results. *J Physiother*. 2015 Oct;61(4):231. doi: 10.1016/j.jphys.2015.03.006.
160. Fu TC, Wong AM, Pei YC, Chou SW, Lin YC. Effect of Kinesio taping on muscle strength in athletes-a pilot study. *J Sci Med Sport*. 2008; 11:198–201. doi: 10.1016/j.jsams.2007.02.011.
161. Selva F, Pardo A, Aguado X, Montava I, Gil-Santos L, Barrios C. A study of reproducibility of kinesiology tape applications: review, reliability and validity. *BMC Musculoskelet Disord*. 2019 Apr 9;20(1):153. doi: 10.1186/s12891-019-2533-0.
162. Castro-Sánchez AM, Lara-Palomo IC, Matarán-Peñarocha GA, Fernández-Sánchez M, Sánchez-Labraca N, Arroyo-Morales M. Kinesio Taping reduces disability and pain slightly in chronic non-specific low back pain: a randomised trial. *J Physiother*. 2012;58(2):89-95. doi: 10.1016/S1836-9553(12)70088-7.
163. Saavedra-Hernández M, Castro-Sánchez AM, Arroyo-Morales M, Cleland JA, Lara-Palomo IC, Fernández-de-Las-Perías C. Short-term effects of kinesio taping versus cervical thrust manipulation in patients with mechanical neck pain: a randomized clinical trial. *J Orthop Sports Phys Ther*. 2012 Aug;42(8):724-30. doi: 10.2519/jospt.2012.4086.
164. Abbasi S, Rohhani-Shirazi Z, Shokri E, García-Muro San José F. The effect of Kinesio Taping on postural control in subjects with non-specific chronic low back pain. *J Bodyw Mov Ther*. 2018 Apr;22(2):487-492. doi: 10.1016/j.jbmt.2017.06.003. Epub 2017 Jun 13.
165. Toprak Celenay S, Ozer Kaya D. Immediate effects of kinesio taping on pain and postural stability in patients with chronic low back pain. *J Bodyw Mov Ther*. 2019 Jan;23(1):206-210.
166. Bernardelli RS, Scheeren EM, Fuentes Filho AR, Pereira PA, Gariba MA, Moser ADL, Bichinho GL. Effects of Kinesio Taping on postural balance in patients with low back pain, a randomized controlled trial. *J Bodyw Mov Ther*. 2019 Jul;23(3):508-514.
167. Shacklock M. Neural mobilization: a systematic review of randomized controlled trials with an analysis of therapeutic efficacy. *J Man Manip Ther*. 2008;16(1):23–24.
168. Scrimshaw S.V., Maher C.G. Randomized controlled trial of neural mobilization after spinal surgery. *Spine (Phila Pa 1976)* 2001;26(24):2647–2652.
169. Olesen T. Auriculotherapy stimulation for neuro-rehabilitation. *NeuroRehabilitation*, 2002, 17:49-62.
170. Vickers AJ, Vertosick EA, Lewith G, MacPherson H, Foster NE, Sherman KJ, Irnich D, Witt CM, Linde K. Acupuncture Trialists C. Acupuncture for chronic pain: update of an individual patient data meta-analysis. *J Pain*. 2018;19(5):455–474. doi: 10.1016/j.jpain.2017.11.00.
171. Ezzo J, Berman B, Hadhazy VA, et al: Is acupuncture effective for the treatment of chronic pain? A systematic review; *Pain*, 2000,86:217-225.
172. Yuan J, Purepong N, Kerr DP, Park J, Bradbury I, McDonough S. Effectiveness of acupuncture for low back pain: a systematic review. *Spine J*. 2008;33(23): E887–E900. doi: 10.1097/BRS.0b013e318186b276.
173. Haake M, Müller HH, Schadebrittinger C, Basler HD, Schäfer H, Maier C, et al. German acupuncture trials (gerac) for chronic low back pain: randomized, multicenter, blinded, parallel-group trial with 3 groups. *Arch Intern Med*. 2007;167(17):1892–1898. doi: 10.1001/Archinte.167.17.1892.
174. Leibing E, Leonhardt U, Köster G, Goerlitz A, Rosenfeldt JA, Hilgers R, Ramadori G. Acupuncture treatment of chronic low-back pain – a randomized, blinded, placebo-controlled trial with 9-month follow-up. *Pain*. 2002;96(1–2):189–196. doi: 10.1016/S0304-3959(01)00444-4.
175. Tulder Mv, Cherkin DC, Berman B, et al: Acupuncture for low back pain. *Cochrane Database Syst Rev* 2:CD001351, 2000.
176. Lee I.S., Lee S.H., Kim S.Y., Lee H.J., Park H.J., Chae Y.Y. Visualization of the meridian system based on biomedical information about acupuncture treatment. *Evid. Based Complement. Alternat. Med*. 2013;2013: 872142.
177. Shin J.Y., Ku B., Kim J.U., Lee Y.J., Kang J.H., Heo H., Choi H.-J., Lee J.-H. Short-term effect of laser acupuncture on lower back pain: A randomized, placebo-controlled, double-blind trial. *Evid. Based Complement. Alternat. Med*. 2016
178. Tiaw-Kee Lim, Yan Ma, Frederic Berger, and Gerhard Litscher, Acupuncture and Neural Mechanism in the Management of Low Back Pain—An Update, *Medicines (Basel)*. 2018 Sep; 5(3): 6.
179. Chia KL. Electroacupuncture treatment of acute low back pain: unlikely to be a placebo response. *Acupunct Med*. 2014;32(4):354–355. doi: 10.1136/acupmed-2014-010582.
180. Kim TH, Kang JW, Lee MS. What is lost in the acupuncture trial when using a sham intervention? *Acupunct Med*. 2017;35(5):384–386. doi: 10.1136/acupmed-2016-011333.
181. Cramer H, Lauche R, Haller H, Dobos G. A systematic review and meta-analysis of yoga for low back pain. *Clin J Pain*. 2013;29(5):450–460.
182. Nambi GS, Inbasekaran D, Khuman R, Devi S, Shanmuganathan, Jagannathan K. Changes in pain intensity and health related quality of life with Iyengar yoga in nonspecific chronic low back pain: a randomized controlled study. *Int J Yoga* 2014; 7:48-53.
183. Bussing A, Ostermann T, Ludtke R, Michalsen A. Effects of yoga interventions on pain and pain associated disability: a meta analysis. *Journal of Pain* 2012;13(1):1–9.
184. Bahçecioglu Turan G. The effect of yoga on respiratory functions, symptom control and life quality of asthma patients: A randomized controlled study. *Complement Ther Clin Pract*. 2019, Oct 28:101070.
185. Cramer H, Ward L, Steel A, Lauche R, Dobos G, Zhang Y. Prevalence, patterns, and predictors of yoga use: results of a U.S. nationally representative survey. *American Journal of Preventive Medicine* 2016;50(2):230–5.
186. Jackson J. K., Shepherd T. R., Kell R. T. The influence of periodized resistance training on recreationally active males with chronic nonspecific low back pain. *Journal of Strength and Conditioning Research*. 2011;25(1):242–251. doi: 10.1519/JSC.0b013e3181b2c83d.
187. Noomohammadpour P, Kordi M, Mansoumia M, Akbari-Fakhrabadi M, Kordi R. Exercise Program in the Treatment of Nurses with Chronic Low Back Pain: A Single-Blinded Randomized Controlled Trial. *Asian Spine J*. 2018 Jun; 12(3): 490–502. Published online 2018 Jun 4. doi: 10.4184/asj.2018.12.3.490.
188. Shamsi M., Sarrafzadeh J., Jamshidi A., Arjmand N., Arjmand F. Comparison of spinal stability following motor control and general exercises in nonspecific chronic low back pain patients. *Clinical Biomechanics*. 2017; 48:42–48. doi: 10.1016/j.clinbiomech.2017.07.006.
189. Hides J. A., Stanton W. R., McMahon S., Sims K, Richardson C. Effect of stabilization training on multifidus muscle cross-sectional area among young elite cricketers with low back pain. *Journal of Orthopaedic & Sports Physical Therapy*. 2008;38(3):101–108. doi: 10.2519/jospt.2008.2658.
190. Shamsi M., Sarrafzadeh J., Jamshidi A., Zarabi V., Pourahmadi M. R. The effect of core stability and general exercise on abdominal muscle thickness in non-specific chronic low back pain using ultrasound imaging. *Physiotherapy Theory and Practice*. 2016;32(4):277–283. doi: 10.3109/09593985.2016.1138559.
191. Waseem Akhtar M, Karimi H, Amir Gilani S. Effectiveness of core stabilization exercises and routine exercise therapy in management of pain in chronic non-specific low back pain: A randomized controlled clinical trial. *Pak J Med Sci*. 2017 Jul-Aug; 33(4): 1002–1006. doi: 10.12669/pjms.334.12664.
192. Roelofs PD, Deyo RA, Koes BW, Scholten RJ, van Tul-der MW. Non-steroidal anti-inflammatory drugs for low back pain. *Cochrane Database Syst Rev*. 2008;(1):CD000396.
193. Hancock MJ, Maher CG, Latimer J, et al. Assessment of diclofenac or spinal manipulative therapy, or both, in addition to recommended first-line treatment for acute low back pain: a randomised controlled trial. *Lancet*. 2007;370(9599):1638–1643.

194. Hancock MJ, Maher CG, Latimer J, McLachlan AJ, Day RO, Davies RA. Can predictors of response to NSAIDs be identified in patients with acute low back pain? *Clin J Pain*. 2009;25(8):659–665.
195. Van Tulder MW, Touray T, Furlan AD, Solway S, Bouter LM. Muscle relaxants for non-specific low back pain. *Cochrane Database Syst Rev*. 2003;(2):CD004252.
196. Kellgren JH. On the distribution of pain arising from deep somatic structures with charts of segmental pain areas. *Clin Sci* 1939;4: 35-46.
197. Schilder A, Hoheisel U, Magerl W, et al. Sensory findings after stimulation of the thoracolumbar fascia with hypertonic saline suggest its contribution to low back pain. *Pain* 2014;155: 222-231.
198. Panjabi M. M. A hypothesis of chronic back pain: ligament subfailure injuries lead to muscle control dysfunction. *European Spine Journal*. 2006;15(5):668–676. doi: 10.1007/s00586-005-0925-3.
199. Schleip R., Vleeming A., Lehmann-Horn F., Klingler W. Letter to the editor concerning 'A hypothesis of chronic back pain: ligament subfailure injuries lead to muscle control dysfunction' (M. Panjabi) *European Spine Journal*. 2007;16(10):1733–173.
200. Langevin H. M., Fox J. R., Koptiuch C., et al. Reduced thoracolumbar fascia shear strain in human chronic low back pain. *BMC Musculoskeletal Disorders*. 2011,12,201,
201. Masi AT, Hannon JC: Human resting muscle tone (HRMT): narrative introduction and modern concepts; *J Bodyw Mov Ther* 2008;12(4):320-32.
202. Wilke J, Krause F, Vogt L, Banzer W. What is evidence-based about myofascial chains: a systematic review. *Archives of physical medicine and rehabilitation*; *Arch Phys Med Rehabil* 2016;97(3):454-61.
203. Pardehshenas H, Maroufi N, Sanjari M, Pamiampour M, Levin SM: Lumbopelvic muscle activation patterns in three stances under graded loading conditions: Proposing a tensegrity model for load transfer through the sacroiliac joints; *J Bodyw Mov Ther* 2014;18(4):633-42.
204. Kassolik K, Jaskólska A, Kisiel-Sajewicz K, Marusiak J i wsp: Tensegrity principle in massage demonstrated by electro- and mechanomyography; *J Bodyw Mov Ther* 2009;13(2):164-70.
205. Kassolik K, Andrzejewski W: Tensegration massage; *Physiotherapy* 2010;18(1): 67-72.
206. Piron A: The tensegrity concept applied to the laryngeal biodynamics; *Rev Laryngol Otol Rhinol* 2007;128(5):273-8.
207. Stecco C, Stern R, Porzionato A, et al. Hyaluronan within fascia in the etiology of myofascial pain. *Surg Radiol Anat* 2011; 33:891-6.
208. Langevin HM, Fox JR, Koptiuch C, et al. Reduced thoracolumbar fascia shear strain in human chronic low back pain. *BMC Musculoskelet Disord* 2011; 12:203.
209. Ajimsha MS, Binsu D, Chithra S. Effectiveness of Myofascial release in the management of chronic low back pain in nursing professionals. *J Bodyw Mov Ther* 2014; 18:273-81.
210. Stecco A, Gilliar W, Hill R, et al. The anatomical and functional relation between gluteus maximus and fascia lata. *J Bodyw Mov Ther* 2013; 17:512-7.
211. Stecco C, Porzionato A, Macchi V, et al. The Expansions of the Pectoral Girdle Muscles onto the Brachial Fascia: Morphological Aspects and Spatial Disposition. *Cells Tissues Organs* 2007;188: 320-9.
212. Casato G, Stecco C, Busin R, Role of fasciae in nonspecific low back pain, *Eur J Transl Myol*. 2019 Aug 6;29(3):8330.
213. Branchini M1, Lopopolo F2, Andreoli E3, Loreti I4, Marchand AM5, Stecco A, Fascial Manipulation® for chronic specific low back pain: a single blinded randomized controlled trial, Version 2. *F1000Res*. 2015 Nov 3 [revised 2016 Jan 8]; 4:1208.
214. Enwemeka CS., Bonet IM., Ingle JA., Prudhithumrong S., Ogbahon FE., Gbenedio NA.: Postural Correction in Persons with Neck Pain. II. Integrated Electromyography of the Upper Trapezius in Three Simulated Neck Positions. *J Orthop Sport Phys Ther*. 2013; 11, 240-242.
215. Sugimoto D., Myer GD., Foss KDB., Hewett TE.: Specific exercise effects of preventive neuromuscular training intervention on anterior cruciate ligament injury risk reduction in young females: Meta-analysis and subgroup analysis. *British Journal of Sports Medicine*. 2015, Mar;49(5):282-9.
216. Hewett TE., Myer GD., Ford KR., Heidt RS., Colosimo AJ., McLean SG., et al.: Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: A prospective study. *Am J Sports Med*. 2005, Apr;33(4):492-501.
217. Pardehshenas H, Maroufi N., Sanjari MA., Pamiampour M., Levin SM.: Lumbopelvic muscle activation patterns in three stances under graded loading conditions: Proposing a tensegrity model for load transfer through the sacroiliac joints. *J Bodyw Mov Ther*. 2014, Oct;18(4):633-42.
218. Rugh JD., Drago CJ.: Vertical dimension: A study of clinical rest position and jaw muscle activity. *J Prosthet Dent*. 1981, Jun, 45(6):670-5.
219. Wahlund K.: Temporomandibular disorders in adolescents. Epidemiological and methodological studies and a randomized controlled trial. *Swed Dent J Suppl*. 2003;(164), 2-64.
220. Ställ AT.: The Philosophy and Mechanical Principle of Osteopathy. Hudson-Kimberly Pub Co. 1902 - 319.
221. Fukunaga T., Kawakami Y., Kubo K., Kanehisa H.: Muscle and Tendon Interaction During Human Movements. *Exerc Sport Sci Rev*. 2002 Jul;30(3):106-10.
222. Tak I., Glasgow P., Langhout R., Weir A., Kerkhoffs G., Agricola R.: Hip Range of Motion Is Lower in Professional Soccer Players with Hip and Groin Symptoms or Previous Injuries, Independent of Cam Deformities. *Am J Sports Med*. 2016; Mar;44(3):682-8.
223. Weiss HR., Negrini S., Rigo M., Kotwicki T., Hawes MC., Grivas TB., et al.: Indications for conservative management of scoliosis (guidelines). *Scoliosis*. 2006 May 8; 1:5. doi: 10.1186/1748-7161-1-5.
224. Canavese F., Kaelin A.: Adolescent idiopathic scoliosis: Indications and efficacy of nonoperative treatment. *Indian J Orthop.*, 2011 Jan-Mar; 45(1): 7–14.
225. Lambek LC, van Mechelen W, Knol DL, Loisel P, Anema JR. Randomised controlled trial of integrated care to reduce disability from chronic low back pain in working and private life. *BMJ* 2010;340: d750.
226. Pellisé F, Balagué F, Rajmil L, Cedraschi C, Aguirre M, Fontecha CG, Pasarín M, Ferrer M., Prevalence of low back pain and its effect on health-related quality of life in adolescents., *Arch Pediatr Adolesc Med*. 2009 Jan;163(1):65-71.
227. Hodges PW, Smeets RJ. Interaction Between Pain, Movement, and Physical Activity. *Clin J Pain*. 2015; 31:97–10.
228. Sahrman SA.: Diagnosis by the physical therapist - A prerequisite for treatment. A special communication. *Physical Therapy*. 1988, 68, 11, 1703-1706.
229. Cole TJ., Bellizzi MC., Flegal KM., Dietz WH.: Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ*. 2000; 320:1240.
230. Jeffries L, Milanese S, Grimmer-Somers K; Epidemiology of Adolescent Spinal Pain: A Systematic Overview of the Research Literature, 2007. *Spine*. 32(23):2630-2637.
231. Canavese F., Kaelin A.: Adolescent idiopathic scoliosis: Indications and efficacy of nonoperative treatment. *Indian J Orthop.*, 2011 Jan-Mar; 45(1): 7–14.
232. Hewett TE., Myer GD., Ford KR., Heidt RS., Colosimo AJ., McLean SG., et al.: Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: A prospective study. *Am J Sports Med*. 2005, Apr;33(4):492-501.
233. Sugimoto D., Myer GD., Foss KDB., Hewett TE.: Specific exercise effects of preventive neuromuscular training intervention on anterior cruciate ligament injury risk reduction in young females: Meta-analysis and subgroup analysis. *British Journal of Sports Medicine*. 2015, Mar;49(5):282-9.
234. Strojek K., Bulatowicz I., Radziwińska A., Kaźmierczak U., Siedlaczek M., Lipiec M., Dzierżanowski M ŻW.: Assessment of body posture in children of Kindergarten Age. *Heal Sci*. 2014; 4:229–40.
235. Mieszowska M., Magdalena K., Garbin M., Srokowski G., Tomczyk M., Piśkorska E.: Characteristics of body posture of Bydgoszcz region pre-school children. *J Educ Heal Sport*. 2016;6 (8): 637-652.
236. Cole TJ., Bellizzi MC., Flegal KM., Dietz WH.: Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ*. 2000; 320:1240.
237. Dionne CE, Dunn KM, Croft PR. Does back pain prevalence really decrease with increasing age? A systematic review. *Age Ageing* 2006; 35: 229– 34.
238. Coenen P, Smith A, Paananen M, O'Sullivan P, Beales D, Straker L. Trajectories of low back pain from adolescence to young adulthood. *Arthritis Care Res (Hoboken)*. 2017; 69:403–412.
239. Janiszewska R., Tuzinek S., Nowak S., Ratyńska A., Biniaszewski T.: Abnormalities of posture in 6-12 year-old children – pupils of primary schools from Radom – a pilot study. *Probl Hig Epidemiol*. 2009;90(3):342–346.
240. Dekker J., van Baar ME., Curfs EC., Kerssens JJ.: Diagnosis and Treatment in Physical Therapy: An Investigation of Their Relationship. *Phys Ther.*, 1993, Sep;73(9):568-77.
241. Gauer RL., Semidey MJ.: Diagnosis and treatment of temporomandibular disorders. *Am Fam Physician*. 2015;91(6):378–86.
242. Blagrove P.: Diagnostik und Behandlung der Art. Temporomandibularis. 2002, 1: 24–29.
243. Romero-Reyes M., Uyanik JM.: Orofacial pain management: Current perspectives. *Journal of Pain Research*. 2014, Feb 21;7:99-115.
244. McNeely ML., Armijo OS., Magee DJ.: A systematic review of the effectiveness of physical therapy interventions for temporomandibular disorders. *Phys Ther*. 2006; May;86(5):710-25.
245. Armijo OS., Pitance L., Singh V., Neto F., Thie N., Michelotti A.: Effectiveness of Manual Therapy and Therapeutic Exercise for Temporomandibular Disorders: Systematic Review and Meta-Analysis. *Phys Ther.*, 2016, Jun, 96 (1):9-25.
246. Saddu SC, Dyasanoor S, Valappila NJ, Ravi BV: The evaluation of head and craniocervical posture among patients with and without temporomandibular joint disorders- a comparative study; *J Clin Diagn Res* 2015;9(8): ZC55-8.
247. Nicolakis P, Nicolakis M, Piehsinger E, Ebenbichler G i wsp.: Relationship between craniomandibular disorders and poor posture; *Cranio* 2000;18(2):106-12.
248. Huggare JA, Raustia AM: Head posture and cervicovertebral and craniofacial morphology in patients with craniomandibular dysfunction; *Cranio* 1992;10(3):173-179.
249. Lee WY, Okeson JP, Lindroth J: The relationship between forward head posture and temporomandibular disorders; *J Orofac Pain* 1995;9(2):161-7.
250. Weber P, Corrêa ECR, Ferreira FS, Soares JC i wsp.: Cervical spine dysfunction signs and symptoms in individuals with temporomandibular disorder; *J Soc Bras Fonoaudiol* 2012;24(2):134-9.
251. Bragatto MM, Bevilacqua-Grossi D, Regalo SC, Sousa JD i wsp.: Associations among temporomandibular disorders, chronic neck pain and neck pain disability in computer office workers: a pilot study; *J Oral Rehabil* 2016;43(5):321-32.
252. Myers Thomas W. "Anatomy Trains" 2009 ISBN 978-443-10283-7.
253. Tanamas SK, Wluka AE, Berry P et al. Relationship between obesity and foot pain and its association with fat mass, fat distribution, and muscle mass. *Arthritis Care Res* 2012; 64:262-268.
254. Stumieks DL, George R, Lord SR. Balance disorders in the elderly. *Neurophysiol Clin* 2008; 38:467–478.
255. Spink MJ, Fotooha MR. Foot and Ankle Strength, Range of Motion, Posture, and Deformity Are Associated With Balance and Functional Ability in Older Adults. *Arch Phys Med Rehabil* 2011; 92:68-75.
256. Drzał-Grabiec J, Rachwał M, Trzaskoma Z i wsp. The foot deformity versus postural control in females aged over 65 years. *Acta Bioeng Biomech* 2014;16(4):73-80.
257. Hals E. K. B. and Stubhaug A., "Mental and somatic comorbidities in chronic orofacial pain conditions: pain patients in need of multiprofessional team approach," *Scandinavian Journal of Pain*, vol. 14, no. 4, pp. 153-154, 2011.
258. Castillo-Morales R: Die orofaziale Regulationstherapie. Kommunikation; Pflaum Verlag. München 1992.
259. Mehta NR, Forgione AG, Rosenbaum RS, Holmberg R: TMJ triad of dysfunctions: a biologic basis of diagnosis and treatment; *J Mass Dent Soc* 1984;33(4):173-6,212-3.

260. Gonzalez HE, Manns A: Forward head posture: its structural and functional influence on the stomatognathic system, a conceptual study; *Cranio* 1996;14(1):71-80.
261. Grossi DB: Physiotherapeutic treatment for temporomandibular disorders (tmd); *Braz J Oral Sci* 2004;3(10):492-7.
262. Ohrbach R, Fillingim R. B., Mulkey F et al.: Clinical findings and pain symptoms as potential risk factors for chronic TMD: descriptive data and empirically identified domains from the OPPERA case-control study; *Journal of Pain*, vol. 14, no. 11, pp. T27-T45, 2011.
263. Herb K., Cho S., Stiles MA.: Temporomandibular joint pain and dysfunction. *Current Pain and Headache Reports*. 2006, Dec;10(6):408-14.
264. Alomar X, Medrano J, Cabratosa J, Clavero JA, Lorente M, Serra I, Monill JM, Salvador A. Anatomy of the temporomandibular joint. *Semin Ultrasound CT MR*. 2007; 28(3):170-83.
265. Al-Ani Z., Gray R.J., Davies S.J., Sloan P., Glenn A-M. "Stabilization splint therapy for the treatment of temporomandibular myofascial pain" *J Dent Educ*. 2005; 69(11):1242-50.
266. Emshoff R. "Clinical factors affecting the outcome of occlusal splint therapy of temporomandibular joint disorders" *J Oral Rehabil*. 2006;33(6):393-401.
267. Matida Hamata M., Junqueira Zuim P.R., Garcia A.R.; Comparative evaluation of the efficacy of occlusal splints fabricated in centric relation or maximum intercuspation in temporomandibular disorders patients, *J Appl Oral Sci*. 2009; 17(1):32-8.
268. Stiesch-Scholz M., Kempert J., Wolter S., Tschernitschek H., Rossbach A. "Comparative prospective study on splint therapy of anterior disc displacement without reduction" *J Oral Rehabil*. 2005; 32(7):474-479.
269. Alencar F. Jr, Becker A. "Evaluation of different occlusal splints and counselling in the management of myofascial pain dysfunction" *J Oral Rehabil*. 2009; 36(2):79-85.
270. Al-Ani MZ, Davies SJ, Gray RJ, Sloan P, Glenn AM.; "Stabilisation splint therapy for temporomandibular pain dysfunction syndrome." *Cochrane Database Syst Rev*. 2004; (1):CD002778.
271. Zakrzewska JM.: Orofacial pain management current perspectives. *J Pain Res*. 2014; 7: 99-115.
272. Furnish T.: Temporomandibular joint disorders. In: *Case Studies in Pain Management*. (pp. 333-340). Cambridge, 2014.
273. Nobrega J. C., Tesseroli de Siquera S. R., and Tesseroli de Siquera J. T., Differential diagnosis in atypical pain: a Clinical study, *Arquivos de Neuro-Psiquiatria*, vol. 65, no. 2, pp. 256-261, 2007.
274. Janal M. N., Raphael K. G., Nayak S., and Klausner J., Prevalence of myofascial temporomandibular disorder in US community women, *Journal of Oral Rehabilitation*, vol. 14, no. 11, pp. 801-809, 2008.
275. Pedroni CR., De Oliveira AS., Guaratini MI.: Prevalence study of signs and symptoms of temporomandibular disorders in university students. *J Oral Rehabil*. 2003, Mar;30(3):283-289.
276. Tassorelli C., Tramontano M., Berlangieri M. et al., Assessing and treating primary headaches and cranio-facial pain in patients undergoing rehabilitation for neurological diseases, *Journal of Headache and Pain*, vol. 18, no. 1, p. 99, 2017.
277. Karch D., Groß-Selbeck G., Pietz J., Schlack H-G.: Orofaziale Regulationstherapie nach Castillo Morales. *Monatsschrift Kinderheilkd*. 2005; Mar. 782-785.
278. Feteih RM. Signs and symptoms of temporomandibular disorders and oral parafunctions in urban Saudi Arabian adolescents: a research report. *Head & Face Med*. 2006; 2:25 doi:10.1186/1746-160X-2-25.
279. Kijak E., Lietz-Kijak D., Sliwinski Z., Fraczak B.: Muscle activity in the course of rehabilitation of masticatory motor system functional disorders, *Advances in Hygiene and Experimental Medicine*, vol. 67, pp. 507-516, 2013.
280. Weiss HR., Negrini S., Rigo M., Kotwicki T., Hawes MC., Grivas TB., et al.: Indications for conservative management of scoliosis (guidelines). *Scoliosis*. 2006 May 8; 1:5. doi: 10.1186/1748-7161-1-5.
281. Arnett GW, Milam SB, Gottesman L. Progressive mandibular retrusion - idiopathic condylar resorption. Part I. *Am J Orthod Dentofacial Orthop*. 1996a; 110:8-15.
282. Arnett GW, Milam SB, Gottesman L. Progressive mandibular retrusion - idiopathic condylar resorption. Part II. *Am J Orthod Dentofacial Orthop*. 1996b; 110:117-127.
283. Leitfaden CF.: Osteopathie. Leitfaden Osteopathie. 2016.
284. Chaitow L.: Cranial Manipulation: Theory and Practice: Osseous and Soft Tissue Approaches. 2005.
285. Matheus RA, Ramos-Perez FM, Menezes AV, Ambrosano GM i wsp.: The relationship between temporomandibular dysfunction and head and cervical posture; *J Appl Oral Sci* 2009;17(3):204-8.
286. Ayub E: Glasheen-Wray M: Kraus S.: Head Posture: A Case Study of the Effects on the Rest Position of the Mandible. *J Orthop Sport Phys Ther*. 1984, Vol.:5, 4, 179-183.
287. Thompson JR: Brodie AG.: Factors in the Position of the Mandible. *J Am Dent Assoc.*, 2014;4(3):68-77.
288. Preiskel HW.: Some observations on the postural position of the mandible. *J Prosthet Dent*. 1965, Jul-Aug;15:625-33.
289. Harris HL.: Effect of Loss of Vertical Dimension on Anatomic Structures of the Head and Neck. *J Am Dent Assoc Dent Cosm*. 2015;
290. Fuentes FR., Freesmeyer W., Henríquez P J.: Influencia de la postura corporal en la prevalencia de las disfunciones craneomandibulares. *Rev Med Chil*. 1999, vol. 127, 9, 1079-1085.
291. Maeda N, Sakaguchi K, Mehta NR, Abdallah EG i wsp.: Effects of experimental leg length discrepancies on body posture and dental occlusion. *Cranio - j craniomandib pract*; *Cranio* 2011;29(3):194-203.
292. Bonato LL, Quinelato V, de Felipe Cordeiro PC i wsp.: Association between temporomandibular disorders and pain in other regions of the body; *J Oral Rehabil* 2017;44(1):9-15.
293. Souza JA, Pasinato F, Corrêa EC, da Silva AM: Global body posture and plantar pressure distribution in individuals with and without temporomandibular disorder: a preliminary study; *J Manipulative Physiol Ther* 2014;37(6):407-14.
294. Saito ET, Akashi PM, Sacco Ide C: Global body posture evaluation in patients with temporomandibular joint disorder; *Clinics (Sao Paulo)* 2009;64(1):35-9.
295. Valentino B, Valentino T, Melito F: Correlation between interdental occlusal plane and plantar arches; *Bull Group Int Rech Sci Stomatol Odontol* 2002;44(1):10-3.
296. Visscher CM., Lobbezoo F., De Boer W., Van Der Zaag J., Naeije M.: Prevalence of cervical spinal pain in craniomandibular pain patients. *Eur J Oral Sci*. 2001, Apr;109(2):76-80.
297. Armijo-Olivo S., Jara X., Castillo N., Alfonso L., Schilling A., Valenzuela E., et al.: A comparison of the head and cervical posture between the self-balanced position and the Frankfurt method. *J Oral Rehabil*. 2006, Mar;33(3):194-201.
298. Hackney J., Bade D., Clawson A.: Relationship between forward head posture and diagnosed internal derangement of the temporomandibular joint. *J Orofac Pain*. 1993;
299. Sonnesen L., Bakke M., Solow B.: Temporomandibular disorders in relation to craniofacial dimensions, head posture and bite force in children selected for orthodontic treatment. *Eur J Orthod*. 2001, Apr;23(2):179-92.
300. McNeill C., Mohl ND., Rugh JD., Tanaka TT.: Temporomandibular Disorders: Diagnosis, Management, Education, and Research. *J Am Dent Assoc*. 2015, Mar;120(3):253-257.
301. Johnson GM.: The correlation between surface measurement of head and neck posture and the anatomic position of the upper cervical vertebrae. *Spine (Phila Pa 1976)*. 1998, Apr. 15;23(8):921-927.
302. Saddu SC, Dyananoor S, Valappila NJ, Ravi BV: The evaluation of head and craniocervical posture among patients with and without temporomandibular joint disorders- A comparative study. *J Clin Diagnostic Res*. 2015.
303. De Wijer A., Steenkens MH., De Leeuw JR.J., Bosman F., Helden P.J.M.: Symptoms of the cervical spine in temporomandibular and Cervical Spine Disorders. *J Oral Rehabil*. 1996, Nov;23(11):742-750.
304. Lee B-K.: Influence of proprioceptive neuromuscular facilitation therapeutic exercise on woman with temporomandibular joint disorder: a case study. *J Exerc Rehabil*. 2018, Dec 27;14(6):1074-1079.
305. Walczyńska-Dragon K, Baron S, Nitecka-Buchta A, Tkacz E. Correlation between TMD and cervical spine pain and mobility: Is the whole body balance TMJ related? *Biomed Res Int*. 2014.
306. Myers TW.: Anatomy Trains: Myofascial Meridians for Manual and Movement Therapists. DB Publishing; 2019.
307. Mannheimer JS., Rosenthal RM.: Acute and chronic postural abnormalities as related to craniofacial pain and temporomandibular disorders. *Dental clinics of North America*. 1991, 35(1):185-208.
308. Dewar L. *The Myofascial Release Manual*. Vol. 87, Physiotherapy. 2005. 330 p.
309. Barnes MF.: The basic science of myofascial release: morphologic change in connective tissue. *J Bodyw Mov Ther*. 1997, 1(4), 231-238.a
310. Barnes JF.: Myofascial release for craniomandibular pain and dysfunction. *Int J Orofacial Myology [Internet]*. 1997, 1(4), 231-238.b
311. Fernández De Las Peñas C., Palomeque Del Cerro L., Fernández Camero J.: Manual treatment of post-whiplash injury. *J Bodyw Mov Ther*. 2005.
312. LeBauer A., Braitlik R., Stowe K.: The effect of myofascial release (MFR) on an adult with idiopathic scoliosis. *J Bodyw Mov Ther*. 2008, Oct;12(4):356-363.
313. Thumati P., Kerstein R., Thumati R.: Disclusion time reduction therapy in treating occluso-muscular pains. *J Indian Prosthodont Soc*. 2016, Jan-Mar; 17(1): 95-98.
314. Qadeer S., Abbas AA., Sarinaphakom L., Kerstein RB.: Comparison of excursive occlusal force parameters in post-orthodontic and non-orthodontic subjects using T-Scan® III. *Cranio - J Craniomandib Pract*. 2018, Jan;36(1):11-18.
315. Friction J.: Myofascial Pain: Mechanisms to Management. *Oral and Maxillofacial Surgery Clinics of North America*. 2016, Aug;28(3):289-311.
316. Darlow LA., Pesco J GM.: The relationship of posture to myofascial pain dysfunction syndrome. *J Am Dent Assoc*. 1987;114(1):73-5.
317. Zonnenberg A.J.J., Van Maanen C.J., Oostendorp R.A.B., Elvers J.W.H.: Body posture photographs as a diagnostic aid for musculoskeletal disorders related to temporomandibular disorders (TMD). *Cranio*. 1996, Jan; 64(1): 35-39.
318. Wright EF., Domenech MA., Fischer JR.: Usefulness of posture training for patients with temporomandibular disorders. *J Am Dent Assoc*. 2000, May;131(5):564- 568.
319. Siqueira III WCMPMJJTT de. Radiographic evaluation of cervical spine of subjects with temporomandibular joint internal disorder. *Braz Oral Res*. 2004;14(4).
320. Dischiavi SL., Wright AA., Hegedus E.J., Bleakley CM.: Biotensegrity and myofascial chains: A global approach to an integrated kinetic chain. *Med Hypotheses*. 2018, Jan;110:90-96.
321. Saito ET, Akashi PMH., de Camargo Neves Sacco I.: Global Body Posture Evaluation in Patients with Temporomandibular Joint Disorder. *Clinics (Sao Paulo)*. 2009, 64(1):35-39.
322. Park Y., Bae Y.: Change of Range of Motion of the Temporomandibular Joint after Correction of Mild Scoliosis. *J Phys Ther Sci*. 2014, Aug; 26(8): 1157-1160.
323. Ikemitsu H., Izumi RZY.: The relationship between jaw deformity and scoliosis. *Oral Radiol*. 2006;22(1):14-17.
324. McClean L.F., et al.: Effects of changing body position on dental occlusion. *J Dent Res*. 1973;52(5):1041-1045.
325. Zyznawska MK.: Assessment of temporomandibular joint dysfunctions and the position of pMk. *J Orthop Trauma Surg*. 2013 (12):42-8.
326. Rosenbloom L.: The Chailey approach to postural management. *Arch Dis Child*. 2002, 44:158-163.
327. Thomas GD., Segal SS.: Neural control of muscle blood flow during exercise. *J Appl Physiol*. 2004, Aug;97(2):731-738.
328. Roe SM., Johnson CD., Tansley EA.: Investigation of physiological properties of nerves and muscles using electromyography. *Adv Physiol Educ*. 2014, Dec;38(4):348-354.
329. Munhoz WC., Marques AP., Siqueira JT.: Radiographic evaluation of cervical spine of subjects with temporomandibular joint internal disorder. *Braz Oral Res*. 2004, Oct-Dec;18(4):283-289.
330. Fuentes FR, Freesmeyer W, Henríquez P J. Influencia de la postura corporal en la prevalencia de las disfunciones craneomandibulares. *Rev Med Chil*. 1999